

CHAPTER 2

INVERSE TRIGONOMETRIC FUNCTIONS

VERY SHORT ANSWER TYPE QUESTIONS (1 MARK)

1. Write the principal value of

(i) $\sin^{-1}(-\sqrt{3}/2)$

(ii) $\cos^{-1}(\sqrt{3}/2)$.

(iii) $\tan^{-1}\left(-\frac{1}{\sqrt{3}}\right)$

(iv) $\operatorname{cosec}^{-1}(-2)$.

(v) $\cot^{-1}\left(\frac{1}{\sqrt{3}}\right)$.

(vi) $\sec^{-1}(-2)$.

(vii) $\sin^{-1}\left(\frac{-\sqrt{3}}{2}\right) + \cos^{-1}\left(\frac{-1}{2}\right) + \tan^{-1}(-1/\sqrt{3})$

2. What is value of the following functions (using principal value).

(i) $\tan^{-1}\left(\frac{1}{\sqrt{3}}\right) - \sec^{-1}\left(\frac{2}{\sqrt{3}}\right)$.

(ii) $\sin^{-1}\left(-\frac{1}{2}\right) - \cos^{-1}\left(\frac{\sqrt{3}}{2}\right)$.

(iii) $\tan^{-1}(1) - \cot^{-1}(-1)$.

(iv) $\operatorname{cosec}^{-1}(\sqrt{2}) + \sec^{-1}(\sqrt{2})$.

(v) $\tan^{-1}(1) + \cot^{-1}(1) + \sin^{-1}(1)$.

(vi) $\sin^{-1}\left(\sin \frac{4\pi}{5}\right)$.

(vii) $\tan^{-1}\left(\tan \frac{5\pi}{6}\right)$.

(viii) $\operatorname{cosec}^{-1}\left(\operatorname{cosec} \frac{3\pi}{4}\right)$.

SHORT ANSWER TYPE QUESTIONS (4 MARKS)

3. Show that $\tan^{-1}\left(\frac{\sqrt{1+\cos x} + \sqrt{1-\cos x}}{\sqrt{1+\cos x} - \sqrt{1-\cos x}}\right) = \frac{\pi}{4} + \frac{x}{2}$. $x \in [0, \pi]$



4. Prove

$$\tan^{-1}\left(\frac{\cos x}{1 - \sin x}\right) - \cot^{-1}\left(\sqrt{\frac{1 + \cos x}{1 - \cos x}}\right) = \frac{\pi}{4} \quad x \in (0, \pi/2).$$

5. Prove $\tan^{-1}\left(\frac{x}{\sqrt{a^2 - x^2}}\right) = \sin^{-1}\frac{x}{a} = \cos^{-1}\left(\frac{\sqrt{a^2 - x^2}}{a}\right).$

6. Prove

$$\cot^{-1}\left[2 \tan\left(\cos^{-1}\frac{8}{17}\right)\right] + \tan^{-1}\left[2 \tan\left(\sin^{-1}\frac{8}{17}\right)\right] = \tan^{-1}\left(\frac{300}{161}\right).$$

7. Prove $\tan^{-1}\left(\frac{\sqrt{1+x^2} + \sqrt{1-x^2}}{\sqrt{1+x^2} - \sqrt{1-x^2}}\right) = \frac{\pi}{4} + \frac{1}{2}\cos^{-1}x^2.$

8. Solve $\cot^{-1} 2x + \cot^{-1} 3x = \frac{\pi}{4}.$

9. Prove that $\tan^{-1}\left(\frac{m}{n}\right) - \tan^{-1}\left(\frac{m-n}{m+n}\right) = \frac{\pi}{4}, m, n > 0$

10. Prove that $\tan\left[\frac{1}{2}\sin^{-1}\left(\frac{2x}{1+x^2}\right) + \frac{1}{2}\cos^{-1}\left(\frac{1-y^2}{1+y^2}\right)\right] = \frac{x+y}{1-xy}$

11. Solve for x , $\cos^{-1}\left(\frac{x^2-1}{x^2+1}\right) + \frac{1}{2}\tan^{-1}\left(\frac{-2x}{1-x^2}\right) = \frac{2\pi}{3}$

12. Prove that $\tan^{-1}\frac{1}{3} + \tan^{-1}\frac{1}{5} + \tan^{-1}\frac{1}{7} + \tan^{-1}\frac{1}{8} = \frac{\pi}{4}$

13. Solve for x , $\tan(\cos^{-1}x) = \sin(\tan^{-1}2); x > 0$

14. Prove that $2\tan^{-1}\left(\frac{1}{5}\right) + \tan^{-1}\left(\frac{1}{4}\right) = \tan^{-1}\left(\frac{32}{43}\right)$



15. Evaluate $\tan\left[\frac{1}{2}\cos^{-1}\left(\frac{3}{\sqrt{11}}\right)\right]$
16. Prove that $\tan^{-1}\left(\frac{a\cos x - b\sin x}{b\cos x + a\sin x}\right) = \tan^{-1}\left(\frac{a}{b}\right) - x$
17. Prove that $\cot\left\{\tan^{-1}x + \tan^{-1}\left(\frac{1}{x}\right)\right\} + \cos^{-1}(1 - 2x^2) + \cos^{-1}(2x^2 - 1) = \pi, x > 0$
18. Prove that $\tan^{-1}\left(\frac{a-b}{1+ab}\right) + \tan^{-1}\left(\frac{b-c}{1+bc}\right) + \tan^{-1}\left(\frac{c-a}{1+ca}\right) = 0$ where $a, b, c > 0$
19. Solve for x , $2 \tan^{-1}(\cos x) = \tan^{-1}(2 \operatorname{cosec} x)$
20. Express $\sin^{-1}(x\sqrt{1-x} - \sqrt{x}\sqrt{1-x^2})$ in simplest form.
21. If $\tan^{-1}a + \tan^{-1}b + \tan^{-1}c = \pi$, then prove that $a + b + c = abc$
22. If $\sin^{-1}x > \cos^{-1}x$, then x belongs to which interval?

ANSWERS

1. (i) $-\frac{\pi}{3}$ (ii) $\frac{\pi}{6}$ (iii) $\frac{-\pi}{6}$ (iv) $\frac{-\pi}{6}$
 (v) $\frac{\pi}{3}$ (vi) $\frac{2\pi}{3}$ (vii) $\frac{\pi}{6}$.
2. (i) 0 (ii) $\frac{-\pi}{3}$ (iii) $-\frac{\pi}{2}$ (iv) $\frac{\pi}{2}$
 (v) π (vi) $\frac{\pi}{5}$ (vii) $\frac{-\pi}{6}$ (viii) $\frac{\pi}{4}$.



8. 1

11. $\tan \frac{\pi}{12} = 2 - \sqrt{3}$

13. $\frac{\sqrt{5}}{3}$

15. $\sqrt{\frac{\sqrt{11}-3}{3+\sqrt{11}}}$

19. $x = \frac{\pi}{4}.$

20. $\sin^{-1} x - \sin^{-1} \sqrt{x}.$

22. $\left(\frac{1}{\sqrt{2}}, 1 \right]$

21. **Hint:** Let $\tan^{-1} a = \alpha$
 $\tan^{-1} b = \beta$
 $\tan^{-1} c = \gamma$

then given, $\alpha + \beta + \gamma = \pi$

$\therefore \alpha + \beta = \pi - \gamma$

take tangent on both sides,

$\tan (\alpha + \beta) = \tan (\pi - \gamma)$

