

CBSE Class 11 Mathematics

Important Questions

Chapter 13

Limits and Derivatives

1 Marks Questions

1. Evaluate $\lim_{x \rightarrow 3} \left[\frac{x^2 - 9}{x - 3} \right]$

Ans. $\lim_{x \rightarrow 3} \frac{x^2 - 9}{x - 3} \frac{0}{0} \text{ form}$

$$\lim_{x \rightarrow 3} \frac{(x+3)(\cancel{x-3})}{(\cancel{x-3})} = 3+3 = 6$$

2. Evaluate $\lim_{x \rightarrow 0} \frac{\sin 3x}{5x}$

Ans. $\lim_{x \rightarrow 0} \frac{\sin 3x}{5x}$

$$= \lim_{3x \rightarrow 0} \frac{\sin 3x}{3x} \times \frac{3}{5}$$

$$= 1 \times \frac{3}{5} = \frac{3}{5} \left[\because \lim_{x \rightarrow 0} \frac{\sin x}{x} = 1 \right]$$

3. Find derivative of 2^x

Ans. Let $y = 2^x$

$$\frac{dy}{dx} = \frac{d}{dx} 2 = 2^x \log 2$$

4. Find derivative of $\sqrt{\sin 2x}$

$$\begin{aligned}\text{Ans. } \frac{d}{dx} \sqrt{\sin 2x} &= \frac{1}{2\sqrt{\sin 2x}} \frac{d}{dx} \sin 2x \\&= \frac{1}{2\sqrt{\sin 2x}} \times 2 \cos 2x \\&= \frac{\cos 2x}{\sqrt{\sin 2x}}\end{aligned}$$

5. Evaluate $\lim_{x \rightarrow 0} \frac{\sin^2 4x}{x^2}$

Ans.

$$\begin{aligned}\lim_{x \rightarrow 0} \frac{\sin^2 4x}{x^2 4^2} \times 4^2 &= \lim_{4x \rightarrow 0} \left(\frac{\sin 4x}{4x} \right)^2 \times 16 \\&= 1 \times 16 = 16\end{aligned}$$

6. What is the value of $\lim_{x \rightarrow a} \left(\frac{x^n - a^n}{x - a} \right)$

$$\text{Ans. } \lim_{x \rightarrow a} \frac{x^n - a^n}{x - a} = 1$$

7. Differentiate $\frac{2^x}{x}$

$$\begin{aligned}\text{Ans. } \frac{d}{dx} \frac{2^x}{x} &= \frac{x \frac{d}{dx} 2^x - 2^x \frac{d}{dx} x}{x^2} \\&= \frac{x \times 2^x \log 2 - 2^x \times 1}{x^2}\end{aligned}$$

$$= 2x \frac{[x+10g^2-1]}{x^2}$$

8. If $y = e^{\sin x}$ find $\frac{dy}{dx}$

Ans. $y = e^{\sin x}$

$$\frac{dy}{dx} = \frac{d}{dx} e^{\sin x}$$

$$= e^{\sin x} \times \cos x = \cos x e^{\sin x}$$

9. Evaluate $\lim_{x \rightarrow 1} \frac{x^{15} - 1}{x^{10} - 1}$

Ans. $\lim_{x \rightarrow 1} \frac{x^{15} - 1}{x^{10} - 1}$

$$= \frac{\lim_{x \rightarrow 1} \frac{x^{15} - 1^{15}}{x - 1}}{\lim_{x \rightarrow 1} \frac{x^{10} - 1^{10}}{x - 1}} = \frac{15 \times 1^{14}}{10 \times 1^9}$$

$$= \frac{15}{10} = \frac{3}{2}$$

10. Differentiate $x \sin x$ with respect to x

Ans. $\frac{d}{dx} x \sin x = x \cos x + \sin x \cdot 1$

$$= x \cos x + \sin x$$

11. Evaluate $\lim_{x \rightarrow 1} \frac{x^2 + 1}{x + 100}$

Ans. $\lim_{x \rightarrow 1} \frac{x^2 + 1}{x + 100} = \frac{2}{101}$

12. Evaluate $\lim_{x \rightarrow 0} [\operatorname{cosec} x - \cot x]$

Ans. $\lim_{x \rightarrow 0} [\operatorname{cosec} x - \cot x]$

$$= \lim_{x \rightarrow 0} \left[\frac{1}{\sin x} - \frac{\cos x}{\sin x} \right]$$

$$= \lim_{x \rightarrow 0} \frac{1 - \cos x}{\sin x} = \lim_{x \rightarrow 0} \frac{\cancel{2} \sin^2 \frac{x}{2}}{\cancel{2} \sin \frac{x}{2} \cos \frac{x}{2}}$$

$$= \lim_{x \rightarrow 0} \tan \frac{x}{2} = 0$$

13. Find $f^{-1}(x)$ at $x = 100$

if $f(x) = 99x$

Ans. $f(x) = 99x$

$$f^{-1}(x) = 99 \quad \text{at } x = 100$$

$$f^{-1}(x) = 99$$

14. Evaluate $\lim_{x \rightarrow -2} \frac{\tan \pi x}{x + 2}$

Ans. $\lim_{x \rightarrow -2} \frac{\tan \pi x}{x+2} \quad \frac{1}{0} \text{ form}$

let $x+2 = y$

$x = y - 2$

$$\lim_{y \rightarrow 0} \frac{\tan \pi(y-2)}{y}$$

$$\lim_{y \rightarrow 0} \frac{-\tan \pi(2-y)}{y} = \lim_{y \rightarrow 0} \frac{\tan[2\pi - 2y]}{y}$$

$$= \lim_{2y \rightarrow 0} \frac{+\tan 2y}{2y} \times 2$$

$$= 1 \times 2 = 2$$

15. Find derivative of $\sin^n x$

Ans. $\frac{d}{dx} \sin^n x$

$$= n \sin^{n-1} x \cdot \frac{d}{dx} \sin x$$

$$= n \sin^{n-1} x \cos x$$

16. Find derivative of $1 + x + x^2 + x^3 + \dots + x^{50}$ at $x = 1$

Ans. $f(x) = 1 + x + x^2 + x^3 + \dots + x^{50}$

$$f^1(x) = 1 + 2x + 3x^2 + \dots + 50x^{49}$$

at $x = 1$

$$f^1(1) = 1 + 2 + 3 + \dots + 50 = \frac{50(50+1)}{2}$$

$$= 25 \times 51 = 1275$$

17. The value of $\lim_{h \rightarrow 0} \frac{e^{2h} - 1}{h}$

Ans. $\lim_{2h \rightarrow 0} \frac{e^{2h} - 1}{2h} \times 2$

$$= 1 \times 2 = 2$$

18. Evaluate $\lim_{x \rightarrow 0} \frac{(1+x)^6 - 1}{(1+x)^2 - 1}$

Ans. $\lim_{x \rightarrow 0} \left[\frac{(1+x)^6 - 1}{(1+x)^2 - 1} \right]$

let $1+x = y$

$x \rightarrow 0, y \rightarrow 1$

$$\lim_{y \rightarrow 1} \frac{y^6 - 1}{y^2 - 1} = \lim_{y \rightarrow 1} \frac{\frac{y^6 - 1^6}{y - 1}}{\frac{y^2 - 1^2}{y - 1}}$$

$$= \frac{6 \times 1^5}{2 \times 1} = \frac{6}{2} = 3$$

19. $\lim_{x \rightarrow a} \frac{x^7 + a^7}{x + a} = 7$ find the value of 'a'

Ans. $\lim_{x \rightarrow a} \frac{x^7 + a^7}{x + a} = 7$

$$= \frac{a^7 + a^7}{a + a} = 7$$

$$= \frac{2a^7}{2a} = 7$$

$$= a^6 = 7$$

$$= a = \sqrt[6]{7}$$

20. Differentiate $x^{-3}(5+3x)$

Ans. $\frac{d}{dx} x^{-3}(5+3x)$

$$= \frac{d}{dx} [5x^{-3} + 3x^{-2}]$$

$$= 5 \times -3x^{-4} + 3 \times -2x^{-3}$$

$$= \frac{-15}{x^4} - \frac{6}{x^3}$$

CBSE Class 12 Mathematics

Important Questions

Chapter 13

Limits and Derivatives

4 Marks Questions

1. Prove that $\lim_{x \rightarrow 0} \left(\frac{e^x - 1}{x} \right) = 1$

Ans. We have

$$\lim_{x \rightarrow 0} \frac{e^x - 1}{x}$$

$$\lim_{x \rightarrow 0} \left\{ \frac{\left[1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots \right] - 1}{x} \right\} \left[\because e^x = 1 + x + \frac{x^2}{2!} + \dots \right]$$

$$\lim_{x \rightarrow 0} \left\{ \frac{x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots}{x} \right\}$$

$$\lim_{x \rightarrow 0} x \left\{ \frac{1 + \frac{x}{2!} + \frac{x^2}{3!} + \dots}{x} \right\}$$

$$= 1 + 0 = 1$$

2. Evaluate $\lim_{x \rightarrow 1} \frac{(2x-3)(\sqrt{x}-1)}{(2x^2+x-3)}$

Ans. $\lim_{x \rightarrow 1} \frac{(2x-3)(\sqrt{x}-1)}{(2x^2+x-3)}$

$$= \lim_{x \rightarrow 1} \frac{(2x-3)(\sqrt{x}-1)}{(2x+3)(x-1)}$$

$$\lim_{x \rightarrow 1} \frac{(2x-3)(\sqrt{x}-1)}{(2x+3)(x-1)} \times \frac{(\sqrt{x}+1)}{(\sqrt{x}+1)}$$

$$\lim_{x \rightarrow 1} \frac{(2x-3)(\cancel{x-1})}{(2x+3)(\cancel{x-1})(\sqrt{x}+1)}$$

$$\lim_{x \rightarrow 1} \frac{(2x-3)}{(2x+3)(\sqrt{x}+1)} = \frac{2 \times 1 - 3}{(2 \times 1 + 3)(\sqrt{1} + 1)}$$

$$= \frac{-1}{10}$$

3. Evaluate $\lim_{x \rightarrow 0} \frac{x \tan 4x}{1 - \cos 4x}$

Ans. $\lim_{x \rightarrow 0} \frac{x \tan 4x}{1 - \cos 4x}$

$$= \lim_{x \rightarrow 0} \frac{x \sin 4x}{\cos 4x [2 \sin^2 2x]}$$

$$= \lim_{x \rightarrow 0} \frac{2x \cancel{\sin 2x} \cos 2x}{\cos 4x (2 \sin^3 2x)}$$

$$= \lim_{x \rightarrow 0} \left(\frac{\cos 2x}{\cos 4x} \cdot \frac{2x}{\sin 2x} \times \frac{1}{2} \right)$$

$$= \frac{1}{2} \frac{\lim_{2x \rightarrow 0} \cos 2x}{\lim_{4x \rightarrow 0} \cos 4x} \times \lim_{2x \rightarrow 0} \left(\frac{2x}{\sin 2x} \right) = \frac{1}{2} \times 1 = \frac{1}{2}$$

4. It $y = \frac{(1 - \tan x)}{(1 + \tan x)}$. Show that $\frac{dy}{dx} = \frac{-2}{(1 + \sin 2x)}$

Ans. $y = \frac{(1 - \tan x)}{(1 + \tan x)}$

$$\frac{dy}{dx} = \frac{(1 + \tan x) \frac{d}{dx} (1 - \tan x) - (1 - \tan x) \frac{d}{dx} (1 + \tan x)}{(1 + \tan x)^2}$$

$$= \frac{(1 + \tan x)(-\sec^2 x) - (1 - \tan x)\sec^2 x}{(1 + \tan x)^2}$$

$$= \frac{-\sec^2 x - \cancel{\tan x \sec^2 x} - \sec^2 x + \cancel{\tan x \sec^2 x}}{(1 + \tan x)^2}$$

$$= \frac{-2\sec^2 x}{(1 + \tan x)^2} = \frac{-2}{\cos^2 x \left[1 + \frac{\sin x}{\cos x} \right]^2}$$

$$= \frac{-2}{\cancel{\cos^2 x} \left[\frac{\cos x + \sin x}{\cancel{\cos^2 x}} \right]^2}$$

$$= \frac{-2}{\cos^2 x + \sin^2 x + 2 \sin x \cos x} = \frac{-2}{1 + \sin^2 x}$$

$$\therefore \frac{dy}{dx} = \frac{-2}{1 + \sin 2x} \quad \text{Hence proved}$$

5. Differentiate $e^{\sqrt{\cot x}}$

Ans. Let $y = e^{\sqrt{\cot x}}$

$$\frac{dy}{dx} = \frac{d}{dx} e^{\sqrt{\cot x}} = e^{\sqrt{\cot x}} \frac{d}{dx} \sqrt{\cot x}$$

$$= e^{\sqrt{\cot x}} \times \frac{1}{2\sqrt{\cot x}} \cdot \frac{d}{dx} \cot x$$

$$= \frac{e^{\sqrt{\cot x}}}{2\sqrt{\cot x}} - \operatorname{cosec}^2 x$$

$$= \frac{-\operatorname{cosec}^2 x e^{\sqrt{\cot x}}}{2\sqrt{\cot x}}$$

6. Let $f(x) = \begin{cases} a + bx, & x < 1 \\ 4, & x = 1 \\ b - ax, & x > 1 \end{cases}$ and if $\lim_{x \rightarrow 1} f(x) = f(1)$ What are the possible value of a

and b ?

Ans. Given $f(1) = 4$

$$\therefore \lim_{x \rightarrow 1} f(x) = f(1) = 4$$

$$\therefore \lim_{x \rightarrow 1} f(x) \text{ exist}$$

$$\therefore \lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^+} f(x) = 4 \text{ --- (1)}$$

$$\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} (a + bx) \quad \left[\begin{array}{l} \because \text{for } x < 1 \\ f(x) = a + bx \end{array} \right]$$

$$= a + b$$

$$\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} (b - ax) \quad \left[\begin{array}{l} \because \text{for } x > 1 \\ f(x) = b - ax \end{array} \right]$$

$$= b - a$$

By eq. (1)

$$a + b = b - a = 4$$

$$a + b = 4$$

$$b - a = 4$$

$$\therefore a = 0 \text{ and } b = 4$$

7. If $y = \frac{1}{\sqrt{a^2 - x^2}}$, find $\frac{dy}{dx}$

Ans. $y = \frac{1}{\sqrt{a^2 - x^2}}$

put $(a^2 - x^2) = t$

$$y = \frac{1}{\sqrt{t}} \text{ and } t = a^2 - x^2$$

$$\frac{dy}{dt} = \frac{d}{dt} t^{\frac{-1}{2}}$$

$$= \frac{-1}{2} t^{\frac{-1}{2} - 1}$$

$$= \frac{-1}{2} t^{\frac{-3}{2}}$$

$$\frac{dt}{dx} = -2x$$

so,

$$\frac{dy}{dx} = \frac{dy}{dt} \times \frac{dt}{dx}$$

$$= \frac{-1}{2} t^{\frac{-3}{2}} \times (-2x) = x t^{\frac{-3}{2}}$$

$$= x(a^2 - x^2)^{\frac{-3}{2}}$$

8. Differentiate $\sqrt{\frac{1 - \tan x}{1 + \tan x}}$

Ans. let $y = \sqrt{\frac{1 - \tan x}{1 + \tan x}}$

put $\frac{1 - \tan x}{1 + \tan x} = t$

$$y = \sqrt{t} \text{ and } t = \frac{1 - \tan x}{1 + \tan x}$$

$$\frac{dy}{dt} = \frac{d}{dt} t^{\frac{1}{2}}$$

$$= \frac{1}{2} t^{\frac{1}{2}-1} = \frac{1}{2} t^{\frac{-1}{2}}$$

$$= \frac{1}{2\sqrt{t}} = \frac{1}{2\sqrt{\frac{1 - \tan x}{1 + \tan x}}} = \frac{1}{2} \sqrt{\frac{1 + \tan x}{1 - \tan x}}$$

$$\frac{dt}{dx} = \frac{(1+\tan x) \frac{d}{dx}(1-\tan x) - (1-\tan x) \frac{d}{dx}(1+\tan x)}{(1+\tan x)^2}$$

$$= \frac{(1+\tan x)(0 - \sec^2 x) - (1-\tan x)(0 + \sec^2 x)}{(1+\tan x)^2}$$

$$= \frac{\sec^2 x [-1 - \cancel{\tan x} - 1 + \cancel{\tan x}]}{(1+\tan x)^2}$$

$$= \frac{-2 \sec^2 x}{(1+\tan x)^2}$$

$$\cancel{\frac{dy}{dx}} = \cancel{\frac{dy}{dt}} \times \cancel{\frac{dt}{dx}} = \frac{1}{2} \sqrt{\frac{1+\tan x}{1-\tan x}}$$

$$\frac{dy}{dx} = \frac{dy}{dt} \times \frac{dt}{dx}$$

$$= \frac{-2 \sec^2 x}{(1+\tan x)^2} \times \frac{1}{2} \sqrt{\frac{1+\tan x}{1-\tan x}}$$

$$= \frac{-\sec^2 x}{(1+\tan x)^{\frac{3}{2}} (1-\tan x)^{\frac{1}{2}}}$$

9. Differentiate (i) $\left(\frac{\sin x + \cos x}{\sin x - \cos x} \right)$ (ii) $\left(\frac{\sin x - 1}{\sec x + 1} \right)$

Ans. (i) $\frac{d}{dx} \left(\frac{\sin x + \cos x}{\sin x - \cos x} \right)$

$$= \frac{(\sin x - \cos x) \cdot \frac{d}{dx}(\sin x + \cos x) - (\sin x + \cos x) \cdot \frac{d}{dx}(\sin x - \cos x)}{(\sin x - \cos x)^2}$$

$$= \frac{(\sin x - \cos x)(\cos x - \sin x) - (\sin x + \cos x)(\cos x + \sin x)}{(\sin x - \cos x)^2}$$

$$= \frac{-[(\sin x - \cos x)^2 + (\sin x + \cos x)^2]}{(\sin x - \cos x)^2}$$

$$= \frac{-2(\sin^2 x + \cos^2 x)}{(\sin^2 x + \cos^2 x - 2\sin x \cos x)}$$

$$= \frac{-2}{(1 - \sin 2x)}$$

$$(ii) \frac{d}{dx} \left[\frac{\sec x - 1}{\sec x + 1} \right]$$

$$= \frac{(\sec x + 1) \cdot \frac{d}{dx}(\sec x - 1) - (\sec x - 1) \cdot \frac{d}{dx}(\sec x + 1)}{(\sec x + 1)^2}$$

$$= \frac{(\sec x + 1) \sec x \tan x - (\sec x - 1) \sec x \tan x}{(\sec x + 1)^2}$$

$$= \frac{2 \sec x \tan x}{(\sec x + 1)^2}$$

10. Evaluate $\lim_{x \rightarrow \frac{\pi}{4}} \frac{\sin x - \cos x}{\left(x - \frac{\pi}{4}\right)}$

Ans. put $\left(x - \frac{\pi}{4}\right) = y$, so that when $x \rightarrow \frac{\pi}{4}$ then $y \rightarrow 0$

$$\begin{aligned} \therefore \lim_{x \rightarrow \frac{\pi}{4}} \frac{\sin x - \cos x}{\left(x - \frac{\pi}{4}\right)} \\ = \lim_{y \rightarrow 0} \frac{\left[\sin\left(\frac{\pi}{4} + y\right) - \cos\left(\frac{\pi}{4} + y\right)\right]}{y} \left[\text{putting } \left(x - \frac{\pi}{4} = y\right)\right] \\ = \lim_{y \rightarrow 0} \frac{\left[\left(\sin \frac{\pi}{4} \cos y + \cos \frac{\pi}{4} \sin y - \sin \frac{\pi}{4} \sin y - \cos \frac{\pi}{4} \cos y\right)\right]}{y} \\ = \frac{2}{\sqrt{2}} \times \lim_{y \rightarrow 0} \left(\frac{\sin y}{y}\right) = (\sqrt{2} \times 1 = \sqrt{2}) \end{aligned}$$

11. Evaluate $\lim_{x \rightarrow 0} \frac{(1+x)^6 - 1}{(1+x)^5 - 1}$

Ans. put $(1+x) = y$, so that when $x \rightarrow 0$ then $y \rightarrow 1$

$$\begin{aligned} \therefore \lim_{x \rightarrow 0} \frac{(1+x)^6 - 1}{(1+x)^5 - 1} \\ = \lim_{y \rightarrow 1} \left[\frac{y^6 - 1}{y^5 - 1} \right] = \frac{\lim_{y \rightarrow 1} \left(\frac{y^6 - 1}{y - 1} \right)}{\lim_{y \rightarrow 1} \left(\frac{y^5 - 1}{y - 1} \right)} \end{aligned}$$

$$\begin{aligned}
 &= \frac{\lim_{y \rightarrow 1} \left(\frac{y^6 - 1}{y - 1} \right)}{\lim_{y \rightarrow 1} \left(\frac{y^5 - 1^5}{y - 1} \right)} = \frac{6 \times 1^{(6-1)}}{5 \times 1^{(5-1)}} \\
 &= \frac{6 \times 1^5}{5 \times 1^4} = \frac{6}{5} \left[\because \lim_{x \rightarrow a} \left(\frac{x^n - a^n}{x - a} \right) = na^{n-1} \right]
 \end{aligned}$$

12. Evaluate $\lim_{x \rightarrow a} \frac{\sqrt{a+2x} - \sqrt{3x}}{\sqrt{3a+x} - 2\sqrt{x}}$

Ans. $\lim_{x \rightarrow a} \frac{(\sqrt{a+2x} - \sqrt{3x})}{(\sqrt{3a+x} - 2\sqrt{x})}$

$$= \lim_{x \rightarrow a} \frac{(\sqrt{a+2x} - \sqrt{3x})}{(\sqrt{3a+x} - 2\sqrt{x})} \times \frac{(\sqrt{3a+x} + 2\sqrt{x})}{(\sqrt{3a+x} + 2\sqrt{x})} \times \frac{(\sqrt{a+2x} + \sqrt{3x})}{(\sqrt{a+2x} + \sqrt{3x})}$$

$$= \lim_{x \rightarrow a} \frac{[(a+2x) - 3x] \times (\sqrt{3a+x} + 2\sqrt{x})}{[(3a+x) - 4x] \times (\sqrt{a+2x} + \sqrt{3x})}$$

$$= \lim_{x \rightarrow a} \frac{(\cancel{a-x}) \times (\sqrt{3a+x} + 2\sqrt{x})}{3(\cancel{a-x}) \times (\sqrt{a+2x} + \sqrt{3x})}$$

$$= \lim_{x \rightarrow a} \frac{(\sqrt{3a+x} + 2\sqrt{x})}{3[\sqrt{a+2x} + \sqrt{3x}]}$$

$$= \frac{(\sqrt{4a} + 2\sqrt{a})}{3(\sqrt{3a} + \sqrt{3a})} = \frac{4\sqrt{a}}{6\sqrt{3}\sqrt{a}} = \frac{\sqrt{2}}{3\sqrt{3}}$$

13. Find the derivative of $f(x) = 1 + x + x^2 + \dots + x^{50}$ at $x = 1$

Ans. $f(x) = 1 + x + x^2 + \dots + x^{50}$

$$f'(x) = \frac{d}{dx}(1 + x + x^2 + \dots + x^{50})$$

$$= 0 + 1 + 2x + 3x^2 + \dots + 50x^{49}$$

At $x = 1$

$$f'(1) = 1 + 2 + 3 + \dots + 50$$

$$= \frac{50(50+1)}{2} = 25 \times 51 \left[\frac{1+2+3+\dots+n}{= \frac{n(n+1)}{2}} \right]$$

$$= 1305$$

14. Find the derivative of $\sin^2 x$ with respect to x using product rule

Ans. let

$$y = \sin^2 x$$

$$y = \sin x \times \sin x$$

$$\frac{dy}{dx} = \frac{d}{dx} \sin x \times \sin x$$

$$= \sin x \frac{d}{dx} \sin x + \sin x \frac{d}{dx} \sin x$$

$$= \sin x \cdot \cos x + \sin x \cdot \cos x$$

$$= 2 \sin x \cdot \cos x = \sin 2x$$

15. Find the derivative of $\frac{x^5 - \cos x}{\sin x}$ with respect to x

Ans. let

$$y = \frac{x^5 - \cos x}{\sin x}$$

$$\frac{dy}{dx} = \frac{d}{dx} \left(\frac{x^5 - \cos x}{\sin x} \right)$$

$$= \frac{\sin x \frac{d}{dx} (x^5 - \cos x) - (x^5 - \cos x) \frac{d}{dx} \sin x}{\sin^2 x}$$

$$= \frac{\sin x [5x^4 + \sin x] - (x^5 - \cos x) \cos x}{\sin^2 x}$$

$$= \frac{5x^4 \sin x + \sin^2 x - x^5 \cos x + \cos^2 x}{\sin^2 x}$$

$$= \frac{5x^4 \sin x - x^5 \cos x + 1}{\sin^2 x}$$

16. Find $\lim_{x \rightarrow 0} f(x)$.

$$\text{when } f(x) = \begin{cases} \frac{|x|}{x}; x \neq 0 \\ 0; x = 0 \end{cases}$$

$$\text{Ans. } f(x) = \begin{cases} \frac{|x|}{x}; x \neq 0 \\ 0; x = 0 \end{cases}$$

We know that $|x| = \begin{cases} x, & x \geq 0 \\ -x, & x < 0 \end{cases}$

$$\therefore f(x) = \begin{cases} \frac{x}{x} = 1, & x > 0 \\ \frac{-x}{x} = -1, & x < 0 \\ 0, & x = 0 \end{cases}$$

L. H. L. $\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0} -1 = -1$

R. H. L. $\lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0} 1 = 1$

L. H. L. \neq R. H. L. $\therefore \lim_{x \rightarrow 0} f(x)$ does not exist

17. Find the derivative of the function $f(x) = 2x^2 + 3x - 5$ at $x = -1$. Also show that $f'(0) + 3f'(-1) = 0$

Ans. $f(x) = 2x^2 + 3x - 5$

$$f^1(x) = \frac{d}{dx}(2x^2 + 3x - 5)$$

$$= 4x + 3$$

At $x = -1$

$$f^1(-1) = 4 \times -1 + 3 = -4 + 3 = -1$$

$$f^1(0) = 4 \times 0 + 3 = 3$$

$$f^1(0) + 3f^1(-1) = 3 + 3 \times -1$$

$$= 3 - 3 = 0 \text{ Hence proved}$$

18. Evaluate $\lim_{x \rightarrow 0} \frac{ax + x \cos x}{b \sin x}$

Ans. $\lim_{x \rightarrow 0} \frac{ax + x \cos x}{b \sin x}$

$$= \lim_{x \rightarrow 0} \frac{x[a + \cos x]}{xb \frac{\sin x}{x}}$$

$$\frac{\lim_{x \rightarrow 0} (a + \cos x)}{\lim_{x \rightarrow 0} b \frac{\sin x}{x}}$$

$$\frac{a+1}{b \times 1} = \frac{a+1}{b} \left[\begin{array}{l} \because \lim_{x \rightarrow 0} \cos x = 1 \\ \lim_{x \rightarrow 0} \frac{\sin x}{x} = 1 \end{array} \right]$$

19. Find derivative of $\tan x$ by first principle

Ans. let $f(x) = \tan x$

$$f(x+h) = \tan(x+h)$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{\tan(x+h) - \tan x}{h}$$

$$= \lim_{h \rightarrow 0} \frac{\frac{\sin(x+h)}{\cos(x+h)} - \frac{\sin x}{\cos x}}{h}$$

$$\begin{aligned}
&= \lim_{h \rightarrow 0} \frac{\sin(x+h)\cos x - \cos(x+h)\sin x}{\cos(x+h)\cos x h} \\
&= \lim_{h \rightarrow 0} \frac{\sin[x+h-x]}{\cos(x+h)\cos x h} \\
&= \frac{\lim_{h \rightarrow 0} \frac{\sin h}{h}}{\lim_{h \rightarrow 0} \cos(x+h)\cos x} = \frac{1}{\cos(x+0)\cos x} \\
&= \frac{1}{\cos^2 x} = \sec^2 x
\end{aligned}$$

20. Evaluate $\lim_{x \rightarrow 1} \frac{x + x^2 + x^3 + \dots + x^n - n}{(x-1)}$

Ans. $\lim_{x \rightarrow 1} \frac{x + x^2 + x^3 + \dots + x^n - 1}{(x-1)}$

$$= \lim_{x \rightarrow 1} \frac{(x-1) + (x^2-1) + (x^3-1) + \dots + (x^n-1)}{(x-1)}$$

$$= \lim_{x \rightarrow 1} \frac{\cancel{(x-1)} [1 + (x+1) + (x^2+x+1) + \dots + x^{n-1} + x^{n-2} + \dots + 1]}{\cancel{(x-1)}}$$

$$= 1 + 2 + 3 + \dots + n = \frac{n(n+1)}{2}$$

21. Evaluate $\lim_{x \rightarrow 4} \frac{|4-x|}{x-4}$ (if it exist)

Ans. $\lim_{x \rightarrow 4} \frac{|4-x|}{x-4}$

$$L.H.L. \quad \lim_{x \rightarrow 4^-} \frac{-(4-x)}{x-4} = \lim_{x \rightarrow 4} \frac{-(4-x)}{-(4-x)} = 1$$

$$R.H.L. \quad \lim_{x \rightarrow 4^+} \frac{4-x}{x-4} = \lim_{x \rightarrow 4} \frac{-(x-4)}{(x-4)} = -1$$

$$L.H.L. \neq R.H.L.$$

$$\therefore \lim_{x \rightarrow 4} \frac{|4-x|}{x-4} \text{ does not exist}$$

22. For what integers m and n does both

$\lim_{x \rightarrow 0} f(x)$ and $\lim_{x \rightarrow 1} f(x)$ exist

$$f(x) = \begin{cases} mx^2 + n; & x < 0 \\ nx + m; & 0 \leq x \leq 1 \\ nx^3 + m; & x > 1 \end{cases}$$

Ans. for $x = 0$

$$L.H.L. \quad \lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0} mx^2 + n \\ = n$$

$$R.H.L. \quad \lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} nx + m \\ = m$$

$$\therefore \lim_{x \rightarrow 0} f(x) \text{ exist}$$

$$\therefore \lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^+} f(x) \\ n = m$$

For all real number $m = n$ $\lim_{x \rightarrow 0} f(x)$ exist

For $x = 1$

$$L.H.L. \lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} nx + m$$

$$= n + m$$

$$R.H.L. \lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} nx^3 + m$$

$$= n + m$$

$$\therefore \lim_{x \rightarrow 1} f(x) = \lim_{x \rightarrow 1^+} f(x)$$

$$m + n = m + n$$

\therefore all integral values of $m + n \lim_{x \rightarrow 1} f(x)$ exist

23. If $y = \sqrt{x} + \frac{1}{\sqrt{x}}$, prove that $2x \frac{dy}{dx} + y = 2\sqrt{x}$

Ans. $y = \sqrt{x} + \frac{1}{\sqrt{x}} = x^{\frac{1}{2}} + x^{-\frac{1}{2}}$

Differentiating w. r. t. x we get

$$\frac{dy}{dx} = \frac{1}{2} x^{-\frac{1}{2}} + \left(\frac{-1}{2} \right) x^{-\frac{3}{2}}$$

$$= \frac{1}{2\sqrt{x}} - \frac{1}{2x^{\frac{3}{2}}}$$

$$2x \frac{dy}{dx} = \sqrt{x} - \frac{1}{\sqrt{x}}$$

$$2x \frac{dy}{dx} + y = \left(\sqrt{x} - \frac{1}{\sqrt{x}} \right) + \left(\sqrt{x} + \frac{1}{\sqrt{x}} \right)$$

$$2x \frac{dy}{dx} + y = 2\sqrt{x} \text{ Hence proved}$$

24. Evaluate $\lim_{x \rightarrow \frac{\pi}{2}} \frac{1 + \cos 2x}{(\pi - 2x)^2}$

Ans. $\lim_{x \rightarrow \frac{\pi}{2}} \frac{1 + \cos 2x}{(\pi - 2x)^2}$

let $\pi - 2x = y$

$2x = \pi - y$

$x \rightarrow \frac{\pi}{2}, y \rightarrow 0$

$$\lim_{y \rightarrow 0} \frac{1 + \cos(\pi - y)}{y^2} = \lim_{y \rightarrow 0} \frac{1 - \cos y}{y^2} \cdot \frac{1}{2}$$

$$= \lim_{y \rightarrow 0} \frac{2 \sin^2 \frac{y}{2}}{4 \times \frac{y^2}{4}}$$

$$= \lim_{y \rightarrow 0} \frac{1}{2} \times \frac{\sin^2 \frac{y}{2}}{\left(\frac{y}{2}\right)^2} =$$

$$= \frac{1}{2} \lim_{\frac{y}{2} \rightarrow 0} \left[\frac{\sin \frac{y}{2}}{\frac{y}{2}} \right]^2$$

$$= \frac{1}{2} \times 1$$

25. Differentiate the function $y = \frac{(x+2)(3x-1)}{(2x+5)}$ with respect to x

Ans. $y = \frac{(x+2)(3x-1)}{(2x+5)}$

$$\begin{aligned}\frac{dy}{dx} &= \frac{d}{dx} \frac{(x+2)(3x-1)}{(2x+5)} \\&= \frac{(2x+5) \frac{d}{dx} (x+2)(3x-1) - (x+2)(3x-1) \frac{d}{dx} (2x+5)}{(2x+5)^2} \\&= \frac{(2x+5) \left[(x+2) \frac{d}{dx} (3x-1) + (3x-1) \frac{d}{dx} (x+2) \right] - (x+2)(3x-1)[2+0]}{(2x+5)^2} \\&= \frac{(2x+5) [(x+2) \times 3 + (3x-1) \times 1] - 2[3x^2 + 6x - x - 2]}{(2x+5)^2} \\&= \frac{(2x+5)[3x+6+3x-1] - 6x^2 - 12x + 2x + 4}{(2x+5)^2} \\&= \frac{12x^2 + 30x + 10x + 25 - 6x^2 - 10x + 4}{(2x+5)^2} \\&= \frac{6x^2 + 30x + 29}{(2x+5)^2}\end{aligned}$$

26. Find $\lim_{x \rightarrow 5} |x| - 5$

Ans. $L.H.S. \lim_{x \rightarrow 5} f(x)$

$$x = 5 - h$$

$$x \rightarrow 5, h \rightarrow 0$$

$$\lim_{h \rightarrow 0} f(5 - h)$$

$$\lim_{h \rightarrow 0} |5 - h| - 5$$

$$= 0$$

$$R.H.S. \lim_{x \rightarrow 5^+} f(x)$$

$$\text{put } x = 5 + h$$

$$x \rightarrow 5, h \rightarrow 0$$

$$\lim_{h \rightarrow 0} f(5 + h) = \lim_{h \rightarrow 0} |5 + h| - 5$$

$$= 0$$

$$R.H.S. = R.H.S.$$

$$\therefore \lim_{x \rightarrow 5^-} f(x) = \lim_{x \rightarrow 5^+} f(x)$$

$$\therefore \lim_{x \rightarrow 5} f(x) \text{ exist}$$

$$\therefore \lim_{x \rightarrow 5} f(x) = 0$$

$$27. \text{ Find } \lim_{x \rightarrow 0} f(x) \text{ and } \lim_{x \rightarrow 1} f(x) \text{ where } f(x) = \begin{cases} 2x+3; x \leq 0 \\ 3(x+1); (x > 0) \end{cases}$$

$$\text{Ans. given } f(x) = \begin{cases} 2x+3, x \leq 0 \\ 3(x+1), x > 0 \end{cases}$$

$$\text{for } x = 0$$

$$L.H.S. \lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0} 2x + 3 = 3$$

$$R.H.S. \lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0} 3(x+1) = 3$$

$$L.H.S. = R.H.S.$$

$$\therefore \lim_{x \rightarrow 0} f(x) \text{ exist}$$

$$\therefore \lim_{x \rightarrow 0} f(x) = 3$$

$$\text{for } x=1$$

$$L.H.S.$$

$$\therefore \lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1} 3(x+1)$$

$$= 3(1+1) = 6$$

$$R.H.S.$$

$$\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1} 3(x+1)$$

$$= 3(1+1) = 6$$

$$L.H.S. = R.H.S.$$

$$\therefore \lim_{x \rightarrow 1} f(x) \text{ exist}$$

$$\lim_{x \rightarrow 1} f(x) = 6$$

28. Find derivative of $\sec x$ by first principle

Ans. let $f(x) = \sec x$

$$f(x+h) = \sec(x+h)$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

$$\begin{aligned}
&= \lim_{h \rightarrow 0} \frac{\sec(x+h) - \sec x}{h} \\
&= \lim_{h \rightarrow 0} \frac{\frac{1}{\cos(x+h)} - \frac{1}{\cos x}}{h} \\
&= \lim_{h \rightarrow 0} \frac{\cos x - \cos(x+h)}{\cos(x+h) \cos x h} \\
&= \lim_{h \rightarrow 0} \frac{-2 \sin \left[\frac{2x+h}{2} \right] \sin \left[\frac{-h}{2} \right]}{\cos(x+h) \cos x h} \\
&= \lim_{h \rightarrow 0} \frac{2 \sin \left[\frac{2x+h}{2} \right] \sin \frac{h}{2}}{\cos(x+h) \cos x h} \quad [\sin(-\theta) = -\sin \theta] \\
&= \lim_{h \rightarrow 0} \frac{2 \sin \frac{h}{2}}{2 \frac{h}{2}} \times \frac{\lim_{h \rightarrow 0} \sin \frac{(2x+h)}{2}}{\lim_{h \rightarrow 0} \cos(x+h) \cos x} \\
&= 1 \times \frac{\sin \left(\frac{2x+0}{2} \right)}{\cos(x+0) \cos x} = \frac{\sin x}{\cos x \cos x} \\
&= \tan x \sec x
\end{aligned}$$

29. Find derivative of $f(x) = \frac{4x + 5 \sin x}{3x + 7 \cos x}$

Ans. $f(x) = \frac{4x + 5 \sin x}{3x + 7 \cos x}$

$$\begin{aligned}
 f^1(x) &= \frac{(3x+7\cos x)\frac{d}{dx}(4x+5\sin x) - (4x+5\sin x) \times \frac{d}{dx}(3x+7\cos x)}{(3x+7\cos x)^2} \\
 &= \frac{(3x+7\cos x)(4+5\cos x) - (4x+5\sin x)(3-7\sin x)}{(3x+7\cos x)^2} \\
 &= \frac{\cancel{12x} + 15x\cos x + 28\cos x + 35\cos^2 x - \cancel{12x} + 28\sin x + 15\sin x + 35\sin^2 x}{(3x+7\cos x)^2} \\
 &= \frac{15x\cos x + 35[\sin^2 x + \cos^2 x] + 28 + \cos x + 43\sin x}{(3x+7\cos x)^2} \\
 &= \frac{15x\cos x + 35 + 28\cos x + 43\sin x}{(3x+7\cos x)^2}
 \end{aligned}$$

30. Find derivative of $\frac{x^n - a^n}{x - a}$

$$\begin{aligned}
 \text{Ans. } & \frac{d}{dx} \frac{x^n - a^n}{x - a} \\
 &= \frac{(x-a)\frac{d}{dx}(x^n - a^n) - (x^n - a^n)\frac{d}{dx}(x-a)}{(x-a)^2} \\
 &= \frac{(x-a)[nx^{n-1} - 0] - (x^n - a^n)[1-0]}{(x-a)^2} \\
 &= \frac{nx^{n-1}(x-a) - x^n + a^n}{(x-a)^2} \\
 &= \frac{nx^n - nax^{n-1} - x^n + a^n}{(x-a)^2} = \frac{x^n(n-1) - nax^{n-1} + a^n}{(x-a)^2}
 \end{aligned}$$

CBSE Class 12 Mathematics
Important Questions
Chapter 13
Limits and Derivatives

6 Marks Questions

1. Differentiate $\tan x$ from first principle.

Ans. $f(x) = \tan x$

$$f(x+h) = \tan(x+h)$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{\tan(x+h) - \tan x}{h}$$

$$= \lim_{h \rightarrow 0} \frac{\frac{\sin(x+h)}{\cos(x+h)} - \frac{\sin x}{\cos x}}{h}$$

$$= \lim_{h \rightarrow 0} \frac{\sin(x+h)\cos x - \cos(x+h)\sin x}{h \cos(x+h)\cos x}$$

$$= \lim_{h \rightarrow 0} \frac{\sin[x+h-x]}{h \cos(x+h)\cos x} \left[\because \sin(A-B) = \sin A \cos B - \cos A \sin B \right]$$

$$= \lim_{h \rightarrow 0} \frac{\sin h}{h \cos(x+h)\cos x}$$



$$= \frac{\lim_{h \rightarrow 0} \frac{\sinh}{h}}{\lim_{h \rightarrow 0} \cos(x+h) \cos x} = \frac{1}{\cos x \cdot \cos x} \left[\because \lim_{h \rightarrow 0} \frac{\sinh}{h} = 1 \right]$$

$$= \frac{1}{\cos^2 x} = \sec^2 x$$

2. Differentiate $(x+4)^6$ From first principle.

Ans. let $f(x) = (x+4)^6$

$$f(x+h) = (x+h+4)^6$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{(x+h+4)^6 - (x+4)^6}{h}$$

$$= \lim_{(x+h+4) \rightarrow (x+4)} \frac{(x+h+4)^6 - (x+4)^6}{(x+h+4) - (x+4)}$$

$$= 6(x+4)^{(6-1)} \left[\because \lim_{x \rightarrow a} \frac{x^n - a^n}{x - a} = na^{n-1} \right]$$

$$= 6(x+4)^5$$

3. Find derivative of cosec x by first principle

Ans. proof let $f(x) = \text{cosec } x$

$$\text{By def, } f(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(h)}{h}$$

$$\begin{aligned}
&= \lim_{h \rightarrow 0} \frac{\operatorname{cosec}(x+h) - \operatorname{cosec} x}{h} \\
&= \lim_{h \rightarrow 0} \frac{\frac{1}{\sin(x+h)} - \frac{1}{\sin x}}{h} = \lim_{h \rightarrow 0} \frac{\sin x - \sin(x+h)}{h \sin(x+h) \sin x} \\
&= \lim_{h \rightarrow 0} \frac{2 \cos \frac{x+x+h}{2} \sin \frac{x-x+h}{2}}{h \sin(x+h) \sin x} \\
&= \lim_{h \rightarrow 0} \frac{2 \cos \left(x + \frac{h}{2}\right) \sin \left(-\frac{h}{2}\right)}{h \sin(x+h) \sin x} \\
&= \frac{\lim_{h \rightarrow 0} \frac{2 \cos \left(x + \frac{h}{2}\right) \sin \frac{h}{2}}{h}}{\cos x \cdot \lim_{h \rightarrow 0} \sin(x+h)} = \frac{\lim_{h \rightarrow 0} \frac{\sin \frac{h}{2}}{\frac{h}{2}}}{\cos x} \\
&= -\frac{\cos x}{\sin x \cdot \sin x} \cdot 1 = -\operatorname{cosec} x \cot x
\end{aligned}$$

4. Find the derivatives of the following functions:

$$(i) \left(x - \frac{1}{x}\right)^3 \quad (ii) \frac{(3x+1)(2\sqrt{x-1})}{\sqrt{x}}$$

Ans. (i) let $f(x) = \left(x - \frac{1}{x}\right)^3 = x^3 - \frac{1}{x^3} - 3x + \frac{1}{x} \left(x - \frac{1}{x}\right)$

$= x^3 - x^{-3} - 3x + 3x^{-1}$. Differentiating w.r.t. x , we get

$$f'(x) = 3x^2 - (-3)x^{-4} - 3 \times 1 + 3 \times (-1)x^{-2}$$

$$= 3x^2 + \frac{3}{x^4} - 3 - \frac{3}{x^2}.$$

$$(ii) \text{ let } f(x) = \frac{(3x+1)(2\sqrt{x}-1)}{\sqrt{x}} = \frac{6x^{\frac{3}{2}} - 3x + 2\sqrt{x} - 1}{\sqrt{x}}$$

$$= 6x - 3x^{\frac{1}{2}} + 2 - x^{-\frac{1}{2}}, d: f \text{ w.r.t. } x. \text{ we get}$$

$$f(x) = 6 \times 1 - 3 \times \frac{1}{2} \times x^{-\frac{1}{2}} + 0 - \left(-\frac{1}{2}\right) x^{-\frac{3}{2}}$$

$$= 6 - \frac{3}{2\sqrt{x}} + \frac{1}{2x^{\frac{3}{2}}}.$$

5. If $f(x) = \begin{cases} |x| + a; x < 0 \\ 0; x = 0 \\ |x| - a; x > 0 \end{cases}$ for what Values of 'a' does $\lim_{x \rightarrow 0} f(x)$ exist

Ans. given $f(x) = \begin{cases} |x| + a; x < 0 \\ 0; x = 0 \\ |x| - a; x > 0 \end{cases}$

$$a=0$$

$$\text{L.H.L. } \lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} |x| + a$$

$$= \lim_{x \rightarrow 0} -x + a = a$$

$$\text{R.H.L. } \lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} |x| - a$$

$$\therefore \lim_{x \rightarrow 0} f(x) \text{ exist}$$

$$\therefore \lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^+} f(x)$$

$$a = -a$$

$$2a = 0$$

$$a = 0$$

At $a = 0 \lim_{x \rightarrow 0} f(x)$ exist

6. Find the derivative of $\sin(x+1)$, with respect to x , from first principle.

Ans. let $f(x) = \sin(x+1)$

$$f(x+h) = \sin(x+h+1)$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{\sin(x+h+1) - \sin(x+1)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{2 \cos \left[\frac{x+h+1+x+1}{2} \right] \sin \left[\frac{x+h+1-x-1}{2} \right]}{h}$$

$$= \lim_{h \rightarrow 0} \frac{2 \cos \left[x+1+\frac{h}{2} \right] \sin \frac{h}{2}}{h}$$

$$= \lim_{h \rightarrow 0} 2 \cos \left(x+1+\frac{h}{2} \right) \times \lim_{h \rightarrow 0} \frac{\sin \frac{h}{2}}{\frac{h}{2}}$$

$$= \cos(x+1) \times 1 = \cos(x+1)$$

7. Find the derivative of $\sin x + \cos x$ from first principle

Ans. let $f(x) = \sin x + \cos x$

$$f(x+h) = \sin(x+h) + \cos(x+h)$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{[\sin(x+h) \cos(x+h)] - [\sin x + \cos x]}{h}$$

$$= \lim_{h \rightarrow 0} \frac{[\sin(x+h) - \sin x] + [\cos(x+h) - \cos x]}{h}$$

$$= \lim_{h \rightarrow 0} \frac{2 \cos \left[\frac{x+h+x}{2} \right] \sin \frac{(x+h-x)}{2} - 2 \sin \frac{(x+h+x)}{2} \times \sin \left[\frac{x+h-x}{2} \right]}{h}$$

$$= \lim_{h \rightarrow 0} \frac{2 \cos \left(x + \frac{h}{2} \right) \sin \frac{h}{2}}{h} + \lim_{h \rightarrow 0} \frac{-2 \sin \left(x + \frac{h}{2} \right) \sin \frac{h}{2}}{h}$$

$$= \lim_{h \rightarrow 0} \cancel{2} \cos \left(x + \frac{h}{2} \right) \frac{\sin \frac{h}{2}}{\cancel{2} \frac{h}{2}} + \lim_{h \rightarrow 0} - \cancel{2} \sin \left(x + \frac{h}{2} \right) \frac{\sin \frac{h}{2}}{\cancel{2} \frac{h}{2}}$$

$$= \cos(x+0) \times 1 - \sin(0+x) \times 1$$

$$= \cos x - \sin x$$

8. Find derivative of

$$(i) \frac{x \sin x}{1 + \cos x} \quad (ii) (ax+b)(x+d)^2$$

Ans. (i) $\frac{d}{dx} \frac{x \sin x}{1 + \cos x}$

$$\begin{aligned}
 &= \frac{(1 + \cos x) \frac{d}{dx} (x \sin x) - x \sin x \frac{d}{dx} (1 + \cos x)}{(1 + \cos x)^2} \\
 &= \frac{(1 + \cos x) \left[x \frac{d}{dx} (\sin x) + \sin x \frac{d}{dx} (x) \right] - x \sin x [0 - \sin x]}{(1 + \cos x)^2} \\
 &= \frac{(1 + \cos x) [x \cos x + \sin x \times 1] + x \sin^2 x}{(1 + \cos x)^2} \\
 &= \frac{x \cos x + x \cos^2 x + \sin x + \sin x \cos x + x \sin^2 x}{(1 + \cos x)^2} \\
 &= \frac{x(\cos^2 x + \sin^2 x) + x \cos x + \sin x + \sin x \cos x}{(1 + \cos x)^2} \\
 &= \frac{x + x \cos x + \sin x + \sin x \cos x}{(1 + \cos x)^2}
 \end{aligned}$$

(ii) $\frac{d}{dx} (ax + b)(cx + d)^2$

$$\begin{aligned}
 &= (ax + b) \frac{d}{dx} (cx + d)^2 + (cx + d)^2 \frac{d}{dx} (ax + b) \\
 &= (ax + b) 2(cx + d) \frac{d}{dx} (cx + d) + (cx + d)^2 \times a \\
 &= 2(ax + b)(cx + d) \times c + a(cx + d)^2 \\
 &= (cx + d) [2c(ax + b) + a(cx + d)]
 \end{aligned}$$

$$= (cx + d)[2acx + 2bc + acx + ad]$$

$$= (cx + d)[3acx + 2abc + ad]$$

9. Evaluate $\lim_{h \rightarrow 0} \frac{(a+h)^2 \sin(a+h) - a^2 \sin a}{h}$

Ans. $\lim_{h \rightarrow 0} \frac{(a+h)^2 \sin(a+h) - a^2 \sin a}{h}$

$$= \lim_{h \rightarrow 0} \frac{(a^2 + 2ah + h^2) \sin(a+h) - a^2 \sin a}{h}$$

$$= \lim_{h \rightarrow 0} \frac{a^2 \sin(a+h) + 2ah \sin(a+h) + h^2 \sin(a+h) - a^2 \sin a}{h}$$

$$= \lim_{h \rightarrow 0} \frac{a^2 [\sin(a+h) - \sin a] + 2ah \sin(a+h) + h^2 \sin(a+h)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{a^2 2 \cos \left[\frac{2a+h}{2} \right] \sin \frac{h}{2}}{2 \frac{h}{2}} + \lim_{h \rightarrow 0} 2a \sin(a+h) + \lim_{h \rightarrow 0} h \sin(a+h)$$

$$= a^2 \cos \left[\frac{2a+0}{2} \right] \times 1 + 2a \sin[a+0] + 0 \times \sin a$$

$$= a^2 \cos a + 2a \sin a$$

10. Differentiate

(i) $\left(\frac{a}{x^4} \right) - \frac{b}{x^2} + \cos x$ (ii) $(x + \cos x)(x - \tan x)$

Ans. (i) $\frac{d}{dx} \left[\frac{a}{x^4} - \frac{b}{x^2} + \cos x \right]$

$$= \frac{d}{dx} ax^{-4} - \frac{d}{dx} bx^{-2} + \frac{d}{dx} \cos x$$

$$= a(-4x^{-5}) - b(-2x^{-3}) - \sin x$$

$$= \frac{-4a}{x^5} + \frac{2b}{x^3} - \sin x$$

(ii) $\frac{d}{dx} (x + \cos x)(x - \tan x)$

$$= (x + \cos x) \frac{d}{dx} (x - \tan x) + (x - \tan x) \frac{d}{dx} (x + \cos x)$$

$$= (x + \cos x)(1 - \sec^2 x) + (x - \tan x)(1 - \sin x)$$

$$= x - x \sec^2 x + \cos x - \cos x \sec^2 x + x - x \sin x - \tan x + \tan x \sin x$$

$$= 2x - x \sec^2 x + \cancel{\cos x} - \cancel{\cos x} - x \sin x - \tan x + \tan x \sin x$$

$$= 2x - x \sec^2 x - x \sin x - \tan x + \tan x \sin x$$