

WITH GRAPH PAPER

केन्द्रीय माध्यमिक शिक्षा बोर्ड, दिल्ली  
विशेष स्कूल सर्टिफिकेट परीक्षा (कक्षा दसवीं)  
परीक्षार्थी प्रवेश-पत्र के अनुसार करें

Mathematics

Subject Code: 041

Date of the Exam:

Date of the Exam: Wednesday 18.03.2015

Time:

Medium of the question paper: English

Code:

Code Number

65/2/C

Set Number

Ⓐ Ⓑ Ⓒ Ⓓ

Mark the correct answer to  
each of the question paper

सही उत्तर चुनिए। प्रत्येक प्रश्न के लिए सही उत्तर चुनिए।

सही उत्तर चुनिए। प्रत्येक प्रश्न के लिए सही उत्तर चुनिए।

सही उत्तर चुनिए। प्रत्येक प्रश्न के लिए सही उत्तर चुनिए।

सही उत्तर चुनिए। प्रत्येक प्रश्न के लिए सही उत्तर चुनिए।

सही उत्तर चुनिए। प्रत्येक प्रश्न के लिए सही उत्तर चुनिए।

सही उत्तर चुनिए। प्रत्येक प्रश्न के लिए सही उत्तर चुनिए।

सही उत्तर चुनिए। प्रत्येक प्रश्न के लिए सही उत्तर चुनिए।

सही उत्तर चुनिए। प्रत्येक प्रश्न के लिए सही उत्तर चुनिए।

सही उत्तर चुनिए। प्रत्येक प्रश्न के लिए सही उत्तर चुनिए।

सही उत्तर चुनिए। प्रत्येक प्रश्न के लिए सही उत्तर चुनिए।

B O H S C A

सही उत्तर चुनिए। प्रत्येक प्रश्न के लिए सही उत्तर चुनिए।

सही उत्तर चुनिए। प्रत्येक प्रश्न के लिए सही उत्तर चुनिए।

सही उत्तर चुनिए। प्रत्येक प्रश्न के लिए सही उत्तर चुनिए।

सही उत्तर चुनिए। प्रत्येक प्रश्न के लिए सही उत्तर चुनिए।

सही उत्तर चुनिए। प्रत्येक प्रश्न के लिए सही उत्तर चुनिए।

सही उत्तर चुनिए। प्रत्येक प्रश्न के लिए सही उत्तर चुनिए।

सही उत्तर चुनिए। प्रत्येक प्रश्न के लिए सही उत्तर चुनिए।

सही उत्तर चुनिए। प्रत्येक प्रश्न के लिए सही उत्तर चुनिए।

सही उत्तर चुनिए। प्रत्येक प्रश्न के लिए सही उत्तर चुनिए।

941-0852 4474974

सही उत्तर चुनिए। प्रत्येक प्रश्न के लिए सही उत्तर चुनिए।

सही उत्तर चुनिए। प्रत्येक प्रश्न के लिए सही उत्तर चुनिए।

य  
3A1  
NR S  
री  
04  
0  
14  
4  
24  
6  
34  
—  
—

17  
18  
19  
20  
21  
22  
23  
24

Linear differential Equations

2015

## Section - C

Q:20

$$(1+x^2) \frac{dy}{dx} = e^{m \tan^{-1} x} - y$$

$$(1+x^2) \frac{dy}{dx} + y = e^{m \tan^{-1} x}$$

$$\frac{dy}{dx} + \frac{y}{1+x^2} = \frac{e^{m \tan^{-1} x}}{1+x^2}$$

comparing it with

$$\frac{dy}{dx} + Py = Q \rightarrow \text{linear differential equation}$$

$$P = \frac{1}{1+x^2}$$

$$\begin{aligned} \text{I.F.} &= e^{\int P dx} \\ &= e^{\int \frac{1}{1+x^2} dx} \end{aligned}$$

$$\text{I.F.} = e^{\tan^{-1} x}$$

$$\left\{ \int \frac{1}{x^2+1} dx = \tan^{-1} x \right.$$



So the equation becomes

$$y \text{ I.F} = \int Q \times \text{I.F} \, dx + c$$

$$y \times e^{\tan^{-1}x} = \int \frac{e^{\tan^{-1}x}}{1+x^2} \times e^{\tan^{-1}x} \, dx + c$$

$$y \times e^{\tan^{-1}x} = \int \frac{e^{(m+1)\tan^{-1}x}}{1+x^2} \, dx + c$$

put  $\tan^{-1}x = t$

$$\frac{1}{1+x^2} \, dx = dt$$

$$y e^{\tan^{-1}x} = \int \frac{e^{(m+1)t}}{1+x^2} \, dx + c$$

$$y e^{\tan^{-1}x} = \frac{e^{(m+1)t}}{(m+1)} + c$$

$$y \text{ or } y e^{\tan^{-1}x} = \frac{e^{(m+1)\tan^{-1}x}}{m+1} + c$$

when  $x=0$   $y=1$

$$ye^{\tan^{-1}x} = \frac{e^{(m+1)\tan^{-1}x}}{(m+1)} + c$$

when  $x=0$   $y=1$

$$1 \times e^{\tan^{-1}0} = \frac{e^{(m+1)\tan^{-1}0}}{(m+1)} + c$$

$$1 \times e^0 = \frac{e^{(m+1)0}}{m+1} + c$$

$$e^0 = 1$$

$$1 = \frac{1}{m+1} + c$$

$$1 - \frac{1}{m+1} = c$$

$$\frac{m+1-1}{m+1} = c$$

$$c = \frac{m}{m+1}$$

So equation is

$$\left[ ye^{\tan^{-1}x} = \frac{e^{(m+1)\tan^{-1}x}}{(1+m)} + \frac{m}{m+1} \right]$$



Q: 21.

$$f(x) = \sin^2 x - \cos x \quad x \in [0, \pi]$$

$$f'(x) = 2 \sin x \cos x - (-\sin x)$$

$$f'(x) = 2 \sin x \cos x + \sin x$$

$$\therefore \text{put } f'(x) = 0$$

$$\text{then } (2 \sin x \cos x + \sin x) = 0$$

$$\sin x (2 \cos x + 1) = 0$$

$$\sin x = 0 \quad \text{or} \quad \cos x = -\frac{1}{2}$$

$$x = 0, \pi \quad \text{or} \quad x = \frac{2\pi}{3}$$

$$\begin{aligned} \text{so } f(0) &= [\sin(0)]^2 - \cos 0 \\ &= 0 - 1 \\ &= -1 \end{aligned}$$

$$\begin{aligned} f(\pi) &= \sin^2 \pi - \cos \pi \\ &= 0 - (-1) \\ &= 1 \end{aligned}$$

$$\begin{aligned} f\left(\frac{2\pi}{3}\right) &= \sin^2 \frac{2\pi}{3} - \cos \frac{2\pi}{3} \\ &= \left(\frac{\sqrt{3}}{2}\right)^2 - \left(-\frac{1}{2}\right) \end{aligned}$$

$$= \frac{3}{4} + \frac{1}{2} = \frac{10}{8} = \frac{5}{4}$$

both the extreme value are automatically included



$$f(0) = -1$$

$$f(\pi) = 1$$

$$f\left(\frac{2\pi}{3}\right) = \frac{5}{4}$$

So Absolute maxima is  $\frac{5}{4}$  at  $x = \frac{2\pi}{3}$

Absolute minima is  $-1$  at  $x = 0$

Q: 22

$$r = \hat{i} + \hat{j} + \hat{k} + \lambda(\hat{i} - \hat{j} + \hat{k})$$

$$r = 4\hat{j} + 2\hat{k} + \mu(2\hat{i} - \hat{j} + 3\hat{k})$$

These lines are coplanar if they are parallel or they are intersecting

But these lines are not parallel.

So they are coplanar if shortest distance between them is zero.

$$\begin{aligned}
 a_1 &= (1, 1, 1) & (a_2 - a_1) &= (-1, 3, 1) \\
 a_2 &= (0, 4, 2) \\
 b_1 &= \langle 1, -1, 1 \rangle \\
 b_2 &= \langle 2, -1, 3 \rangle
 \end{aligned}$$

$$\text{Shortest distance} = \frac{|(a_2 - a_1) \cdot (b_1 \times b_2)|}{|b_1 \times b_2|}$$

$$\text{Let us find } |(a_2 - a_1) \cdot (b_1 \times b_2)|$$

$$(a_2 - a_1) \cdot (b_1 \times b_2) = \begin{vmatrix} -1 & 3 & 1 \\ 1 & -1 & 1 \\ 2 & -1 & 3 \end{vmatrix}$$

(expanding by 1st row)

$$\begin{aligned}
 &= -1(-3+1) - 3(3-2) + 1(-1+2) \\
 &= -1(-2) - 3(1) + 1 \\
 &= 2 + 1 - 3 \\
 &= 0
 \end{aligned}$$

So shortest distance between the lines is zero  
So they are coplanar.

Let  $\langle a, b, c \rangle$  be the direction ratio of normal to the plane.

So the dot product of direction ratio of normal to plane and the direction ratio of line is 0.

$$\begin{aligned} \text{So } a - b + c &= 0 \\ 2a - b + 3c &= 0 \end{aligned}$$

$$\begin{array}{cccc} \underline{a} & \underline{b} & \underline{c} & \\ -1 & 1 & 1 & -1 \\ -1 & 3 & 2 & -1 \end{array}$$

$$\frac{a}{-3+1} = \frac{b}{2-3} = \frac{c}{-1+2}$$

$$\text{So } \langle a, b, c \rangle = \langle -2, -1, 1 \rangle$$

So passing point of line is also the passing point of plane

$$\text{So passing point} = (1, 1, 1)$$





So equation becomes

$$-2(x-1) - 1(y-1) + 1(z-1) = 0$$

$$-2x + 2 - y + 1 + z - 1 = 0$$

$$\boxed{-2x - y + z + 2 = 0}$$

equation of plane.

Q: 23

$f: W \rightarrow W$

$$f(n) = \begin{cases} n-1 & \text{if } n \text{ is odd} \\ n+1 & \text{if } n \text{ is even} \end{cases}$$

→ prove that it is one-one:

Let us suppose that  $n_1, n_2 \in W$   
 $n_1 \neq n_2$  but  $f(n_1) = f(n_2)$

(i) 1st case: if  $n_1$  &  $n_2$  are odd

$$f(n_1) = f(n_2)$$

$$n_1 - 1 = n_2 - 1$$

$$n_1 = n_2$$

which is contradiction.

(ii) 2nd case :  $n_1, n_2$  are even

$$f(n_1) = f(n_2)$$

$$n_1 + 2 = n_2 + 2$$

$$n_1 = n_2$$

which is contradiction.

(iii) 3rd case if  $n_1$  is even &  $n_2$  is odd

$$f(n_1) = f(n_2)$$

$$n_1 + 1 = n_2 - 1$$

$$n_1 + 2 = n_2$$

↓  
even

↓  
odd

if we add 2 in an even no<sup>n</sup> then we get an even no<sup>n</sup> but  $n_2$  is odd which is again contradiction

So from these 3 point we see that

$f(n_1) = f(n_2)$  only if  $n_1 = n_2$   
 $\Rightarrow f$  is one-one.

To prove that  $f$  is onto:

let  $y \in W$

and  $y$  is even

then  $y+1$  is odd and it belongs in whole no.

$$f(y+1) = y+1-1 = y$$

So for every  $y \in W$  which is even there exist a preimage  $y+1$  which is odd and  $(y+1) \in W$

let  $y \in W$

$y$  is odd

then  $y-1$  is even and  $y-1 \in W$

$$f(y-1) = y-1+1 = y$$

So for every  $y \in W$  which is odd there exist a pre image  $y-1 \in W$  which is even.

So for all  $y \in W$  there exist pre image in whole no.  
So  $f$  is onto.

$f$  is one-one and onto so it is invertible

To find  $f^{-1}(x)$

$$ny = x - 1 \quad \text{if } x \text{ is odd}$$

$$y + 1 = x$$

$y$  here is even

so  $x = y + 1$  if  $y$  is even:

$$f^{-1}(x) = x + 1 \quad \text{if } x \text{ is even}$$

$$ny = x + 1 \quad \text{if } x \text{ is even}$$


$y$  is odd here

$$y - 1 = x$$

so  $f^{-1}(x) = x - 1$  if  $x$  is odd

$$\text{so } f^{-1}(x) = \begin{cases} x - 1 & \text{if } x \text{ is odd} \\ x + 1 & \text{if } x \text{ is even} \end{cases}$$

$$\boxed{x \in W} \quad \text{so } \boxed{f^{-1} = f}$$

240

$$y = x - 1 \quad \text{if } x \geq 1$$

$$y = -x + 1 \quad \text{if } x < 1$$

intersection point :

$$y = \sqrt{5-x^2}$$

$$y = x - 1$$

so

$$\sqrt{5-x^2} = (x-1)$$

$$5-x^2 = x^2 + 1 - 2x$$

$$2x^2 - 2x - 4 = 0$$

$$x^2 - x - 2 = 0$$

$$x^2 - 2x + x - 2 = 0$$

$$x(x-2) + 1(x-2) = 0$$

$$\text{so } x = 2 \quad \text{or } x = -1$$

$$x = 2$$

↓ rejected as  $x \geq 1$



$$y = \sqrt{5-x^2}$$

$$y = -x+1$$

$$y \text{ if } x < 1$$

$$5-x^2 = x^2+1-2x$$

$$\text{So } x = 2, -1$$

$$x = 2$$

↓  
rejected

$$\boxed{x = -1}$$

Required Area

$$(A) = \int_{-1}^2 \sqrt{5-x^2} dx - \left[ \int_{-1}^1 (-x+1) dx + \int_1^2 (x-1) dx \right]$$

$$A = \left[ \frac{x\sqrt{5-x^2}}{2} + \frac{5 \sin^{-1} x}{\sqrt{5}} \right]_{-1}^2 - \left[ \frac{-x^2+x}{2} \Big|_{-1}^1 + \frac{x^2-x}{2} \Big|_1^2 \right]$$

$$A = 1 \times 1 + \frac{5 \sin^{-1} 2}{\sqrt{5}} - \left( \frac{-1 \times 2}{2} + \frac{5 \sin^{-1}(-1)}{\sqrt{5}} \right) - \left[ \frac{-1}{2} [1-1] + 2 \right. \\ \left. + \frac{1}{2} [4-1] - 1 \right]$$

$$A = 1 + \frac{5 \sin^2 2}{2} + 1 + \frac{5 \sin^2 1}{2} - \left[ 0 + 2 + \frac{3}{2} - 1 \right]$$

$$A = 2 + \frac{5}{2} \left[ \frac{\sin^2 2}{\sqrt{5}} + \frac{\sin^2 1}{\sqrt{5}} \right] - \frac{5}{2}$$

$$A = \frac{5 \times \pi}{2} + \frac{2-5}{2}$$

$$A = \frac{5\pi}{4} - \frac{1}{2} \text{ sq units}$$

$$\left\{ \begin{aligned} & \frac{\sin^2 2}{\sqrt{5}} + \frac{\sin^2 1}{\sqrt{5}} \\ &= \frac{\sin^2 2 + \sin^2 1}{\sqrt{5}} \\ &= \frac{\pi}{2} \end{aligned} \right\}$$

(1) dx

25°

1st 6 positive integers = (1, 2, 3, 4, 5, 6)

Random Variable = X = larger of 2 numbers.

$x \begin{matrix} |^2 \\ |^1 \end{matrix}$

Sample space S = { (1,2) (1,3) (1,4) (1,5) (1,6)  
 (2,1) (3,1) (4,1) (5,1) (6,1)  
 (2,3) (2,4) (2,5) (2,6) (3,2) (4,2) (5,2)  
 (6,2) (3,4) (3,5) (3,6) (4,3) (5,3) (6,3)  
 (4,5) (4,6), (5,4), (6,4) (5,6) (6,5) }

2

$[4-1]-1$

X	P(X)	$X_i P_i$	$X_i^2 P_i$	Rough
2	$\frac{2}{30} = \frac{1}{15}$	$\frac{2}{15}$	$\frac{4}{15}$	2+6
3	$\frac{4}{30} = \frac{2}{15}$	$\frac{6}{15}$	$\frac{18}{15}$	
4	$\frac{6}{30} = \frac{3}{15}$	$\frac{12}{15}$	$\frac{48}{15}$	$\frac{36 \times 5}{180}$
5	$\frac{8}{30} = \frac{4}{15}$	$\frac{20}{15}$	$\frac{100}{15}$	$\frac{22 \times 48}{70}$
6	$\frac{10}{30} = \frac{5}{15}$	$\frac{30}{15}$	$\frac{180}{15}$	$\frac{200 \times 150}{1500}$

$$\sum X_i P_i = \frac{70}{15}$$

$$\sum X_i^2 P_i = \frac{350}{15}$$

$$\text{Mean} = \frac{\sum X_i P_i}{3} = \frac{14}{3} = 4.66$$

$$\text{Variance} = \sum X_i^2 P_i - (\mu)^2$$

$$= \frac{350}{15} - \frac{70 \times 70}{15 \times 15}$$





$$\sigma^2 = \frac{70}{15} \left[ \frac{5 - 70}{15} \right]$$

$$= \frac{70}{15} \left[ \frac{75 - 70}{15} \right]$$

$$= \frac{70 \times 5}{15 \times 15}$$

$$= \frac{350}{225}$$

$$= \frac{14}{9}$$

$$\sigma^2 = 1.55$$

$$\boxed{\text{Variance} = 1.55}$$

### Section - B

Q:19

$$x^x + x^y + y^x = a^b$$

$$\begin{array}{ccc} \downarrow & \downarrow & \downarrow \\ P & Q & R \end{array}$$

$$\frac{d}{dx} [P + Q + R] = 0$$

$$\left[ \frac{dP}{dx} + \frac{dQ}{dx} + \frac{dR}{dx} = 0 \right]$$

$$P = x^x$$

$$\log P = x \log x$$

$$\frac{1}{P} \frac{dP}{dx} = x \left[ \frac{1}{x} \right] + \log x$$

$$\frac{dP}{dx} = x^x (1 + \log x) \quad - (1)$$

$$Q = x^y$$

$$\log Q = y \log x$$

$$\frac{1}{Q} \frac{dQ}{dx} = \left( \frac{y}{x} + \log x \frac{dy}{dx} \right)$$

$$\frac{dQ}{dx} = x^y \left[ \frac{y}{x} + \log x \frac{dy}{dx} \right] \quad - (2)$$

$$Q = y^x$$

$$R = y^x$$

$$\log R = x \log y$$

$$\frac{1}{R} \frac{dR}{dx} = \frac{x}{y} \frac{dy}{dx} + \log y$$

$$\frac{dx}{dx} = y^n \left[ \frac{x}{y} \frac{dy}{dx} + \log y \right] \quad \text{--- (3)}$$

adding (1) & (2) & (3) we get.

$$x^x + x^x \log x + x^{y-1} y + x^y \log x \frac{dy}{dx} + y^{n-1} x \frac{dy}{dx} + y^n \log y = 0$$

$$\frac{dy}{dx} = - \frac{[x^x(1+\log x) + x^{y-1}y + y^n \log y]}{x^y \log x + y^{n-1}x}$$

Q:18

$$y = e^{ax} \cos bx$$

$$\frac{dy}{dx} = e^{ax} (-\sin bx) b + \cos bx \times e^{ax} \times a$$

$$\frac{dy}{dx} = -b e^{ax} \sin bx + a y \Rightarrow \left[ \frac{1}{b} \left( \frac{dy}{dx} - ay \right) = -e^{ax} \sin bx \right] \quad \text{--- (1)}$$

$$\frac{d^2y}{dx^2} = -b [e^{ax} \cos bx \times b + \sin bx \times e^{ax} \times a] + a \frac{dy}{dx}$$

$$\frac{d^2y}{dx^2} = -b \left[ by - \frac{a}{b} \left( \frac{dy}{dx} - ay \right) \right] + a \frac{dy}{dx} \quad \text{from } \textcircled{1}$$

$$\frac{d^2y}{dx^2} = -b^2y + a \frac{dy}{dx} - a^2y + a \frac{dy}{dx}$$

$$\frac{d^2y}{dx^2} + (a^2 + b^2)y - 2a \frac{dy}{dx} = 0$$

hence proved.

26°

$$Z = 5x + 2y$$

↓  
objective function to be made maximum and minimum

$$x - 2y \leq 2$$

$$3x + 2y \leq 12$$

$$-3x + 2y \leq 3$$

$$x \geq 0, y \geq 0$$



corresponding equations

$$x - 2y = 2$$

put  $(0, 0)$  in it

$$0 - 0 = 0 \leq 2$$

which is true so the region is towards

origin

$$3x + 2y \leq 12 \Rightarrow 3x + 2y = 12$$

put  $(0, 0)$

$$0 \leq 12 \text{ which is true}$$

so region towards origin

$$-3x + 2y = 3$$

put  $(0, 0)$

$$0 \leq 3$$

which is true

so region is towards origin

$x > 0$   $y > 0$ . so it means 1st quadrant.

at A (2,

at B

at C

at

intersection point of

$$-3x + 2y = 3$$

$$3x + 2y = 12$$

$$4y = 15$$

$$y = \frac{15}{4}$$

$$-3x + 2 \times \frac{15}{4} = 3$$

$$-3x = 3 - \frac{15}{2}$$

$$-x = 1 - \frac{5}{2}$$

$$-x = -\frac{3}{2}$$

$$x = \frac{3}{2}$$

$$Z = 5x + 2y$$

at A (2, 0)  $Z = 10$

at B (0, 0)  $Z = 0$

at C  $(0, \frac{3}{2})$   $Z = 3$

at D  $(\frac{3}{2}, \frac{15}{4})$   $Z = \frac{15}{2} + \frac{15}{2} = 15$

intersection point

of:

$$x - 2y = 2$$

$$3x + 2y = 12$$

$$4x = 14$$

$$x = \frac{7}{2}$$

$$\frac{7}{2} - 2y = 2$$

$$\frac{7}{2} - 2 = 2y$$

$$\frac{3}{2} = 2y$$


$$y = \frac{3}{4}$$

at E  $(\frac{3}{2}, \frac{3}{4})$

$$Z = \frac{15}{2} + \frac{3}{2}$$

$$Z = 9$$

Q. 2.6



0 small box + small

So  $Z$  is maximum

$$\text{at } D \begin{pmatrix} 3 \\ 2 \\ 4 \end{pmatrix}$$

$$Z = 15$$

$Z$  is minimum at  $(0, 0)$

$$Z = 0$$

170

	Jans	Mats	Toys	
A =	30	12	70	→ school X
	40	15	55	→ school Y
	35	20	75	→ school Z
	3x3			

B =	25	→ cost of Jans
	100	→ cost of Mats
	50	→ cost of Toys





$$AB = \begin{bmatrix} 30 & 12 & 70 \\ 40 & 15 & 55 \\ 30 & 20 & 75 \end{bmatrix} \begin{bmatrix} 25 \\ 100 \\ 50 \end{bmatrix}$$

$$X = AB = \begin{bmatrix} 750 + 1200 + 3500 \\ 1000 + 1500 + 2750 \\ 875 + 2000 + 3750 \end{bmatrix}$$

$$X = \begin{bmatrix} 5450 \\ 5250 \\ 6625 \end{bmatrix} \begin{array}{l} \rightarrow \text{fund collected by school X} \\ \rightarrow \text{fund collected by school Y} \\ \rightarrow \text{fund collected by school Z} \end{array}$$

fund by X = Rs. 5450

fund by Y = Rs. 5250

fund by Z = Rs. 6625

Total fund = Rs. 17325

They are helping victims and hence



the value of helping others is generated.

$$\underline{16^0} \quad I = \int \frac{x+3}{(x+5)^3} e^x dx$$

$$I = \int \frac{x+5}{(x+5)^3} e^x dx - \int \frac{2}{(x+5)^3} e^x dx$$

$$I = \int \frac{1}{(x+5)^2} e^x dx - \int \frac{2}{(x+5)^3} e^x dx$$

↓  
integrating this by parts  
is taken as 1st function

$$I = \frac{1}{(x+5)^2} e^x - \int \frac{d}{dx} \frac{1}{(x+5)^2} e^x dx - \int \frac{2}{(x+5)^3} e^x dx$$

$$I = \frac{1}{(x+5)^2} e^x + \frac{2}{(x+5)^3} e^x - \int \frac{2}{(x+5)^3} e^x dx$$



$$\text{So } I = \frac{e^x}{(x+5)^2} + C$$

15°

$$x = a \sin at (1 + \cos at)$$

$$\frac{dx}{dt} = a [(\sin at) (-\sin at)(2) + (1 + \cos at) \cos at \times 2]$$

$$\frac{dx}{dt} = 2a [\cos^2 at + \cos at - \sin^2 at]$$

$$\frac{dx}{dt} = 2a [-\cos 4t + \cos at] \quad [\cos^2 x - \sin^2 x = \cos 2x]$$

$$y = b \cos at (1 - \cos at)$$

$$y = b \cos at - b \cos^2 at$$

$$\frac{dy}{dt} = -b \sin at \times 2 + b \times 2 \cos at \sin at \times 2$$

$$\frac{dy}{dt} = 2b [-\sin at + \sin 4t]$$



$$\frac{dy}{dx} = \frac{dy}{dt} \cdot \frac{dx}{dt}$$

$$\frac{dy}{dx} = \frac{2b}{2a} \left[ \frac{\sin 4t - \sin 2t}{\cos 2t + \cos 4t} \right]$$

$$\text{at } t = \frac{\pi}{4}$$

$$\frac{dy}{dx} = \frac{b}{a} \left[ \frac{\sin \pi - \sin \frac{\pi}{2}}{\cos \frac{\pi}{2} + \cos \pi} \right]$$

$$= \frac{b}{a} \left[ \frac{0 - 1}{0 - 1} \right]$$

$$\left. \frac{dy}{dx} \right|_{t = \frac{\pi}{4}} = \frac{b}{a}$$

140

$$I = \frac{\pi}{2} \int_0^{\frac{\pi}{2}} \frac{\sin^2 x}{\sin x + \cos x} dx \quad \text{--- (1)}$$

$$\left[ \int_0^a f(x) dx = \int_0^a f(a-x) dx \right]$$

$$I = \frac{\pi}{2} \int_0^{\frac{\pi}{2}} \frac{\sin^2(a-x)}{\sin(a-x) + \cos(a-x)} dx$$

$$I = \frac{\pi}{2} \int_0^{\frac{\pi}{2}} \frac{\cos^2 x}{\cos x + \sin x} dx \quad \text{--- (2)}$$

adding (1) & (2) we get

$$2I = \frac{\pi}{2} \int_0^{\frac{\pi}{2}} \frac{\sin^2 x + \cos^2 x}{\sin x + \cos x} dx$$

$$2I = \frac{\pi}{2} \int_0^{\frac{\pi}{2}} \frac{1}{(\sin x + \cos x)} dx$$



$$2I = \int_0^{\frac{\pi}{2}} \frac{1 + \tan^2 \frac{x}{2}}{2 \tan \frac{x}{2} + 1 - \tan^2 \frac{x}{2}} dx$$

$$2I = \int_0^{\frac{\pi}{2}} \frac{\sec^2 \frac{x}{2}}{2 \tan \frac{x}{2} + 1 - \tan^2 \frac{x}{2}} dx$$

$$\left. \begin{aligned} \text{put } \sin x &= 2 \tan \frac{x}{2} \\ \text{and } \cos x &= \frac{1 + \tan^2 \frac{x}{2}}{1 - \tan^2 \frac{x}{2}} \end{aligned} \right\}$$

$$\text{put } \tan \frac{x}{2} = t$$

$$\sec^2 \frac{x}{2} \times \frac{1}{2} dx = dt$$

$$\begin{aligned} x \rightarrow 0 \quad t \rightarrow 0 \\ \text{when } x \rightarrow \frac{\pi}{2} \quad t \rightarrow 1 \end{aligned}$$

$$2I = \int_0^1 \frac{dt}{2t + 1 - t^2}$$

$$I = \int_0^1 \frac{dt}{2t + 1 + 1 - t^2} \Rightarrow I = \int_0^1 \frac{dt}{2 + (t-1)^2}$$



$$I = \int_0^1 \frac{dt}{(\sqrt{2})^2 - (t-1)^2} = \int \frac{1}{a^2 - x^2} = \frac{1}{2a} \log \left| \frac{a+x}{a-x} \right|$$

$$I = \frac{1}{2\sqrt{2}} \log \left| \frac{\sqrt{2} + t - 1}{\sqrt{2} - t + 1} \right| \Bigg|_0^1$$

$$I = \frac{1}{2\sqrt{2}} \log \left| \frac{\sqrt{2} + 1 - 1}{\sqrt{2} - 1 + 1} \right| - \frac{1}{2\sqrt{2}} \log \left| \frac{\sqrt{2} - 1}{\sqrt{2} + 1} \right|$$

$$I = \frac{1}{2\sqrt{2}} \log 1 - \frac{1}{2\sqrt{2}} \log \left| \frac{\sqrt{2} - 1}{\sqrt{2} + 1} \right|$$

$$= 0 - \frac{1}{2\sqrt{2}} \log \left| \frac{\sqrt{2} - 1}{\sqrt{2} + 1} \right|$$

$$I = -\frac{1}{2\sqrt{2}} \log \left| \frac{\sqrt{2} - 1}{\sqrt{2} + 1} \right|$$

130

$$\Delta = \begin{vmatrix} x+2 & x+6 & x-1 \\ x+6 & x-1 & x+2 \\ x-1 & x+2 & x+6 \end{vmatrix}$$

$$R_1 \rightarrow R_1 + R_2 + R_3$$

$$\Delta = \begin{vmatrix} 3x+7 & 3x+7 & 3x+7 \\ x+6 & x-1 & x+2 \\ x-1 & x+2 & x+6 \end{vmatrix}$$

$$\Delta = 3x+7 \begin{vmatrix} 1 & 1 & 1 \\ x+6 & x-1 & x+2 \\ x-1 & x+2 & x+6 \end{vmatrix}$$

$$C_1 \rightarrow C_1 - C_2 \quad C_2 \rightarrow C_2 - C_3$$

$$\Delta = (3x+7) \begin{vmatrix} 0 & 0 & 1 \\ 7 & -3 & x+2 \\ -3 & -4 & x+6 \end{vmatrix}$$



expanding by  $R_1$

$$\Delta = (3x+7) [-28 - 9]$$

$$\Delta = 0$$

$$\Rightarrow 3x+7=0$$

$$x = \frac{-7}{3}$$

120

$$A = \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & 2 \\ 2 & 2 & 1 \end{bmatrix}$$

$$A^2 = AA = \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & 2 \\ 2 & 2 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & 2 \\ 2 & 2 & 1 \end{bmatrix}$$

$$A^2 = \begin{bmatrix} 1+4+4 & 2+2+4 & 2+4+2 \\ 2+2+4 & 4+1+4 & 4+2+2 \\ 2+4+2 & 4+2+2 & 4+4+1 \end{bmatrix}$$

$$A^2 = \begin{bmatrix} 9 & 8 & 8 \\ 8 & 9 & 8 \\ 8 & 8 & 9 \end{bmatrix}$$

$$A^2 - 4A - 5I = \begin{bmatrix} 9 & 8 & 8 \\ 8 & 9 & 8 \\ 8 & 8 & 9 \end{bmatrix} + \begin{bmatrix} -4 & -8 & -8 \\ -8 & -4 & -8 \\ -8 & -8 & -4 \end{bmatrix} + \begin{bmatrix} -5 & 0 & 0 \\ 0 & -5 & 0 \\ 0 & 0 & -5 \end{bmatrix}$$

$$A^2 - 4A - 5I = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} = 0$$

$$\therefore A^2 - 4A - 5I = 0$$

Hence proved.



$$A^2 - 4A - 5I = 0$$

pre-multiplying by  $A^{-1}$

$$A^{-1}AA - 4A^{-1}A - 5A^{-1}I = 0$$

$$IA - 4I - 5A^{-1} = 0$$

$$A - 4I - 5A^{-1} = 0$$

$$A - 4I = 5A^{-1}$$

$$A^{-1} = \frac{A - 4I}{5}$$

$$A^{-1} = \frac{1}{5} [A - 4I]$$

$$5A^{-1} = \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & 2 \\ 2 & 2 & 1 \end{bmatrix} \times \begin{bmatrix} -4 & 0 & 0 \\ 0 & -4 & 0 \\ 0 & 0 & -4 \end{bmatrix}$$

$$5A^{-1} = \begin{bmatrix} -3 & 2 & 2 \\ 2 & -3 & 2 \\ 2 & 2 & -3 \end{bmatrix}$$

$$A^{-1}A = I$$

$$IA = A$$

$$\text{So } A^{-1} = \frac{1}{5} \begin{bmatrix} -3 & 2 & 2 \\ 2 & -3 & 2 \\ 2 & 2 & -3 \end{bmatrix}$$

11.

$$\sin^{-1}(1-x) - 2\sin^{-1}x = \frac{\pi}{2}$$

$$\sin^{-1}(1-x) = \frac{\pi}{2} + 2\sin^{-1}x$$

$$(1-x) = \sin\left(\frac{\pi}{2} + 2\sin^{-1}x\right)$$

$$(1-x) = \cos(2\sin^{-1}x)$$

$$2\sin^{-1}x = \theta$$

$$x = \sin\theta$$

$$\cos 2\theta = 1 - 2\sin^2\theta$$

$$\cos 2\theta = 1 - 2x^2$$

$$\Rightarrow 1-x = \cos 2\theta$$

$$1-x = 1-2x^2$$

$$\Rightarrow -2x^2 - x = 0$$

$$x(2x-1) = 0$$

$$x=0 \text{ or } x=\frac{1}{2}$$

put  $x=\frac{1}{2}$  in equation

$$x \sin^{-1} \frac{1}{2} - 2 \sin^{-1} \frac{1}{2}$$

$$= \frac{1}{6} - 2 \times \frac{1}{6}$$

$$\neq \frac{1}{2}$$

$$\text{So } x \neq \frac{1}{2}$$

$$\text{So } \boxed{x=0}$$

10°

passing point of line = (4, 2, 2)

since B is || to line

So direction ratio of line =  $\langle 2, 3, 6 \rangle$



So equation of line

$$\frac{x-4}{2} = \frac{y-2}{3} = \frac{z-2}{6} = \lambda$$

general point on line

$$(2\lambda+4, 3\lambda+2, 6\lambda+2)$$

$$P(1, 2, 3)$$

Q

line

Q be the foot of  $\perp$  on line

PQ is  $\perp$  to line

So dot product of direction ratios of line and PQ is 0

direction ratios of

$$= \langle 2\lambda+3, 3\lambda, 6\lambda-1 \rangle$$

also according to question:

$$2(2\lambda+3) + (3\lambda)3 + 6(6\lambda-1) = 0$$

$$4\lambda + 6 + 9\lambda + 36\lambda - 6 = 0$$

$$\boxed{\lambda = 0}$$

also point Q = (4, 2, 2)

also len distance

$$PQ = \sqrt{(4-1)^2 + (2-2)^2 + (2-3)^2}$$

$$= \sqrt{(3)^2 + (0)^2 + (-1)^2}$$

$$= \sqrt{9+1}$$

$$\boxed{\text{length of Lar} = \sqrt{10} \text{ unit}}$$



90

AB  
these A, B, C, D are coplaner :

so  $\vec{AB} \cdot (\vec{BC} \times \vec{CD}) = 0$   
triple product is 0.

$$\vec{AB} = 1\hat{i} + (x-1)\hat{j} + 4\hat{k}$$

$$\vec{BC} = 0\hat{i} + (1-x)\hat{j} - 7\hat{k}$$

$$\vec{CD} = 2\hat{i} + 3\hat{j} + \hat{k}$$

1 3 4  
0 3 2  
2 3 3

$$\vec{AB} \cdot (\vec{BC} \times \vec{CD}) = \begin{vmatrix} 1 & x-1 & 4 \\ 0 & 1-x & -7 \\ 2 & 3 & 1 \end{vmatrix} = 0$$

expanding by R<sub>1</sub>

$$1(1-x+21) - (x-1)14 + 4(2(x-1)) = 0$$
$$22-x - 14x+14 + 8x-8 = 0 \quad \boxed{x=4}$$
$$-7x = -28$$



8°

$$p \text{ (probability of success)} = \frac{1}{2}$$

i.e. that head comes

$$q \text{ (probability of failure)} = \frac{1}{2}$$

i.e. that tail comes

Let the coin be tossed  $n$  times

this event follows the conditions of Bernoulli trial

$X$  be the random variable = no. of heads

$$P(X \geq 1) = P(1) + \dots + P(n)$$

$$P(X \geq 1) = 1 - P(0)$$

$$= 1 - {}^n C_0 \times \left(\frac{1}{2}\right)^0 \left(\frac{1}{2}\right)^n$$

$P(X \geq 1)$  should be more than 80%.

$$P(X \geq 1) > \frac{8}{10}$$

$$\frac{80}{100} < 1 - \left(\frac{1}{2}\right)^n$$

$$1 - \left(\frac{1}{2}\right)^n \geq \frac{8}{10}$$

$$\left(\frac{1-8}{10}\right) \geq \left(\frac{1}{2}\right)^n$$

$$\frac{18}{50} > \left(\frac{1}{2}\right)^n$$

$$5 < 2^n$$

$$n = 3$$

follow the condition  
So the coin should be tossed  
at least 3 times.

7.

$$I = \int \frac{x^2}{x^4 + x^2 - 2} dx$$

$$I = \int \frac{x^2}{x^4 + 2x^2 - x^2 - 2} dx$$

$$I = \int \frac{x^2}{x^2(x^2 + 2) - 1(x^2 + 2)} dx$$



$$I = \int \frac{x^2}{(x^2-1)(x^2+2)} dx$$

put  $x^2 = t$

then 
$$\frac{t}{(t-1)(t+2)} = \frac{A}{t-1} + \frac{B}{t+2}$$

$$t = A(t+2) + B(t-1)$$

put  $t=1$

$$1 = A \times 3$$

$$\boxed{A = \frac{1}{3}}$$

put  $t=-2$

$$-2 = -3B$$

$$\boxed{B = \frac{2}{3}}$$



$$I = \int \frac{1}{3(t-1)} dt + \frac{2}{3} \int \frac{dx}{t+2}$$

$$I = \int \frac{1}{3(x^2-1)} dx + \frac{2}{3} \int \frac{dx}{x^2+2} \quad \int \frac{1}{x^2-a^2} dx = \frac{1}{2a} \log \left| \frac{x-a}{x+a} \right|$$

$$I = \frac{1}{3} \times \frac{1}{2} \log \left| \frac{x-1}{x+1} \right| + \frac{2}{3} \times \frac{1}{\sqrt{2}} \tan^{-1} \frac{x}{\sqrt{2}} + C \quad \int \frac{1}{x^2+a^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}$$

$$I = \frac{1}{6} \log \left| \frac{x-1}{x+1} \right| + \frac{\sqrt{2}}{3} \tan^{-1} \frac{x}{\sqrt{2}} + C$$

⊗

## Section - A

Q: 6 equation of plane

$$6x - 3y + 2z - 4 = 0$$

$$\text{distance} = \frac{|6 \times 2 - 3 \times 5 + 2(-3) - 4|}{\sqrt{36 + 9 + 4}}$$

$$= \frac{|12 - 15 - 6 - 4|}{\sqrt{49}}$$

$$= \frac{|-13|}{7}$$

$$\boxed{\text{distance} = \frac{13}{7} \text{ units}}$$

Q: 5:

$$\vec{a} = \hat{i} - \hat{j}$$

$$|\vec{a}| = \sqrt{2}$$

$$\vec{b} = \hat{j} - \hat{k}$$

$$|\vec{b}| = \sqrt{2}$$

0902

Fictitious Roll No.  
(To be entered by Board)

4474874

अपना अनुक्रमीक इस उत्तर-पुस्तिका

पर न लिखें

Please do not write your

Roll Number on this Answer-Book

अतिरिक्त उत्तर-पुस्तिका (सं.) को सहायक

Supplementary Answer-Books No.

$$\vec{a} \cdot \vec{b} = |\vec{a}| |\vec{b}| \cos \theta$$

$$(\hat{i} - \hat{j}) \cdot (\hat{j} - \hat{k}) = \sqrt{2} \times \sqrt{2} \cos \theta$$

$$\frac{-1}{2} = \cos \theta$$

$$\theta = \frac{2\pi}{3}$$

angle between vectors =  $\frac{2\pi}{3}$ 

4.

$$\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 1 & 3 \\ 3 & 5 & -2 \end{vmatrix}$$

$$\vec{a} \times \vec{b} = \hat{i}(-2-15) - \hat{j}(-4-9) + \hat{k}(10-3)$$

$$\vec{a} \times \vec{b} = -17\hat{i} + 13\hat{j} + 7\hat{k}$$

$$|\vec{a} \times \vec{b}| = \sqrt{(-17)^2 + (13)^2 + (7)^2}$$

$$= \sqrt{289 + 169 + 49}$$

$$= \sqrt{507}$$

$$= \sqrt{3 \times 169}$$

$$|\vec{a} \times \vec{b}| = 13\sqrt{3}$$

Q: 3°

$$x \log x \frac{dy}{dx} + y = 2 \log x$$

$$\frac{dy}{dx} + \frac{y}{x \log x} = 2 \log x$$

compare it with  $\frac{dy}{dx} + Py = Q$

$$I \cdot F = e^{\int P dx}$$

$$= e^{\int \frac{1}{x \log x} dx}$$

put  $\log x = t$   
 $\frac{1}{x} dx = dt$



$$\begin{aligned} I \cdot F &= e^{\int \frac{dt}{t}} \\ &= e^{\log|t|} \\ &= t \end{aligned}$$

$$I \cdot F = \log x$$

2°

general equation of family of lines passing through origin

$$y = mx$$

$$m = \frac{y}{x}$$

$$\frac{dy}{dx} = m$$

$$\frac{dy}{dx} = \frac{y}{x}$$

$$x \frac{dy}{dx} - y = 0$$





Q:10

$$a_{12} = e^{2x} \sin x$$

$$a_{11} = e^{2x} \sin x$$

POANAM 011074009/A  
Evaluation done as per CBSE marking scheme.

XI - U