

MIND MAP : LEARNING MADE SIMPLE

CHAPTER - 3

If $A = [a_{ij}]_{m \times n}$, then its transpose A^T (A^T) = $[a_{ji}]_{n \times m}$ i.e. if $A = \begin{pmatrix} 2 & 1 \end{pmatrix}$ then $A^T = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$.

Also, $(A^T)^T = A$, $(kA)^T = kA^T$, $(A+B)^T = A^T + B^T$, $(AB)^T = B^T A^T$.

- A is symmetric matrix if $A = A^T$ i.e. $A^T = A$.
- A is skew-symmetric if $A = -A^T$ i.e. $A^T = -A$.
- A is any matrix, then $A = \frac{1}{2} \left\{ (A + A^T) + (A - A^T) \right\}$ = sum of a symmetric and a skew-symmetric matrix.

For eg if $A = \begin{pmatrix} 2 & 8 \\ 6 & 4 \end{pmatrix}$, then $A = \frac{1}{2} \left\{ \begin{pmatrix} 2 & 7 \\ 7 & 4 \end{pmatrix} + \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \right\}$.

$$\begin{aligned} R_i &\leftrightarrow R_j \text{ or } C_i \leftrightarrow C_j \\ R_i &\rightarrow kR_i \text{ or } C_i \rightarrow kC_i \\ R_i &\rightarrow R_i + kR_j \text{ or } C_i \rightarrow C_i + kC_j \end{aligned}$$

If A, B are square matrices such that $AB = BA = I$ then $B = A^{-1}$ i.e., A is the inverse of B and vice-versa.

Inverse of a square matrix, if it exists, is unique

$$\text{For eg: If } A = \begin{pmatrix} 1 & 2 \\ 3 & 4 \\ 5 & 6 \end{pmatrix}, \text{ then after } R_1 \leftrightarrow R, A \text{ becomes } \begin{pmatrix} 3 & 4 \\ 1 & 2 \\ 5 & 6 \end{pmatrix}$$

If A and B are invertible matrices of the same order, then $(AB)^{-1} = B^{-1}A^{-1}$. By elementary transformations, we can convert $A = IA$ to $A^{-1}A$. This is one process of finding the inverse of a given square matrix A .

Transpose of a Matrix

Elementary operations on a matrix
Inverse of a matrix

Matrices

Definition and its types

A matrix of order $m \times n$ is an ordered rectangular array of numbers or functions having 'm' rows and 'n' columns. The matrix

$A = [a_{ij}]_{m \times n}$ is given by

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix}_{m \times n}$$

- Column matrix : It is of the form $\begin{bmatrix} a_{ij} \end{bmatrix}_{m \times 1}$
- Row matrix : It is of the form $\begin{bmatrix} a_{ij} \end{bmatrix}_{1 \times n}$
- Square matrix : Here, $m = n$ (no. of rows = no. of columns)
- Diagonal matrix : All non-diagonal entries are zero i.e. $a_{ij} = 0 \forall i \neq j$
- Scalar matrix : $a_{ij} = 0, i \neq j$ and $a_{ii} = k$ (Scalar), $i = j$
- Identity matrix : $a_{ij} = 0, i \neq j$ and $a_{ii} = 1, i = j$
- Zero matrix : All entries are zero.

Equality of two matrix

$$A = \begin{bmatrix} a_{ij} \end{bmatrix} = \begin{bmatrix} b_{ij} \end{bmatrix} = B \text{ if, } A \text{ and } B \text{ are of same order and } a_{ij} = b_{ij} \forall i \text{ and } j.$$

Operations on matrices
Addition

If A, B are two matrices of same order, then $A+B = [a_{ij}+b_{ij}]$. The addition of A and B follows:
 $A+B=B+A$, $(A+B)+C=A+(B+C)$, $-A=(-1)A$, $k(A+B)=kA+kB$, k is scalar and $(k+I)A=kA+IA$, k and I are constants.

Multiplication

- If $A = \begin{pmatrix} 2 & 3 \\ 2 & 4 \end{pmatrix}$, $B = \begin{pmatrix} -3 & 2 \\ -4 & 5 \end{pmatrix}$ then $A+B = \begin{pmatrix} -1 & 5 \\ -2 & 9 \end{pmatrix}$
- If $A = (2 \ 3)_{1 \times 2}$, $B = \begin{pmatrix} 4 \\ 5 \end{pmatrix}_{2 \times 1}$, then $AB = (2 \times 4 + 3 \times 5) = (2 \ 3)_{1 \times 1}$

If $A = [a_{ij}]_{m \times n}$ and $B = [b_{jk}]_{n \times p}$, then $AB = C = [C_{ik}]_{m \times p}$, $[C_{jk}] = \sum_{i=1}^n a_{ij} b_{jk}$. Also,
 $A(BC) = (AB)C$, $A(B+C) = AB + AC$ and $(A+B)C = AC + BC$, but $AB \neq BA$ (always).

Get More Learning Materials Here :

CLICK HERE



www.studentbro.in