

Number Systems

History and Evolution of Irrational Numbers

The existence of irrationality in numbers was accepted by Indian mathematicians as far back as 7th century BC when Manava, an author of the Indian geometric text *Sulbasutras*, discovered (while finding the hypotenuse of a right-angled triangle) that it is not possible to accurately calculate the square roots of numbers like 2 and 8. It is, however, the Pythagorean school of Greek mathematicians, or the Pythagoreans, who are credited with discovering irrational numbers sometime in 400 BC. In 5th century AD, the great Indian mathematician Aryabhata suggested that the value of π is incommensurable. Later, in the 1700s, a Swiss mathematician named Lambert and a French mathematician named Legendre proved π to be irrational.

In this way, a long line of mathematicians helped shed light on the concept of irrational numbers. These mathematicians questioned the rationality of those numbers that cannot be written in the form of a ratio of integers. The Pythagoreans were the first to actually prove a number to be irrational and this number was $\sqrt{2}$. The set of all irrational numbers is denoted by \mathbb{Q}' .


Go through this lesson to get a basic idea about the irrationality of numbers.

Whiz Kid

Golden ratio

Two quantities are said to be in the golden ratio if the ratio of the sum of those quantities to the larger quantity is the same as the ratio of the larger quantity to the smaller one. Let us understand this concept.

Say a and b are two line segments that are in the golden ratio.



Therefore, $\frac{a+b}{a} = \frac{a}{b}$



The golden ratio is represented by the Greek letter ' ϕ ' (phi), where $\phi = \frac{1+\sqrt{5}}{2} = 1.61803...$, an irrational number.

The golden ratio is also known as 'the golden mean' and 'the golden section'. This ratio is used not only in mathematics but also in biology, art, music, architecture and in various other branches of science.

Did You Know?

Pi is a constant value that is equal to the ratio of the circumference of a circle to its diameter. It is an irrational number represented by the Greek letter ' π '. This symbol was proposed by a Welsh mathematician named William Jones in 1706. The value of pi is

approximately equal to $\frac{22}{7}$.

The Great Pyramid of Giza was constructed with a perimeter of about 1760 cubits and a height of about 280 cubits. The ratio of the perimeter to the height, i.e., $\frac{1760}{280}$ is approximately equal to 6.285, which is almost equal to 2π . This is cited by some as the proof that the people who built the pyramid knew about the special ratio represented by π .

Since π is closely related to the circle, it is found in many geometric and trigonometric formulae. It is also used in many other scientific formulae such as in thermodynamics, the number theory, mechanics and electromagnetism.

Know Your Scientist



Aryabhata (476 AD–550 AD) was the first Indian mathematician and astronomer. Belonging to the Indian classical age, he is primarily known for the invention of zero. His works *Aryabhatiya* and *Arya-sidhanta* are incomparable sources of astronomy and mathematics. Some of the concepts put forth by Aryabhata helped in the evolution of modern mathematics and astronomy. These include the place value system, the approximation of pi, the formula for the area of a triangle, the solution of indeterminate equations, the summation of series of cubes and squares, and the explanations for the motion of solar systems, solar and lunar eclipses and sidereal periods.

Know More

1. No rational number (e.g. 5) is irrational and no irrational number (e.g. $\sqrt{5}$) is rational as the properties of one are different from those of the other.
2. There are no smallest or largest numbers in the groups of rational and irrational numbers.

Did You Know?

The need for numbers

Initially, humans used to keep count of different things by using pebbles, sticks or fingers. Later, they began using tally marks for counting purposes. The system of tally marks is not based on the concept of place value. This made the counting and representation of large numbers difficult. Thus grew the need for simpler counting systems and, with the passage of time, the Roman, Hindu-Arabic and other numeral systems came into being.

Solved Examples

Easy

Example 1: Check whether the given numbers are rational or irrational.

i) $(7 - 2\sqrt{5}) + 2\sqrt{5}$

ii) 0

iii) $\sqrt{\frac{49}{81}}$

iv) $\sqrt{289}$

v) 5.012896896896...

vi) 7.1931931934...

vii) 0.0787887888...

Solution:



$$\text{i) } (7 - 2\sqrt{5}) + 2\sqrt{5}$$

$$= 7 - 2\sqrt{5} + 2\sqrt{5}$$

$$= 7$$

$$= \frac{7}{1}$$

$$\frac{p}{q}$$

This number is of the form $\frac{p}{q}$, where $p = 7$ and $q = 1$ are integers and $q \neq 0$.

Hence, $(7 - 2\sqrt{5}) + 2\sqrt{5}$ is a rational number.

$$\text{ii) } 0 = \frac{0}{1}$$

$$\frac{p}{q}$$

This number is of the form $\frac{p}{q}$, where $p = 0$ and $q = 1$ are integers and $q \neq 0$.

Hence, 0 is a rational number.

$$\text{iii) } \sqrt{\frac{49}{81}} = \frac{7}{9}$$

$$\frac{p}{q}$$

This number is of the form $\frac{p}{q}$, where $p = 7$ and $q = 9$ are integers and $q \neq 0$.

Hence, $\sqrt{\frac{49}{81}}$ is a rational number.

$$\text{iv) } \sqrt{289} = 17 = \frac{17}{1}$$

$$\frac{p}{q}$$

This number is of the form $\frac{p}{q}$, where $p = 17$ and $q = 1$ are integers and $q \neq 0$.

Hence, $\sqrt{289}$ is a rational number.

v) 5.012896896896...

This decimal number is non-terminating and repeating (i.e., the group of digits 896 repeats after the decimal point). Hence, it is a rational number.

vi) 7.1931931934...

This decimal number appears to be non-terminating and repeating, but it is in fact non-terminating and non-repeating. The group of digits '193' repeats twice after the decimal point and then a new set of digits '1934' can be seen. There is no one digit or group of digits that keeps repeating itself after the decimal point. Hence, this number is irrational.

vii) 0.0787887888...

This decimal number is non-terminating and non-repeating. In it, the 8s between the 7s keep increasing by one. Hence, this number is irrational.

Medium

Example 1:

Consider the given set of numbers.

$$A = \left\{ 2.6666..., \frac{2157}{625}, 0.181881888..., \frac{35}{16}, 0.14201421..., \frac{\sqrt{2}}{2\sqrt{8}}, \pi \right\}$$

In set A, there are a rational numbers and b irrational numbers. Show that $a^2 - b^2 = a + b$.

Solution:

2.6666... is non-terminating and repeating (i.e., the digit '6' is repeated after the decimal point). Hence, it is a rational number.

$\frac{2157}{625}$ is of the form $\frac{p}{q}$, where $p = 2157$ and $q = 625$ are integers and $q \neq 0$. Hence, it is a rational number.

0.181881888... is non-terminating and non-repeating. In it, the 8s between the 1s keep increasing by one. Hence, it is an irrational number.



$\frac{35}{16}$ is of the form $\frac{p}{q}$, where $p = 35$ and $q = 16$ are integers and $q \neq 0$. Hence, it is a rational number.

0.14201421... is non-terminating and non-repeating. Hence, it is an irrational number.

$\frac{\sqrt{2}}{2\sqrt{8}} = \frac{\sqrt{2}}{4\sqrt{2}} = \frac{1}{4}$, which is of the form $\frac{p}{q}$, where $p = 1$ and $q = 4$ are integers and $q \neq 0$. Hence, it is a rational number.

We know that π is an irrational number.

Thus, in set A, we have $a = 4$ and $b = 3$.

$$a^2 - b^2 = (a + b)(a - b)$$

$$\Rightarrow a^2 - b^2 = (a + b)(4 - 3)$$

$$\Rightarrow a^2 - b^2 = (a + b)(1)$$

$$\Rightarrow a^2 - b^2 = (a + b)$$

Decimal Expansions of Rational Numbers

The Need for Converting Rational Numbers into Decimals

A carpenter wishes to make a point on the **edge** of a wooden plank at 95 mm from any end. He has a centimeter tape, but how can he use that to mark the required point?



Simple! He should convert 95 mm into its corresponding centimeter value, i.e., 9.5 cm and then measure and mark the required length on the wooden plank.

This is just one of the many situations in life when we face the need to convert numbers into decimals. In this lesson, we will learn to convert rational numbers into decimals, observe the types of decimal numbers, and solve a few examples based on this concept.



Two rational numbers $\frac{a}{b}$ and $\frac{c}{d}$ are equal if and only if $ad = bc$.

Take, for example, the rational numbers $\frac{2}{4}$ and $\frac{3}{6}$. Let us see if they are equal or not.

Here, $a = 2$, $b = 4$, $c = 3$ and $d = 6$

Now, we have:

$$ad = 2 \times 6 = 12$$

$$bc = 4 \times 3 = 12$$

Since $ad = bc$, we obtain $\frac{2}{4} = \frac{3}{6}$.

Rational Numbers as Division of Integers

We know that the form $\frac{p}{q}$ represents the division of integer p by the integer q . By solving this division, we can find the decimal equivalent of the rational number $\frac{p}{q}$. Now, let us convert the numbers $\frac{5}{8}$, $\frac{4}{3}$ and $\frac{2}{7}$ into decimals using the long division method.



		$\begin{array}{r} 0.285714... \\ 7 \overline{) 2.000000} \\ \underline{14} \\ 60 \\ \underline{56} \\ 40 \\ \underline{35} \\ 50 \\ \underline{49} \\ 10 \\ \underline{7} \\ 30 \\ \underline{28} \\ 2 \end{array}$
$\begin{array}{r} 0.625 \\ 8 \overline{) 5.000} \\ \underline{48} \\ 20 \\ \underline{16} \\ 40 \\ \underline{40} \\ 0 \end{array}$	$\begin{array}{r} 1.33... \\ 3 \overline{) 4.00} \\ \underline{3} \\ 10 \\ \underline{9} \\ 10 \\ \underline{9} \\ 1 \end{array}$	

While the remainder is zero in the division of 5 by 8, it is not so in case of the other two divisions. Thus, we can get two different cases in the decimal expansions of rational numbers.

Observing the Decimal Expansions of Rational Numbers

We can get the following two cases in the decimal expansions of rational numbers.

Case I: When the remainder is zero

In this case, the remainder becomes zero and the quotient or decimal expansion terminates after a finite number of digits after the decimal point. For example, in the decimal

expansion of $\frac{5}{8}$, we get the remainder as zero and the quotient as 0.625.

Case II: When the remainder is never zero

In this case, the remainder never becomes zero and the corresponding decimal expansion

is non-terminating. For example, in the decimal expansions of $\frac{4}{3}$ and $\frac{2}{7}$, we see that the remainder never becomes zero and their corresponding quotients are **non-terminating decimals**.

When we divide 4 by 3 and 2 by 7, we get 1.3333... and 0.285714285714... as the respective quotients. In these decimal numbers, the digit '3' and the group of digits

'285714' get repeated. Therefore, we can write $\frac{4}{3} = 1.3333... = 1.\bar{3}$ and $\frac{2}{7} = 0.285714285714... = 0.\overline{285714}$. Here, the symbol $\overline{\quad}$ indicates the digit or group of digits that gets repeated.

Solved Examples

Example 1: Write the decimal expansion of $\frac{1237}{25}$ and find if it is terminating or non-terminating and repeating.

Solution: Here is the long division method to find the decimal expansion of $\frac{1237}{25}$.

$$\begin{array}{r} 49.48 \\ 25 \overline{)1237.00} \\ \underline{100} \\ 237 \\ \underline{225} \\ 120 \\ \underline{100} \\ 200 \\ \underline{200} \\ 0 \end{array}$$

Hence, the decimal expansion of $\frac{1237}{25}$ is 49.48. Since the remainder is obtained as zero, the decimal number is terminating.

Example 2:

Write the decimal expansion of $\frac{2358}{27}$ and find if it is terminating or non-terminating and repeating.

Solution:

Here is the long division method to find the decimal expansion of $\frac{2358}{27}$.

$$\begin{array}{r}
 87.33... \\
 27 \overline{) 2358.00} \\
 \underline{216} \\
 198 \\
 \underline{189} \\
 90 \\
 \underline{81} \\
 90 \\
 \underline{81} \\
 9...
 \end{array}$$

Hence, the decimal expansion of $\frac{2358}{27}$ is 87.33.... Since the remainder 9 is obtained again and again, the decimal number is non-terminating and repeating. The decimal number can also be written as $87.\bar{3}$.

Medium

Example 1:

Find the decimal expansion of each of the following rational numbers and write the nature of the same.

1. $\frac{65}{101}$

2. $\frac{923}{400}$

3. $\frac{37}{99}$

4. $\frac{67}{100}$

Solution:

$$\begin{array}{r}
 \text{i) } 101 \overline{) 65.000000} \\
 \underline{606} \\
 440 \\
 \underline{404} \\
 360 \\
 \underline{303} \\
 570 \\
 \underline{505} \\
 650 \\
 \underline{606} \\
 440 \\
 \underline{404} \\
 36
 \end{array}$$

We have $\frac{65}{101} = 0.64356435\ldots = \overline{0.6435}$

The group of digits '6435' repeats after the decimal point. Hence, the decimal expansion of the given rational number is non-terminating and repeating.

$$\begin{array}{r}
 \text{ii) } 400 \overline{) 923.0000} \\
 \underline{800} \\
 1230 \\
 \underline{1200} \\
 300 \\
 \underline{0} \\
 3000 \\
 \underline{2800} \\
 2000 \\
 \underline{2000} \\
 0
 \end{array}$$

We have $\frac{923}{400} = 2.3075$

Hence, the given rational number has a terminating decimal expansion.

$$\begin{array}{r} \text{iii) } 99 \overline{) 37.0000} \\ \underline{297} \\ 730 \\ \underline{693} \\ 370 \\ \underline{297} \\ 730 \\ \underline{693} \\ 37 \end{array}$$

We have $\frac{37}{99} = 0.3737\ldots = 0.\overline{37}$

The pair of digits '37' repeats after the decimal point. Hence, the decimal expansion of the given rational number is non-terminating and repeating.

$$\begin{array}{r} \text{iv) } 100 \overline{) 67.00} \\ \underline{600} \\ 700 \\ \underline{700} \\ 0 \end{array}$$

We have $\frac{67}{100} = 0.67$

Hence, the given rational number has a terminating decimal expansion.

Decimal Expansion of Irrational Numbers

Facing Irrational Numbers in Life

Anu has a piece of land in the backyard of her house. She calls a mason and tells him to construct a square-shaped kitchen garden covering an area of 7m^2 .



After a quick mental calculation, the mason replies that he cannot construct the garden as per Anu's specification. Why do you think the mason says this?

To answer this question, let us first see the calculation performed by the mason in his head. Anu wants a square garden with an area of 7m^2 .

Area of a square = Side \times Side

$$\Rightarrow (\text{Side})^2 = 7\text{ m}^2$$

$$\therefore \text{Side} = \sqrt{7}\text{ m}$$

Now, we can find only an approximate value of $\sqrt{7}$ as it is an irrational number. We cannot ascertain the exact value of an irrational number as its decimal expansion is non-terminating and non-repeating. This is why the mason says that it is not possible to obtain a square garden as per Anu's specification.

Let us learn more about irrational numbers and the method to find their decimal expansions.

Did You Know?

In the 1870s, two mathematicians named Cantor and Dedekind stated that every real number is represented by a unique point on the number line and every point on the number line represents a unique real number.

Know Your Scientist





Georg Ferdinand Ludwig Philipp Cantor (1845–1918) was a German mathematician. He is best known for the invention of 'the set theory', which went on to become 'the fundamental theory of mathematics'. Apart from being a teacher and researcher, he was also an outstanding violinist.



Julius Wilhelm Richard Dedekind (1831–1916) was a German mathematician. He is best known for his work in abstract algebra, algebraic number theory and, especially, the foundation of real numbers.

Irrationality of Square Roots of Non Perfect Square Numbers

We know that the decimal expansion of an irrational number is non-terminating and non-repeating.

Take, for example, the irrational number $\sqrt{2}$. Let us find its decimal expansion.



	1.41421
1	2.0000000000
	1
24	100
	96
281	400
	281
2824	11900
	11296
28282	60400
	56564
282841	383600
	282841
	100759

The decimal expansion of $\sqrt{2}$ is 1.41421.... It is clearly non-terminating and non-repeating. So, $\sqrt{2}$ is an irrational number.

Decimal expansions of few more numbers are given below:

$$\sqrt{7} = 2.64575...; \sqrt{10} = 3.16227...; \sqrt{65} = 8.06225...$$

In all cases, we found an irrational number.

It can be observed that 2, 7, 10 and 65 all are non perfect squares and their square roots are irrational numbers.

So, it can be concluded that **the square roots of non perfect square numbers are irrational numbers.**

Solved Examples

Easy

Example 1: Prove that $\sqrt{3}$ is an irrational number.

Solution:

	1.7320508
1	3.00000000000000
	1
27	200
	189
343	1100
	1029
3462	7100
	6924
346405	1760000
	1732025
34641008	279750000
	277128064
	2621936

The decimal expansion of $\sqrt{3}$ is 1.7320508.... It is clearly non-terminating and non-repeating. So, $\sqrt{3}$ is an irrational number.


Irrationality of n^{th} Root of Numbers

Till now we have studied that the square root of a non-perfect square number is irrational. But what about numbers in which the root is of an order greater than 2? Take, for example, the number $\sqrt[3]{6}$. In this number, the order of the root is 3. Since it is not possible to find the cube root of 6 as a terminating or repeating decimal number, $\sqrt[3]{6}$ is an irrational number. We can conclude similarly about other numbers like $\sqrt[4]{10}$ and $\sqrt[8]{25}$.

The irrationality of such numbers can be stated as follows:

The number $\sqrt[n]{a}$ or $a^{\frac{1}{n}}$ is irrational if it is not possible to represent a in the form b^n , where b is a factor of a .

Know Your Scientist



Plato (428/427 BC–348/347 BC) was classical Greek philosopher and mathematician. He was the founder of the Academy in Athens, which was the first institution of higher learning in the western world. Plato's dialogues have been used in various subjects, including philosophy, logic, ethics, rhetoric and mathematics.

Solved Examples

Example 1: Among the numbers $\sqrt{5}$, $\sqrt{4}$, $\sqrt[5]{12}$ and $\sqrt{7}$, which is/are irrational?

Solution:

We know that \sqrt{a} is an irrational number when 'a' is not a perfect square.

Among the given numbers, 4 is a perfect square. So, $\sqrt{4}$ is a rational number.

On the other hand, 5 and 7 are not perfect squares. Thus, $\sqrt{5}$ and $\sqrt{7}$ are irrational numbers.

$\sqrt[5]{12}$ is also irrational as 12 cannot be written in the form b^6 where b is a factor of 12.

Representing Rational Numbers in Non-terminating Recurring Decimal Form

Rational numbers have terminating decimal expansions, but they can be represented in the form of non-terminating recurring decimal numbers as well.

For example, $\frac{5}{2}$ is a rational number as it is written in the form of $\frac{p}{q}$ where $q \neq 0$.

Now, $\frac{5}{2} = 2.5$ which is a terminating decimal number. 2.5 can also be written as 2.5000... or $2.\overline{50}$.

Thus, we get non-terminating recurring decimal form for the rational number $\frac{5}{2}$.

Similarly, we can represent each rational number in non-terminating recurring decimal form.

Conversely, it can be said that every number in the non-terminating recurring decimal form is a rational number.



Solved Examples

Easy

Example 1:

Represent the following rational numbers in non-terminating recurring decimal form.

a. $\frac{15}{4}$

b. 2.007

c. 0.02

Solution:

a. The given number is $\frac{15}{4}$. It can be written in the non-terminating recurring decimal form as follows:

$$\frac{15}{4} = 3.75 = 3.75000... = 3.75\overline{0}$$

b. The given number is 2.007. It can be written in the non-terminating recurring decimal form as follows:

$$2.007 = 2.007000... = 2.007\overline{0}$$

c. The given number is 0.02. It can be written in the non-terminating recurring decimal form as follows:

$$0.02 = 0.02000... = 0.02\overline{0}$$

Conversion of Decimals into Rational Numbers

The Need for Converting Decimals into Fractions

In daily life, situations can arise wherein we have to separate a part of a whole for some purpose. Say, for example, a farmer wishes to give $(0.3333...)^\text{th}$ part of his three-hectare land to his eldest son.





Now, $0.3333\dots$ is not a terminating decimal. It is not possible to measure this decimal part of three hectares. However, if this non-terminating decimal is expressed as

a **fraction** $\left(\text{i.e., } \frac{1}{3}\right)$, then the calculation becomes a lot easier and more accurate. It is now clear that the farmer wants to give his eldest son one-third of his land, i.e., one hectare out of the three hectares of land.

In this lesson, we will learn how to convert decimals into fractions.

Converting a Terminating Decimal into a Fraction

Solved Examples

Example 1: Express the number 2.25 in the form $\frac{p}{q}$.

Solution:

Let $x = 2.25$

There are two digits after the decimal **point**; so, we can write:

$$x = \frac{225}{10^2} = \frac{225}{100}$$

On simplification, we obtain:

$$x = \frac{225}{100} = \frac{25 \times 9}{25 \times 4} = \frac{9}{4}$$

Thus, the number 2.25 can be written in the form $\frac{p}{q}$ as $\frac{9}{4}$.

Converting a Non-Terminating Repeating Decimal into a Fraction

Solved Examples

Easy

Example 1: Express each of the following decimals as a fraction or a rational number.

1. $0.\overline{7}$
2. $0.\overline{29}$

Solution:

1. Let $x = 0.\overline{7}$

$$\Rightarrow x = 0.7777 \dots \dots (1)$$

We have only one repeating digit after the decimal point. On multiplying equation 1 with $10^1 = 10$, we get:

$$10x = 7.7777 \dots (2)$$

On subtracting equation 1 from equation 2, we get:

$$9x = 7$$

$$\Rightarrow x = \frac{7}{9}$$

$$\therefore 0.\overline{7} = \frac{7}{9}$$

2. Let $x = 0.\overline{29}$

$$\Rightarrow x = 0.292929 \dots \dots (1)$$

We have two repeating digits after the decimal point. On multiplying equation 1 with $10^2 = 100$, we get:

$$100x = 29.2929 \dots \dots (2)$$

On subtracting equation 1 from equation 2, we get:

$$99x = 29$$

$$\Rightarrow x = \frac{29}{99}$$

$$\therefore 0.\overline{29} = \frac{29}{99}$$

Hard

Example 1: Find the rational forms of the following numbers.

1. $2.\overline{35961}$
2. $0.\overline{059}$

Solution:

1. Let $x = 2.\overline{35961}$

$$x = 2.35961961... \dots (1)$$

On multiplying both sides of equation 1 with 100, we obtain:

$$100x = 235.961961961... \dots (2)$$

On multiplying both sides of equation 2 with 1000, we obtain:

$$100000x = 235961.961961961... \dots (3)$$

On subtracting equation 2 from equation 3, we obtain:

$$99900x = 235726$$

$$\Rightarrow x = \frac{235726}{99900}$$

$$\Rightarrow x = \frac{117863}{49950}$$

Thus, the rational form of $2.\overline{35961}$ is $\frac{117863}{49950}$.

2. Let $x = 0.\overline{059}$

$$x = 0.0595959... \dots (1)$$

On multiplying both sides of equation 1 with 10, we obtain:

$$10x = 0.595959... \dots (2)$$

On multiplying both sides of equation 2 with 100, we obtain:

$$1000x = 59.595959... \dots (3)$$

On subtracting equation 2 from equation 3, we obtain:

$$990x = 59$$


$$\Rightarrow x = \frac{59}{990}$$

Thus, the rational form of $0.\overline{059}$ is $\frac{59}{990}$.

Finding Irrational Numbers between Given Rational Numbers

Real Numbers between Any Two Numbers

Consider the given number line.



This number line shows integers from -10 to 10 . While there are clearly 19 integers between -10 and 10 , the number of fractions between the two integers is infinite. So, we can say that there are infinite real numbers between any two numbers. Real numbers include both rational and irrational numbers. So we can say that there are infinite rational and irrational numbers between any two numbers.

In this lesson, we will learn to find irrational numbers between any two rational or irrational numbers.

Finding Irrational Numbers between Pairs of Rational Numbers

We know that between any two numbers, there are infinite rational and irrational numbers.

Let us learn to find irrational numbers between any two rational numbers. Here are the steps to do the same.



Step 1: Find the decimal representation (up to 2 or 3 places of decimal) of the two given rational numbers. Let those decimal representations be a and b , such that $a < b$.

Step 2: Choose the required non-terminating and non-repeating decimal numbers (i.e., irrational numbers) between a and b .

Similarly, we can find irrational numbers between other pairs of rational numbers.

Finding Irrational Numbers between Pairs of Rational Numbers

Solved Examples

Example 1: Find an irrational number between 0.12 and 0.15.

Solution:

Step 1: The rational numbers are already in the decimal form.

Let $a = 0.12$ and $b = 0.15$.

We can see that $a < b$.

Step 2: There is an infinite number of irrational numbers between a and b .

Clearly, the non-terminating and non-repeating number 0.12101001000... lies between 0.12 and 0.15. Hence, it is the required irrational number.

Example 1: Find five irrational numbers between $\frac{1}{3}$ and $\frac{2}{5}$.

Solution: Step 1: The decimal expansion of $\frac{1}{3}$ is 0.333...

The decimal expansion of $\frac{2}{5}$ is 0.4 or 0.400.

We can see that $0.333 < 0.4$.

Step 2: Five irrational numbers between 0.333 and 0.400 are listed below.

0.34560561562563...

0.3574744744474444...

0.369874562301...

0.3710110111011110...

0.39919293...

Representation of Rational Numbers on Number Line Using Successive Magnification

Introduction to Magnification

There are several things around us which are so small that they cannot be seen clearly with the naked eye; for example, the eyes of ants and small insects. These minute items can only be observed with the help of a lens that makes them appear bigger to the eye.



This process of making a thing seem bigger without actually changing its physical size is known as **magnification**. Generally, a magnifying glass is used for such enlargement.

Similarly, on the **number line**, there are infinite smaller numbers lying between any two numbers. These smaller numbers can be of two, three or more decimal places. To see or mark such numbers clearly, we use the process called **successive magnification of the number line**. Here, we use a virtual (imaginary) magnifying glass to enlarge the smaller divisions on the number line

In this lesson, we will learn to represent rational numbers on the number line using the successive magnification method.

Successive Magnification of the Number Line

Watch this video to understand the need for the successive magnification of the number line.

Representing Numbers Using the Successive Magnification Method

Solved Examples



Medium

Example 1: Visualize $6.\overline{285}$ on the number line, up to five decimal places.

Solution:

$6.\overline{285}$ can be expressed up to five decimal places as 6.28585. Here are the steps to visualize this number on the number line.

Step 1: 6.28585 lies between 6 and 7. Divide the number line between 6 and 7 into ten equal parts and magnify the distance between them.

Step 2: 6.28585 lies between 6.2 and 6.3. Divide the number line between 6.2 and 6.3 into ten equal parts and magnify the distance between them.


Step 3: 6.28585 lies between 6.28 and 6.29. Divide the number line between 6.28 and 6.29 into ten equal parts and magnify the distance between them.

Step 4: 6.28585 lies between 6.285 and 6.286. Divide the number line between 6.285 and 6.286 into ten equal parts and magnify the distance between them.

Step 5: 6.28585 lies between 6.2858 and 6.2859. Divide the distance between 6.2858 and 6.2859 into ten equal parts and magnify the distance between them. We can now mark 6.28585 on the number line.

The figures obtained at the end of each step are shown.






Representation of Irrational Numbers on Number Line

Square Root Spiral


Consider the given square root spiral.



This spiral is obtained by geometrical representation of square roots of successive natural numbers 1, 2, 3..., and so on. We can construct this square root spiral if we know how to represent irrational numbers of the form \sqrt{n} on the number line, where n is any positive integer. Since irrational numbers are non-terminating and non-repeating, they are represented on the number line by a special method involving Pythagoras theorem.

In this lesson, we will learn to represent irrational numbers of the form \sqrt{n} on the number line.

Know Your Scientist



Pythagoras (570 BC–495 BC) was a great Greek mathematician and philosopher, often described as the first pure mathematician. He was born on the island of Samos and is best known for the Pythagoras theorem about right-angled triangles. He also made influential contributions to philosophy and religious teaching.


Representing Irrational Numbers of the Form \sqrt{n} on the Number Line

Concept Builder

Pythagoras theorem

In a right-angled triangle, the square of the hypotenuse is equal to the sum of the squares of the other two sides.

Consider the following right-angled triangle PQR.



ΔPQR is right-angled at Q, i.e., $\angle PQR = 90^\circ$. By Pythagoras theorem, we have:

$$\text{Hypotenuse}^2 = \text{Base}^2 + \text{Perpendicular}^2$$

$$\Rightarrow PR^2 = RQ^2 + PQ^2$$

Solved Examples

Example 1: Represent $\sqrt{5}$ on the number line.

Solution:

Here are the steps to represent $\sqrt{5}$ on the number line.

Step 1: Draw a line and mark the integers $-2, -1, 0, 1, 2$, etc. on it, so that the distance between any two consecutive integers is one unit.

Step 2: Mark points O and A at 0 and 1 respectively. From A, draw a line segment AB of unit length and perpendicular to OA. Join O to B to get $\triangle OBA$. Thus, by Pythagoras theorem, we have $OB = \sqrt{2}$.


Step 3: From B, draw a line segment BC of unit length and perpendicular to OB. Join O to C to get $\triangle OCB$. Thus, by Pythagoras theorem, we have $OC = \sqrt{3}$.

Step 4: From C, draw a line segment CD of unit length and perpendicular to OC. Join O to D to get $\triangle ODC$. Thus, by Pythagoras theorem, we have $OD = \sqrt{4}$.

Step 5: From D, draw a line segment DE of unit length and perpendicular to OD. Join O to E to get $\triangle OED$. Thus, by Pythagoras theorem, we have $OE = \sqrt{5}$.

Step 6: Now, taking O as the centre and OE as the radius, draw an arc that cuts the number line at point F. This point represents the irrational number $\sqrt{5}$ on the number line.


The figure for the construction is shown below.



Geometrical Representation of \sqrt{n} when $(n - 1)$ is a Perfect Square

We have seen that to represent the square root of a number on the number line, the square root of its predecessor is used as the base of a right-angled triangle. So, if the number $(n - 1)$ is a perfect square, then we can directly find its square root and use it to represent \sqrt{n} on the number line. Let us represent $\sqrt{5}$ on the number line using this method.

We have $\sqrt{5-1} = \sqrt{4} = 2$. So, we will consider a base of 2 units on the number line. To represent this base, mark points 0 and 2 as O and A respectively. From A, draw a perpendicular AB of unit length. Join O to B to get $\triangle OBA$. Then, taking O as the centre and OB as the radius, draw an arc cutting the number line at point P. Point P represents the irrational number $\sqrt{5}$ on the number line.



Let us verify our construction.

Using Pythagoras theorem in $\triangle OBA$, we obtain:

$$OB^2 = OA^2 + AB^2$$

$$\Rightarrow OB^2 = 2^2 + 1^2 = 4 + 1$$

$$\Rightarrow OB = \sqrt{5}$$

$$\therefore OP = OB = \sqrt{5} \quad (\because OP \text{ and } OB \text{ are radii of the same arc})$$

In this way, we can represent other irrational numbers on the number line.


Geometrical Representation of \sqrt{n} when $(n - 1)$ is a Perfect Square

Solved Examples

Example 1: Represent $\sqrt{17}$ on the number line and justify the construction.

Solution:

We know that $17 = 4^2 + 1^2$. So, mark points 0 and 4 as O and A respectively. From A, draw a perpendicular AB of 1 unit. Join O to B to get $\triangle OBA$. Then, take O as the centre and OB as the radius, and draw an arc that intersects the number line at point P. This point represents $\sqrt{17}$ on the number line. The construction is shown below.



The construction can be verified as is shown.

We have $OA = 4$ units and $AB = 1$ unit. Using Pythagoras theorem in $\triangle OAB$, we have:

$$OB^2 = OA^2 + AB^2$$

$$\Rightarrow OB^2 = 4^2 + 1^2$$

$$\Rightarrow OB^2 = 16 + 1 = 17$$

$$\Rightarrow OB = \sqrt{17}$$

$$\therefore OP = OB = \sqrt{17} \quad (\because OP \text{ and } OB \text{ are radii of the same arc})$$

Geometrical Representation of the Square Root of a Given Positive Real Number

Difficulty in Representing \sqrt{n} for a Positive Real Number on the Number Line

Suppose we have to represent $\sqrt{5.6}$ on the number line. We know that $\sqrt{5.6}$ is irrational since we cannot find a terminating or repetitive decimal number x such that $x^2 = 5.6$.

We know how to represent \sqrt{n} , for any integer n , on the number line. For this, we locate $\sqrt{n-1}$ first and then \sqrt{n} . So, to locate $\sqrt{5.6}$ on the number line by the same method, we have to first locate $\sqrt{4.6}$. Now, $\sqrt{4.6}$ is also an irrational number; to locate it on the number line, we have to first locate $\sqrt{3.6}$, which is again an irrational number. It is clear that this method is not helpful in locating irrational numbers like $\sqrt{5.6}$ on the number line. We use another method to represent such numbers on the number line. Go through this lesson to learn about the same.

Concept Builder

Square and square root

When a number is multiplied with itself, the obtained product is the square of the original number. Say, x is any number and $x \times x = y$. In this case, the number y is the square of x . In exponential form, $x \times x$ is written as x^2 . This is called 'square of x ', ' x square' or ' x raised to the power 2'.

Now, we have assumed that $x^2 = y$. So, x will be equal to the square root of y , which is represented as \sqrt{y} . Here, $\sqrt{\quad}$ is the symbol of square root. Finding the square root of a number means representing the number as the product of another number multiplied with itself. It is the reverse of finding the square of a number. We have assumed y to be the product of $x \times x$; so, $\sqrt{y} = x$.


Locating $\sqrt{5.6}$ on the Number Line

Concept Builder

Pythagoras theorem

In a right-angled triangle, the square of the hypotenuse is equal to the sum of the squares of the other two sides.

Consider the following right-angled triangle PQR.



$\triangle PQR$ is right-angled at Q, i.e., $\angle PQR = 90^\circ$. By Pythagoras theorem, we have:

$$\text{Hypotenuse}^2 = \text{Base}^2 + \text{Perpendicular}^2$$

$$\Rightarrow PR^2 = RQ^2 + PQ^2$$

Know More

Square root

- The square root of every non-negative real number is a unique non-negative real number.
- The square root of 0 is 0.
- The square root of a negative real number is not a real number.

Solved Examples

Medium


Example 1: Represent $\sqrt{6.7}$ on the number line and justify the construction.

Solution: Here are the steps to locate $\sqrt{6.7}$ on the number line.

Step 1: Draw a line and mark a point A on it. Mark points B and C such that AB = 6.7 units and BC = 1 unit.

Step 2: Find the midpoint of AC and mark it as M. Taking M as the centre and MA as the radius, draw a semicircle.

Step 3: From B, draw a perpendicular to AC and let it meet the semicircle at point D. Taking B as the centre and BD as the radius, draw an arc that intersects the line at point E.



Now, the distance BE is $\sqrt{6.7}$ units.

Verification of construction:

We have AB = 6.7 units and BC = 1 unit

$$\therefore AC = AB + BC = 6.7 \text{ units} + 1 \text{ unit} = 7.7 \text{ units}$$

$$\therefore MA = \frac{1}{2} AC = \frac{1}{2} \times 7.7 \text{ units} = 3.85 \text{ units}$$

$$MB = AB - MA = 6.7 \text{ units} - 3.85 \text{ units} = 2.85 \text{ units}$$

Also, $MA = MD = 3.85 \text{ units}$ (\because MA and MD are the radii of the same circle)

On applying Pythagoras theorem in $\triangle MBD$, we obtain:

$$BD^2 = MD^2 - MB^2 = 3.85^2 - 2.85^2 = 14.8225 - 8.1225 = 6.7$$

$$\Rightarrow \therefore BD = \sqrt{6.7} \text{ units}$$

Hence, we get

$$BE = BD = \sqrt{6.7} \text{ units} (\because BE \text{ and } BD \text{ are the radii of the same circle})$$

Thus, our construction is justified.

Operations on Irrational Numbers

Mathematical Operations and Irrational Numbers

We have learnt to perform addition, subtraction, multiplication and division on integers, decimals and fractions. We can also perform these operations on irrational numbers of the form \sqrt{n} , where n is a positive real number.

Performing mathematical operations on irrational numbers is similar to performing these operations on **algebraic expressions**. For example, to add the algebraic expressions $2xy + 3y^2$ and $x^2y - 4xy$, we first check and add the like terms and then write the unlike terms as they are. In our example, $2xy$ and $-4xy$ are like terms as they have the common algebraic part xy .

$$\text{So, } (2xy + 3y^2) + (x^2y - 4xy) = (2xy - 4xy) + 3y^2 + x^2y = -2xy + 3y^2 + x^2y$$

Irrational numbers are also categorized as like and unlike irrational numbers. We can add or subtract like irrational numbers only.

In this lesson, we will learn how to perform the four mathematical operations on irrational numbers.

Like terms: The terms or numbers whose irrational parts are the same are known as like terms. For example, $\frac{2}{7}\sqrt{3}$ and $\frac{3}{5}\sqrt{3}$ are like terms because the irrational parts in these numbers are the same, i.e., $\sqrt{3}$.

Unlike terms: The terms or numbers whose irrational parts are not the same are known as unlike terms. For example, $\frac{31}{8}\sqrt{7}$ and $\frac{11}{13}\sqrt{5}$ are unlike terms because the irrational parts in these numbers are different, i.e., $\sqrt{7}$ and $\sqrt{5}$.

Sometimes, two numbers may appear to have different irrational parts, but on simplification they are found to be the same. For example, $5\sqrt{2}$ and $3\sqrt{8}$ seem to have different irrational parts, i.e., $\sqrt{2}$ and $\sqrt{8}$. However, on simplifying $3\sqrt{8}$, we get $3\sqrt{8} = 3(2\sqrt{2}) = 6\sqrt{2}$. Thus, we see that the two numbers have the same irrational part, i.e., $\sqrt{2}$.

Arithmetic Operations between Rational and Irrational Numbers

We have learnt to perform operations between fractions and integers, decimals and whole numbers and different types of numbers. Now, let us try to perform the same between rational and irrational numbers.

Let us take the rational number 4 and the irrational number $\sqrt{7}$. On applying the four operations on these numbers, we get $4 + \sqrt{7}$, $4 - \sqrt{7}$, $4 \times \sqrt{7}$ and $\frac{4}{\sqrt{7}}$.

Since $\sqrt{7}$ has a non-terminating and non-repeating decimal expansion, the decimal expansions of $4 + \sqrt{7}$, $4 - \sqrt{7}$, $4 \times \sqrt{7}$ and $\frac{4}{\sqrt{7}}$ will also be non-terminating and non-repeating. Hence, these numbers will also be irrational.

So, we can conclude that:

- The sum or difference of a rational and an irrational number is always irrational.
- The product or quotient of a non-zero rational number and an irrational number is always irrational.

Solved Examples

Example 1: Check whether $\pi + 8$ and $\frac{4\sqrt{3}-6}{5}$ are irrational numbers or not.

Solution:

We know that π is an irrational number and 8 is a rational number. The sum of a rational and an irrational number is always irrational. Hence, $\pi + 8$ is an irrational number. It can be proved as follows:

$$\pi = 3.1415...$$

$$\Rightarrow \pi + 8 = 3.1415... + 8 = 11.1415...$$

11.1415... is a non-terminating and non-repeating decimal number, so it is irrational.

We know that $\sqrt{3}$ is an irrational number and 4 is rational number. The product of a non-zero rational number and an irrational number is always irrational. Thus, $4\sqrt{3} = 4 \times \sqrt{3}$ is an irrational number. Similarly, $4\sqrt{3} - 6$ is an irrational number as it is the difference

between a rational and an irrational number. Finally, $\frac{4\sqrt{3}-6}{5}$ is an irrational number as it is the quotient of a non-zero rational number and an irrational number.

Performing Operations on Irrational Numbers

The decimal expansion of an irrational number is non-terminating and non-repeating. For this reason, unlike irrational terms cannot be added or subtracted.

Suppose \sqrt{x} and \sqrt{y} are two unlike irrational numbers. The arithmetic operations between them are shown as follows:

- Addition = $\sqrt{x} + \sqrt{y}$
- Subtraction = $\sqrt{x} - \sqrt{y}$ or $\sqrt{y} - \sqrt{x}$
- Multiplication = $\sqrt{x} \times \sqrt{y} = \sqrt{x \times y} = \sqrt{xy}$
- Division = $\sqrt{x} \div \sqrt{y} = \frac{\sqrt{x}}{\sqrt{y}} = \sqrt{\frac{x}{y}}$

Suppose $a\sqrt{x}$ and $b\sqrt{x}$ are two like irrational numbers. The arithmetic operations between them are shown as follows:

- Addition = $a\sqrt{x} + b\sqrt{x} = a \times \sqrt{x} + b \times \sqrt{x} = (a+b) \times \sqrt{x} = (a+b)\sqrt{x}$
- Subtraction = $a\sqrt{x} - b\sqrt{x} = a \times \sqrt{x} - b \times \sqrt{x} = (a-b) \times \sqrt{x} = (a-b)\sqrt{x}$
- Multiplication = $a\sqrt{x} \times b\sqrt{x} = ab\sqrt{x \times x} = ab\sqrt{x^2} = abx$
- Division = $\frac{a\sqrt{x}}{b\sqrt{x}} = \frac{a \times \sqrt{x}}{b \times \sqrt{x}} = \frac{a}{b} \times \frac{\sqrt{x}}{\sqrt{x}} = \frac{a}{b}$

Solved Examples

Easy

Example 1:

1. Divide $8\sqrt{21}$ by $2\sqrt{7}$.
2. Multiply $8\sqrt{21}$ with $2\sqrt{7}$.

Solution:

$$\begin{aligned} \text{i) } & \frac{8\sqrt{21}}{2\sqrt{7}} \\ &= \frac{8\sqrt{7 \times 3}}{2\sqrt{7}} \\ &= \frac{8\sqrt{7} \times \sqrt{3}}{2\sqrt{7}} \\ &= 4\sqrt{3} \end{aligned}$$

$$\begin{aligned} \text{ii) } & 8\sqrt{21} \times 2\sqrt{7} = 8 \times \sqrt{3 \times 7} \times 2 \times \sqrt{7} \\ &= 8 \times 2 \times \sqrt{7} \times \sqrt{7} \times \sqrt{3} \\ &= 16 \times \sqrt{7 \times 7} \times \sqrt{3} \\ &= 16 \times 7\sqrt{3} \\ &= 112\sqrt{3} \end{aligned}$$

Example 2:

1. Prove that $\sqrt{24} + \sqrt{54} = \sqrt{150}$.

- Subtract $\sqrt{125}$ from $2\sqrt{5}$.

Solution:

- i) We have to prove that $\sqrt{24} + \sqrt{54} = \sqrt{150}$.

$$\begin{aligned}
 \text{LHS} &= \sqrt{24} + \sqrt{54} \\
 &= \sqrt{2 \times 2 \times 2 \times 3} + \sqrt{2 \times 3 \times 3 \times 3} \\
 &= 2\sqrt{2 \times 3} + 3\sqrt{2 \times 3} \\
 &= 2\sqrt{6} + 3\sqrt{6} \\
 &= 5\sqrt{6} \\
 \text{RHS} &= \sqrt{150} \\
 &= \sqrt{2 \times 3 \times 5 \times 5} \\
 &= 5\sqrt{6} \\
 \therefore \text{LHS} &= \text{RHS}
 \end{aligned}$$

$$\begin{aligned}
 \text{ii) } 2\sqrt{5} - \sqrt{125} \\
 &= 2\sqrt{5} - \sqrt{5 \times 5 \times 5} \\
 &= 2\sqrt{5} - 5\sqrt{5} \\
 &= (2 - 5)\sqrt{5} \\
 &= -3\sqrt{5}
 \end{aligned}$$

Example 1: Simplify $\frac{\sqrt{20} \times 2\sqrt{12}}{6\sqrt{15}}$.

Solution:

$$\begin{aligned}
 & \frac{\sqrt{20} \times 2\sqrt{12}}{6\sqrt{15}} \\
 &= \frac{\sqrt{2 \times 2 \times 5} \times 2\sqrt{2 \times 2 \times 3}}{6\sqrt{3 \times 5}} \\
 &= \frac{2\sqrt{5} \times 2 \times 2\sqrt{3}}{6\sqrt{3} \times \sqrt{5}} \\
 &= \frac{8}{6} \times \frac{\sqrt{5} \times \sqrt{3}}{\sqrt{5} \times \sqrt{3}} \\
 &= \frac{4}{3}
 \end{aligned}$$

Concept Builder

There is an order in which calculations should be performed while simplifying an expression. This order of performing operations is called BODMAS, with each letter in this word standing for a particular operation.

B	O	D	M	A	S
----------	----------	----------	----------	----------	----------

Brackets Of Division Multiplication Addition Subtraction

While simplifying an expression, we should first remove the '**brackets**'. Next, we should perform operations involving '**of**', e.g., one fourth of 16, 20% of 100, etc. Then, we should carry out '**division**', '**multiplication**', '**addition**' and '**subtraction**', in that order.

All expressions are solved using the **BODMAS rule**. Take, for example, the expression $36 \div 12 + 7 \times 2 - 7$. We simplify this expression as follows:

$$36 \div 12 + 7 \times 2 - 7$$

$$= 3 + 7 \times 2 - 7 \text{ (Division)}$$

$$= 3 + 14 - 7 \text{ (Multiplication)}$$

$$= 17 - 7 \text{ (Addition)}$$

$$= 10 \text{ (Subtraction)}$$

Solved Examples

Easy

Example 1: Multiply $(3\sqrt{7} + 2\sqrt{13})$ with $\sqrt{5}$.

Solution:

$$\begin{aligned} & (3\sqrt{7} + 2\sqrt{13}) \times \sqrt{5} \\ &= 3\sqrt{7} \times \sqrt{5} + 2\sqrt{13} \times \sqrt{5} \\ &= 3\sqrt{7 \times 5} + 2\sqrt{13 \times 5} \\ &= 3\sqrt{35} + 2\sqrt{65} \end{aligned}$$

Example 2: Simplify $\frac{2\sqrt{150} - 9}{\sqrt{3}}$.

Solution:

$$\begin{aligned} & \frac{2\sqrt{150} - 9}{\sqrt{3}} \\ &= \frac{2\sqrt{2 \times 3 \times 5 \times 5} - 3 \times 3}{\sqrt{3}} \\ &= \frac{2 \times 5 \times \sqrt{2} \times \sqrt{3} - 3 \times \sqrt{3} \times \sqrt{3}}{\sqrt{3}} \\ &= \frac{\sqrt{3}(10\sqrt{2} - 3\sqrt{3})}{\sqrt{3}} \\ &= 10\sqrt{2} - 3\sqrt{3} \end{aligned}$$

Medium

Example 1: Simplify $(5 + \sqrt{7})(\sqrt{7} - 2) + (3 + \sqrt{3})^2$.

Solution:



$$\begin{aligned}
& (5 + \sqrt{7})(\sqrt{7} - 2) + (3 + \sqrt{3})^2 \\
&= (5 + \sqrt{7})(\sqrt{7} - 2) + (3 + \sqrt{3})(3 + \sqrt{3}) \\
&= 5 \times \sqrt{7} - 5 \times 2 + \sqrt{7} \times \sqrt{7} - \sqrt{7} \times 2 + 3 \times 3 + 3 \times \sqrt{3} + \sqrt{3} \times 3 + \sqrt{3} \times \sqrt{3} \\
&= 5\sqrt{7} - 10 + 7 - 2\sqrt{7} + 9 + 6\sqrt{3} + 3 \\
&= 3\sqrt{7} + 6\sqrt{3} + 9
\end{aligned}$$

Example 2: Simplify $(\sqrt{486} \div 2) - (\sqrt{27} \div \sqrt{2})$.

Solution:

$$\begin{aligned}
& (\sqrt{486} \div 2) - (\sqrt{27} \div \sqrt{2}) \\
&= \frac{\sqrt{486}}{2} - \frac{\sqrt{27}}{\sqrt{2}} \\
&= \frac{\sqrt{486}}{2} - \sqrt{\frac{27}{2}} \\
&= \frac{9\sqrt{6}}{2} - 3\sqrt{\frac{3}{2}} \\
&= \frac{9\sqrt{6}}{\sqrt{4}} - 3\sqrt{\frac{3}{2}} \\
&= 9\sqrt{\frac{6}{4}} - 3\sqrt{\frac{3}{2}} \\
&= 9\sqrt{\frac{3}{2}} - 3\sqrt{\frac{3}{2}} \\
&= 6\sqrt{\frac{3}{2}} \\
&= 3 \times 2\sqrt{\frac{3}{2}} \\
&= 3 \times 2 \frac{\sqrt{3}}{\sqrt{2}} \times \frac{\sqrt{2}}{\sqrt{2}} \\
&= 3 \times \frac{2\sqrt{6}}{2} \\
&= 3\sqrt{6}
\end{aligned}$$

Closure Property of Irrational Numbers

Introduction to Closure Property

In order to study the closure property of irrational numbers, we first need to know the meaning of the term 'closure property'. For this purpose, let us consider the two integers -5 and 8 . The sum of these integers is 3 and the difference when 8 is subtracted from -5 is -13 . You can see that the sum and the difference are both integers. What this tells us is that adding or subtracting two integers always gives an integer as the result. In other words, the sum or difference of integers is closed to be an integer. Thus, we can say that: integers are 'closed' with respect to addition and subtraction, or integers satisfy the closure property under addition and subtraction.

In this lesson, we will discuss the closure property of irrational numbers under different algebraic operations such as addition, subtraction, multiplication and division.

Know More

Division by 0:

Division means to divide the dividend (numerator) into as many equal parts as the divisor (denominator). For example, $4 \div 2$ means that we need to divide the number 4 into two equal parts.

Division by 0 means dividing a whole into zero equal parts. There may be infinite number of ways in which we can divide that whole into unequal parts or zero equal parts. Thus, division by 0 is not defined.

Solved Examples

Example 1: State two irrational numbers whose:

1. sum is a rational number
2. sum is an irrational number

Solution:

1. Let us consider the two irrational numbers $2 + \sqrt{3}$ and $2 - \sqrt{3}$.

The sum of these two numbers is found as follows:

$$\begin{aligned} & (2 + \sqrt{3}) + (2 - \sqrt{3}) \\ &= 2 + \sqrt{3} + 2 - \sqrt{3} \\ &= 4 \text{ (a rational number)} \end{aligned}$$

2. Let us consider the two irrational numbers $\sqrt{5} + 3$ and $\sqrt{5} - 3$.



The sum of these two numbers is found as follows:

$$\begin{aligned} & (\sqrt{5} + 3) + (\sqrt{5} - 3) \\ &= \sqrt{5} + 3 + \sqrt{5} - 3 \\ &= 2\sqrt{5} \text{ (an irrational number)} \end{aligned}$$

Example 2: State two irrational numbers whose:

1. difference is a rational number
2. difference is an irrational number

Solution:

1. Let us consider the two irrational numbers $\sqrt{2} + 3$ and $\sqrt{2} - 5$.

The difference between these two numbers is found as follows:

$$\begin{aligned} & (\sqrt{2} + 3) - (\sqrt{2} - 5) \\ &= \sqrt{2} + 3 - \sqrt{2} + 5 \\ &= 8 \text{ (a rational number)} \end{aligned}$$

2. Let us consider the two irrational numbers $\frac{2}{5} - \sqrt{6}$ and $\frac{8}{5} - 2\sqrt{6}$.

The difference between these two numbers is found as follows:

$$\begin{aligned} & \left(\frac{2}{5} - \sqrt{6}\right) - \left(\frac{8}{5} - 2\sqrt{6}\right) \\ &= \frac{2}{5} - \sqrt{6} - \frac{8}{5} + 2\sqrt{6} \\ &= -\frac{6}{5} + \sqrt{6} \text{ (an irrational number)} \end{aligned}$$

Medium

Example 1: State two irrational numbers whose:

1. product is a rational number
2. product is an irrational number

Solution:

1. Let us consider the two irrational numbers $7+4\sqrt{3}$ and $7-4\sqrt{3}$.

The product of these two numbers is found as follows:

$$\begin{aligned} & (7+4\sqrt{3})(7-4\sqrt{3}) \\ &= 7(7-4\sqrt{3}) + 4\sqrt{3}(7-4\sqrt{3}) \\ &= 49 - 28\sqrt{3} + 28\sqrt{3} - (4\sqrt{3} \times 4\sqrt{3}) \\ &= 49 - (4 \times 4 \times 3) \\ &= 49 - 48 \\ &= 1 \text{ (a rational number)} \end{aligned}$$

2. Let us consider the two irrational numbers $2+\sqrt{3}$ and $5+\sqrt{6}$.

The product of these two numbers is found as follows:

$$\begin{aligned} & (2+\sqrt{3})(5+\sqrt{6}) \\ &= 2(5+\sqrt{6}) + \sqrt{3}(5+\sqrt{6}) \\ &= 10 + 2\sqrt{6} + 5\sqrt{3} + \sqrt{18} \\ &= 10 + 2\sqrt{6} + 5\sqrt{3} + 3\sqrt{2} \text{ (an irrational number)} \end{aligned}$$

Example 2: State two irrational numbers whose:

1. quotient is a rational number
2. quotient is an irrational number

Solution:

1. Let us consider the two irrational numbers $4+3\sqrt{11}$ and $12+9\sqrt{11}$.

The quotient of these two numbers is found as follows:

$$\begin{aligned} & \frac{4+3\sqrt{11}}{12+9\sqrt{11}} \\ &= \frac{4+3\sqrt{11}}{3(4+3\sqrt{11})} \\ &= \frac{1}{3} \text{ (a rational number)} \end{aligned}$$

2. Let us consider the two irrational numbers $3\sqrt{33}$ and $6\sqrt{11}$.

The quotient of these two numbers is found as follows:

$$\begin{aligned} & \frac{3\sqrt{33}}{6\sqrt{11}} \\ &= \frac{1}{2} \sqrt{\frac{33}{11}} \\ &= \frac{1}{2} \sqrt{3} \text{ (an irrational number)} \end{aligned}$$

Simplifying Expressions Involving Irrational Numbers Using Identities

Consider the expression $(\sqrt{2} + \sqrt{3})(\sqrt{2} - \sqrt{3})$.

To solve this expression, we need to multiply each term in the first bracketed pair with each term in the second bracketed pair. Let us solve this expression.

$$\begin{aligned} (\sqrt{2} + \sqrt{3})(\sqrt{2} - \sqrt{3}) &= \sqrt{2} \times \sqrt{2} - \sqrt{2} \times \sqrt{3} + \sqrt{3} \times \sqrt{2} - \sqrt{3} \times \sqrt{3} \\ &= 2 - \sqrt{6} + \sqrt{6} - 3 \\ &= 2 - 3 \\ &= -1 \end{aligned}$$

Is there an easier way to simplify the given expression? Yes, there is. We can solve such expressions using the **identity**: $(\sqrt{x} + \sqrt{y})(\sqrt{x} - \sqrt{y}) = x - y$.

On using this identity, we get the same value as obtained via the longer method.

$$(\sqrt{2} + \sqrt{3})(\sqrt{2} - \sqrt{3}) = 2 - 3 = -1$$

There are other identities related to the square roots of **positive real numbers**. Let us first learn some of these identities and then we will apply them to solve problems.

Identities Related to the Square Roots of Positive Real Numbers

Here are some identities that help us solve problems involving irrational numbers.

If we consider x, y, p and q to be positive real numbers, then

- $\sqrt{xy} = \sqrt{x} \times \sqrt{y}$
- $\sqrt{\frac{x}{y}} = \frac{\sqrt{x}}{\sqrt{y}}$
- $(\sqrt{x} + \sqrt{y})(\sqrt{x} - \sqrt{y}) = x - y$
- $(\sqrt{x} + y)(\sqrt{x} - y) = x - y^2$
- $(x - \sqrt{y})(x + \sqrt{y}) = x^2 - y$
- $(\sqrt{x} + \sqrt{y})^2 = x + y + 2\sqrt{xy}$
- $(\sqrt{x} - \sqrt{y})^2 = x + y - 2\sqrt{xy}$
- $(\sqrt{p} + \sqrt{q})(\sqrt{x} + \sqrt{y}) = \sqrt{px} + \sqrt{py} + \sqrt{qx} + \sqrt{qy}$

Concept Builder

Important algebraic identities

Here are some important algebraic identities that help us solve various types of problems.

- $(a + b)^2 = a^2 + 2ab + b^2$
- $(a - b)^2 = a^2 - 2ab + b^2$
- $(a + b)(a - b) = a^2 - b^2$

- $(x + a)(x + b) = x^2 + (a + b)x + ab$
- $(a + b)^3 = a^3 + b^3 + 3ab(a + b)$
- $(a - b)^3 = a^3 - b^3 - 3ab(a - b)$

Solved Examples

Easy

Example 1: Simplify the following expressions using identities.

1. $(\sqrt{13} + \sqrt{10})(\sqrt{13} - \sqrt{10})$

2. $(3 - \sqrt{5})(3 + \sqrt{5})$

3. $(\sqrt{31} + 5)(\sqrt{31} - 5)$

4. $(\sqrt{11} - \sqrt{17})^2$

5. $(\sqrt{6} + \sqrt{11})^2$

6. $(\sqrt{2} + \sqrt{3})(\sqrt{7} + \sqrt{23})$

Solution:

1. $(\sqrt{13} + \sqrt{10})(\sqrt{13} - \sqrt{10})$

This expression is of the form $(\sqrt{x} + \sqrt{y})(\sqrt{x} - \sqrt{y})$, where $x = 13$ and $y = 10$.

On using the identity $(\sqrt{x} + \sqrt{y})(\sqrt{x} - \sqrt{y}) = x - y$, we obtain:

$$(\sqrt{13} + \sqrt{10})(\sqrt{13} - \sqrt{10}) = 13 - 10 = 3$$

2. $(3 - \sqrt{5})(3 + \sqrt{5})$

This expression is of the form $(x - \sqrt{y})(x + \sqrt{y})$, where $x = 3$ and $y = 5$.

On using the identity $(x - \sqrt{y})(x + \sqrt{y}) = x^2 - y$, we obtain:

$$(3 - \sqrt{5})(3 + \sqrt{5}) = 3^2 - 5 = 9 - 5 = 4$$

3. $(\sqrt{31} + 5)(\sqrt{31} - 5)$

This expression is of the form $(\sqrt{x} + y)(\sqrt{x} - y)$, where $x = 31$ and $y = 5$.

On using the identity $(\sqrt{x} + y)(\sqrt{x} - y) = x - y^2$, we obtain:

$$(\sqrt{31} + 5)(\sqrt{31} - 5) = 31 - 5^2 = 31 - 25 = 6$$

4. $(\sqrt{11} - \sqrt{17})^2$

This expression is of the form $(\sqrt{x} - \sqrt{y})^2$, where $x = 11$ and $y = 17$.

On using the identity $(\sqrt{x} - \sqrt{y})^2 = x + y - 2\sqrt{xy}$, we obtain:

$$(\sqrt{11} - \sqrt{17})^2 = 11 + 17 - 2\sqrt{11 \times 17} = 28 - 2\sqrt{187}$$

5. $(\sqrt{6} + \sqrt{11})^2$

This expression is of the form $(\sqrt{x} + \sqrt{y})^2$, where $x = 6$ and $y = 11$.

On using the identity $(\sqrt{x} + \sqrt{y})^2 = x + y + 2\sqrt{xy}$, we obtain:

$$(\sqrt{6} + \sqrt{11})^2 = 6 + 11 + 2\sqrt{11 \times 6} = 17 + 2\sqrt{66}$$

6. $(\sqrt{2} + \sqrt{3})(\sqrt{7} + \sqrt{23})$

This expression is of the form $(\sqrt{p} + \sqrt{q})(\sqrt{x} + \sqrt{y})$, where $p = 2$, $q = 3$, $x = 7$ and $y = 23$.

On using the identity $(\sqrt{p} + \sqrt{q})(\sqrt{x} + \sqrt{y}) = \sqrt{px} + \sqrt{py} + \sqrt{qx} + \sqrt{qy}$, we obtain:

$$(\sqrt{2} + \sqrt{3})(\sqrt{7} + \sqrt{23}) = \sqrt{2 \times 7} + \sqrt{2 \times 23} + \sqrt{3 \times 7} + \sqrt{3 \times 23} = \sqrt{14} + \sqrt{46} + \sqrt{21} + \sqrt{69}$$

Example 1: The diagonals of a rhombus measure $(4\sqrt{5} + 8)$ units and $(4\sqrt{5} - 8)$ units. Find the area of the rhombus.

Solution:

$$\text{Area of a rhombus} = \frac{1}{2} \times \text{Diagonal 1} \times \text{Diagonal 2}$$

$$\Rightarrow \text{Area of given rhombus} = \frac{1}{2} (4\sqrt{5} + 8) (4\sqrt{5} - 8)$$

$$\Rightarrow \text{Area of given rhombus} = \frac{1}{2} \times 16 (\sqrt{5} + 2) (\sqrt{5} - 2)$$

$$\Rightarrow \text{Area of given rhombus} = 8 (\sqrt{5} + 2) (\sqrt{5} - 2)$$

On using the identity, $(\sqrt{x} + y)(\sqrt{x} - y) = x - y^2$ (where, $x = 5$; $y = 2$) we get

$$\text{Area of given rhombus} = 8 (5 - 2^2) \text{ square units}$$

$$\Rightarrow \text{Area of given rhombus} = 8 (5 - 4) \text{ square units}$$

$$\Rightarrow \text{Area of given rhombus} = 8 \text{ square units}$$

Example 2: Simplify the irrational number $\sqrt{6 - \sqrt{20}}$.

Solution:

$$\sqrt{6 - \sqrt{20}}$$

$$= \sqrt{5 + 1 - 2\sqrt{5}}$$

$$= \sqrt{(\sqrt{5})^2 + 1^2 - 2 \times \sqrt{5} \times 1}$$

$$= \sqrt{(\sqrt{5} - 1)^2}$$

$$= \sqrt{5} - 1$$

$$\left[\text{By the identity: } (a - b)^2 = a^2 + b^2 - 2ab \right]$$

Example 1: Simplify the expression $\sqrt{(2 + \sqrt{45}) - (3\sqrt{5} - 6) + \sqrt{60}}$.

Solution:

$$\begin{aligned} & \sqrt{(2 + \sqrt{45}) - (3\sqrt{5} - 6) + \sqrt{60}} \\ &= \sqrt{2 + \sqrt{3 \times 3 \times 5} - 3\sqrt{5} + 6 + \sqrt{2 \times 2 \times 3 \times 5}} \\ &= \sqrt{2 + 3\sqrt{5} - 3\sqrt{5} + 6 + 2\sqrt{3 \times 5}} \\ &= \sqrt{2 + 6 + 2\sqrt{3 \times 5}} \\ &= \sqrt{8 + 2\sqrt{3} \times \sqrt{5}} \\ &= \sqrt{3 + 5 + 2\sqrt{3} \times \sqrt{5}} \\ &= \sqrt{(\sqrt{3})^2 + (\sqrt{5})^2 + 2 \times \sqrt{3} \times \sqrt{5}} \\ &= \sqrt{(\sqrt{3} + \sqrt{5})^2} \quad \left[\text{By the identity: } (a + b)^2 = a^2 + b^2 + 2ab \right] \\ &= \sqrt{3} + \sqrt{5} \end{aligned}$$

Example 2: The length and breadth of a rectangle are $(2 + \sqrt{6})$ units and $(4 + \sqrt{3})$ units. Find the perimeter and area of the rectangle.

Solution:

Length (l) of the rectangle = $(2 + \sqrt{6})$ units

Breadth (b) of the rectangle = $(4 + \sqrt{3})$ units

Perimeter of the rectangle is given as:

$$\begin{aligned} & 2(l + b) \text{ units} \\ &= 2[(2 + \sqrt{6}) + (4 + \sqrt{3})] \text{ units} \\ &= 2[6 + \sqrt{3} + \sqrt{6}] \text{ units} \\ &= (12 + 2\sqrt{3} + 2\sqrt{6}) \text{ units} \end{aligned}$$

Area of the rectangle is given as:

$$\begin{aligned}
 & (1 \times b) \text{ sq. units} \\
 &= (2 + \sqrt{6})(4 + \sqrt{3}) \text{ sq. units} \\
 &= (2 \times 4 + 2 \times \sqrt{3} + \sqrt{6} \times 4 + \sqrt{6} \times \sqrt{3}) \text{ sq. units} \\
 &= (8 + 2\sqrt{3} + 4\sqrt{6} + \sqrt{18}) \text{ sq. units} \\
 &= (8 + 2\sqrt{3} + 4\sqrt{6} + 3\sqrt{2}) \text{ sq. units}
 \end{aligned}$$

Rationalizing the Denominators

Rationalization

So far we have studied different operations on rational and irrational numbers, such as addition and subtraction. Now, what if we have to add two irrational fractions whose denominators are irrational numbers, say of the form $\sqrt{2} + 3$ and $\sqrt{3} + 5$? In such cases, we have to first rationalize the denominators, i.e., we have to make the denominators rational quantities (even if the numerators remain in the irrational form).

What does rationalization mean? Rationalization is the process in which an irrational fraction having a **surd** in the denominator is rewritten to obtain a rational number in the denominator. The surd may be a monomial or a binomial having a square root.

In this lesson, we will learn to:

- Rationalize the denominator of an irrational fraction
- Solve expressions using the rationalization method

Rationalizing the Denominator

Let us understand the method of rationalizing the denominator of an irrational fraction by

taking the example of $\frac{1}{\sqrt{7}}$.

We know that $\sqrt{x} \times \sqrt{x} = x$.

$\therefore \sqrt{7} \times \sqrt{7} = 7$, which is a rational number

So, we can write the expression $\frac{1}{\sqrt{7}}$ as:

$$\begin{aligned}
 & \frac{1}{\sqrt{7}} \times 1 \\
 &= \frac{1}{\sqrt{7}} \times \frac{\sqrt{7}}{\sqrt{7}} \quad \left(\because \frac{\sqrt{7}}{\sqrt{7}} = 1 \right) \\
 &= \frac{\sqrt{7}}{7}
 \end{aligned}$$

Thus, we get a rational number as the denominator.

Let us now understand what is meant by 'conjugate of a number'.

When a number is represented as the sum or difference of a rational number and an irrational number or two irrational numbers, the conjugate of that number just differs by the sign in between. For example, the conjugate of $\sqrt{2} + 3$ is $\sqrt{2} - 3$ and that of $\sqrt{2} + \sqrt{3}$ is $\sqrt{2} - \sqrt{3}$ or $\sqrt{3} - \sqrt{2}$. While solving irrational fractions having such denominators, we multiply and divide the fractions by the conjugates of the denominators.

Solving an Expression Using the Rationalization Method

Solved Examples

Example 1: Rationalize the denominator of $\frac{2}{\sqrt{3}}$.

Solution: We know that $\sqrt{3} \times \sqrt{3} = 3$, which is a rational number.

So, we multiply and divide $\frac{2}{\sqrt{3}}$ by $\sqrt{3}$.

$$\frac{2}{\sqrt{3}} \times \frac{\sqrt{3}}{\sqrt{3}} = \frac{2\sqrt{3}}{3}$$

Thus, we obtain a rational number as the denominator.

Example 2: Rationalize the denominator of $\frac{11\sqrt{3}-6}{\sqrt{3}+2}$.

Solution: To rationalize the denominator, we multiply and divide $\frac{11\sqrt{3}-6}{\sqrt{3}+2}$ by the conjugate of $\sqrt{3}+2$.

The conjugate of $\sqrt{3}+2$ is $\sqrt{3}-2$.

$$\begin{aligned}\therefore \frac{11\sqrt{3}-6}{\sqrt{3}+2} &= \frac{11\sqrt{3}-6}{(\sqrt{3}+2)} \times \frac{(\sqrt{3}-2)}{(\sqrt{3}-2)} \\ &= \frac{(11\sqrt{3} \times \sqrt{3}) - (11\sqrt{3} \times 2) - (6 \times \sqrt{3}) + (6 \times 2)}{(\sqrt{3})^2 - (2)^2} \\ &= \frac{33 - 22\sqrt{3} - 6\sqrt{3} + 12}{3 - 4} \\ &= 28\sqrt{3} - 45\end{aligned}$$

Example 1: Simplify the expression $\frac{6}{2\sqrt{3}-\sqrt{6}} + \frac{\sqrt{6}}{\sqrt{3}+\sqrt{2}} - \frac{4\sqrt{3}}{\sqrt{6}-\sqrt{2}}$.

Solution:

$$\text{Let } \frac{6}{2\sqrt{3}-\sqrt{6}} + \frac{\sqrt{6}}{\sqrt{3}+\sqrt{2}} - \frac{4\sqrt{3}}{\sqrt{6}-\sqrt{2}} = a + b - c$$

Where,

$$a = \frac{6}{2\sqrt{3}-\sqrt{6}}, \quad b = \frac{\sqrt{6}}{\sqrt{3}+\sqrt{2}} \quad \text{and} \quad c = \frac{4\sqrt{3}}{\sqrt{6}-\sqrt{2}}$$

To solve such an expression, we have to first rationalize the denominator of each term.

$$\begin{aligned}a &= \frac{6}{2\sqrt{3}-\sqrt{6}} \\ \Rightarrow a &= \frac{6}{2\sqrt{3}-\sqrt{6}} \times \frac{2\sqrt{3}+\sqrt{6}}{2\sqrt{3}+\sqrt{6}} = \frac{6(2\sqrt{3}+\sqrt{6})}{(2\sqrt{3})^2 - (\sqrt{6})^2} = \frac{6(2\sqrt{3}+\sqrt{6})}{12-6} = 2\sqrt{3}+\sqrt{6}\end{aligned}$$

$$b = \frac{\sqrt{6}}{\sqrt{3} + \sqrt{2}}$$

$$\Rightarrow b = \frac{\sqrt{6}}{\sqrt{3} + \sqrt{2}} \times \frac{\sqrt{3} - \sqrt{2}}{\sqrt{3} - \sqrt{2}} = \frac{\sqrt{6}(\sqrt{3} - \sqrt{2})}{(\sqrt{3})^2 - (\sqrt{2})^2} = \frac{3\sqrt{2} - 2\sqrt{3}}{3 - 2} = 3\sqrt{2} - 2\sqrt{3}$$

$$c = \frac{4\sqrt{3}}{\sqrt{6} - \sqrt{2}}$$

$$\Rightarrow c = \frac{4\sqrt{3}}{\sqrt{6} - \sqrt{2}} \times \frac{\sqrt{6} + \sqrt{2}}{\sqrt{6} + \sqrt{2}} = \frac{4\sqrt{3}(\sqrt{6} + \sqrt{2})}{(\sqrt{6})^2 - (\sqrt{2})^2} = \frac{4(3\sqrt{2} + \sqrt{6})}{6 - 2} = 3\sqrt{2} + \sqrt{6}$$

On substituting the values of a , b and c , we obtain:

$$\begin{aligned} \frac{6}{2\sqrt{3} - \sqrt{6}} + \frac{\sqrt{6}}{\sqrt{3} + \sqrt{2}} - \frac{4\sqrt{3}}{\sqrt{6} - \sqrt{2}} &= (2\sqrt{3} + \sqrt{6}) + (3\sqrt{2} - 2\sqrt{3}) - (3\sqrt{2} + \sqrt{6}) \\ &= 2\sqrt{3} + \sqrt{6} + 3\sqrt{2} - 2\sqrt{3} - 3\sqrt{2} - \sqrt{6} \\ &= 0 \end{aligned}$$

Example 2: Evaluate $\frac{15}{\sqrt{10} + \sqrt{20} + \sqrt{40} - \sqrt{5} - \sqrt{80}}$.

Solution:

Let us first simplify the denominator of the given expression.

$$\begin{aligned} &\sqrt{10} + \sqrt{20} + \sqrt{40} - \sqrt{5} - \sqrt{80} \\ &= \sqrt{10} + \sqrt{2^2 \times 5} + \sqrt{2^2 \times 10} - \sqrt{5} - \sqrt{4^2 \times 5} \\ &= \sqrt{10} + 2\sqrt{5} + 2\sqrt{10} - \sqrt{5} - 4\sqrt{5} \\ &= 3\sqrt{10} - 3\sqrt{5} \\ &= 3(\sqrt{10} - \sqrt{5}) \end{aligned}$$

$$\begin{aligned}
 \therefore \frac{15}{\sqrt{10} + \sqrt{20} + \sqrt{40} - \sqrt{5} - \sqrt{80}} &= \frac{15}{3(\sqrt{10} - \sqrt{5})} \\
 &= \frac{5}{\sqrt{10} - \sqrt{5}} \\
 &= \frac{5}{\sqrt{10} - \sqrt{5}} \times \frac{\sqrt{10} + \sqrt{5}}{\sqrt{10} + \sqrt{5}} \\
 &= \frac{5(\sqrt{10} + \sqrt{5})}{(\sqrt{10})^2 - (\sqrt{5})^2} \\
 &= \frac{5(\sqrt{10} + \sqrt{5})}{10 - 5} \\
 &= \sqrt{10} + \sqrt{5}
 \end{aligned}$$

Example 1: Prove that $\frac{1}{3-\sqrt{8}} - \frac{1}{\sqrt{8}-\sqrt{7}} + \frac{1}{\sqrt{7}-\sqrt{6}} - \frac{1}{\sqrt{6}-\sqrt{5}} + \frac{1}{\sqrt{5}-2} = 5$.

Solution:

$$\begin{aligned}
 \text{LHS} &= \frac{1}{3-\sqrt{8}} - \frac{1}{\sqrt{8}-\sqrt{7}} + \frac{1}{\sqrt{7}-\sqrt{6}} - \frac{1}{\sqrt{6}-\sqrt{5}} + \frac{1}{\sqrt{5}-2} \\
 &= \left(\frac{1}{3-\sqrt{8}} \times \frac{3+\sqrt{8}}{3+\sqrt{8}} \right) - \left(\frac{1}{\sqrt{8}-\sqrt{7}} \times \frac{\sqrt{8}+\sqrt{7}}{\sqrt{8}+\sqrt{7}} \right) + \left(\frac{1}{\sqrt{7}-\sqrt{6}} \times \frac{\sqrt{7}+\sqrt{6}}{\sqrt{7}+\sqrt{6}} \right) \\
 &\quad - \left(\frac{1}{\sqrt{6}-\sqrt{5}} \times \frac{\sqrt{6}+\sqrt{5}}{\sqrt{6}+\sqrt{5}} \right) + \left(\frac{1}{\sqrt{5}-2} \times \frac{\sqrt{5}+2}{\sqrt{5}+2} \right) \\
 &= \frac{3+\sqrt{8}}{(3)^2 - (\sqrt{8})^2} - \frac{\sqrt{8}+\sqrt{7}}{(\sqrt{8})^2 - (\sqrt{7})^2} + \frac{\sqrt{7}+\sqrt{6}}{(\sqrt{7})^2 - (\sqrt{6})^2} \\
 &\quad - \frac{\sqrt{6}+\sqrt{5}}{(\sqrt{6})^2 - (\sqrt{5})^2} + \frac{\sqrt{5}+2}{(\sqrt{5})^2 - (2)^2} \\
 &= \frac{3+\sqrt{8}}{9-8} - \frac{\sqrt{8}+\sqrt{7}}{8-7} + \frac{\sqrt{7}+\sqrt{6}}{7-6} - \frac{\sqrt{6}+\sqrt{5}}{6-5} + \frac{\sqrt{5}+2}{5-4}
 \end{aligned}$$

$$\begin{aligned}
&= (3 + \sqrt{8}) - (\sqrt{8} + \sqrt{7}) + (\sqrt{7} + \sqrt{6}) - (\sqrt{6} + \sqrt{5}) + (\sqrt{5} + 2) \\
&= 3 + \sqrt{8} - \sqrt{8} - \sqrt{7} + \sqrt{7} + \sqrt{6} - \sqrt{6} - \sqrt{5} + \sqrt{5} + 2 \\
&= 3 + 2 \\
&= 5 \\
&= \text{RHS}
\end{aligned}$$

Example 2: If $x = \frac{\sqrt{3}+1}{2}$, then find the value of $4x^3 + 2x^2 - 8x + 7$.

Solution:

$$\begin{aligned}
x &= \frac{\sqrt{3}+1}{2} \\
\Rightarrow 2x &= \sqrt{3}+1 \\
2x-1 &= \sqrt{3} \\
\Rightarrow (2x-1)^2 &= (\sqrt{3})^2 \\
\Rightarrow 4x^2 - 4x + 1 &= 3 \\
\Rightarrow 4x^2 - 4x - 2 &= 0 \\
\Rightarrow 2x^2 - 2x - 1 &= 0 \quad \dots(1)
\end{aligned}$$

$$\begin{aligned}
\therefore 4x^3 + 2x^2 - 8x + 7 &= 2x(2x^2 - 2x - 1) + 3(2x^2 - 2x - 1) + 10 \\
&= 2x(0) + 3(0) + 10 \quad (\text{From equation 1}) \\
&= 0 + 0 + 10 \\
&= 10
\end{aligned}$$

Laws of Rational Exponents of Real Numbers

Exponents or Indices

The term 'exponent' refers to the number of times a quantity is multiplied with itself. It is also called 'index' or 'power'. The exponential form of any number is x^y , where **x is the base and y is the exponent**. It is read as 'yth power of x' or 'x raised to the power y'.

Till now, we have studied the laws of exponents for numbers having non-zero integers as the base and the **integral exponent**. For example, consider the expression $11^5 \div 11^3$. It can be simplified using the law of exponents $a^p \div a^q = a^{(p-q)}$ as follows: $11^5 \div 11^3 = 11^{(5-3)} = 11^2$

But what about numbers whose base is any real number or whose exponent is any rational number?

Can we simplify them using the same laws? Go through this lesson to find out how to simplify such expressions.

Numbers with Fractional Indices

While studying about the exponents, we come across few numbers having fractional indices. For example, $8^{\frac{1}{3}}$, $16^{\frac{1}{2}}$, $81^{\frac{1}{4}}$ etc.

Do you know what these numbers mean?

We know that the square root of a number is represented with the index $\frac{1}{2}$ as well as with the symbol $\sqrt{}$.

For example, the square root of 16 can be represented as $16^{\frac{1}{2}}$ and $\sqrt{16}$.

Similarly, the square root of any real number a can be represented as $a^{\frac{1}{2}}$ and \sqrt{a} .

So, there are two ways to represent the square roots of numbers.

In the same manner, we can represent the **cube root, fourth root, fifth root, ..., n^{th} root** of a real number with the symbol $\sqrt[n]{}$ as well as with fractional indices.

For example, the cube root of 8 can be represented as $\sqrt[3]{8}$ and $8^{\frac{1}{3}}$. The fourth root of 81 can be represented as $\sqrt[4]{81}$ and $81^{\frac{1}{4}}$.

Similarly, the n^{th} root of a number can also be represented as $\sqrt[n]{a}$ and $a^{\frac{1}{n}}$.

Solved Examples

Easy

Example 1: Find the values of the following index numbers.

i. $27^{\frac{1}{3}}$

ii. $256^{\frac{1}{4}}$

iii. $576^{\frac{1}{2}}$

iv. $4096^{\frac{1}{6}}$

Solution:

i. $27^{\frac{1}{3}}$ means the cube root of 27.

Therefore,

$$\begin{aligned} 27^{\frac{1}{3}} &= \sqrt[3]{27} \\ \Rightarrow 27^{\frac{1}{3}} &= \sqrt[3]{3 \times 3 \times 3} \\ \Rightarrow 27^{\frac{1}{3}} &= 3 \end{aligned}$$

ii. $256^{\frac{1}{4}}$ means fourth root of 256.

Therefore,

$$\begin{aligned} 256^{\frac{1}{4}} &= \sqrt[4]{256} \\ \Rightarrow 256^{\frac{1}{4}} &= \sqrt[4]{4 \times 4 \times 4 \times 4} \\ \Rightarrow 256^{\frac{1}{4}} &= 4 \end{aligned}$$

iii. $576^{\frac{1}{2}}$ means square root of 576.

Therefore,

$$\begin{aligned} 576^{\frac{1}{2}} &= \sqrt{576} \\ \Rightarrow 576^{\frac{1}{2}} &= \sqrt{24 \times 24} \\ \Rightarrow 576^{\frac{1}{2}} &= 24 \end{aligned}$$

iv. $4096^{\frac{1}{6}}$ means sixth root of 4096.

Therefore,

$$4096^{\frac{1}{6}} = \sqrt[6]{4096}$$

$$\Rightarrow 4096^{\frac{1}{6}} = \sqrt[6]{4 \times 4 \times 4 \times 4 \times 4 \times 4}$$

$$\Rightarrow 4096^{\frac{1}{6}} = 4$$

Laws of Exponents for Real Numbers

Consider two real numbers a and b and two rational numbers m and n . The laws of exponents involving these real bases and rational exponents can be written as follows:

- $a^m \times a^n = a^{m+n}$
- $a^m \div a^n = \frac{a^m}{a^n} = a^{m-n} \quad (a \neq 0)$
- $(a^m)^n = a^{mn} = (a^n)^m$
- $\frac{a^m}{b^m} = \left(\frac{a}{b}\right)^m$
- $a^m \times b^m = (ab)^m$
- $a^{-m} = \frac{1}{a^m}$
- $a^0 = 1$

Solved Examples

Example 1: Simplify the following expressions.

1. $\left(\frac{27}{125}\right)^{\frac{2}{3}}$
2. $\sqrt[3]{(512)^{-2}}$

Solution:

$$\begin{aligned}
 \text{i) } & \left(\frac{27}{125}\right)^{\frac{2}{3}} \\
 &= \left[\frac{(3)^3}{(5)^3}\right]^{\frac{2}{3}} \\
 &= \left[\left(\frac{3}{5}\right)^3\right]^{\frac{2}{3}} \quad \left[\because \frac{a^m}{b^m} = \left(\frac{a}{b}\right)^m\right] \\
 &= \left(\frac{3}{5}\right)^{3 \times \frac{2}{3}} \quad \left[\because (a^m)^n = a^{m \times n}\right] \\
 &= \left(\frac{3}{5}\right)^2 \\
 &= \frac{(3)^2}{(5)^2} \quad \left[\because \left(\frac{a}{b}\right)^m = \frac{a^m}{b^m}\right] \\
 &= \frac{9}{25}
 \end{aligned}$$

$$\begin{aligned}
 \text{ii) } & \sqrt[3]{(512)^{-2}} \\
 &= \left[(512)^{-2}\right]^{\frac{1}{3}} \\
 &= (512)^{\frac{-2}{3}} \quad \left[\because (a^m)^n = a^{mn}\right] \\
 &= (8^3)^{\frac{-2}{3}} \\
 &= (8)^{3 \times \frac{-2}{3}} \quad \left[\because (a^m)^n = a^{mn}\right] \\
 &= (8)^{-2} \\
 &= \frac{1}{8^2} \quad \left[\because a^{-m} = \frac{1}{a^m}\right] \\
 &= \frac{1}{64}
 \end{aligned}$$

Example 2: Simplify the expression $\left(\sqrt{\frac{2}{3}}\right)^{\frac{3}{5}} \times \left(\sqrt{\frac{1}{7}}\right)^{\frac{3}{5}}$.

Solution:

$$\begin{aligned}
 & \left(\sqrt{\frac{2}{3}}\right)^{\frac{3}{5}} \times \left(\sqrt{\frac{1}{7}}\right)^{\frac{3}{5}} \\
 &= \left(\sqrt{\frac{2}{3}} \times \sqrt{\frac{1}{7}}\right)^{\frac{3}{5}} \quad [\because a^m \times b^m = (ab)^m] \\
 &= \left(\sqrt{\frac{2}{21}}\right)^{\frac{3}{5}}
 \end{aligned}$$

Example 3: Simplify $\sqrt[3]{2^4} \times \sqrt[3]{3^4}$.

Solution:

$$\begin{aligned}
 & \sqrt[3]{2^4} \times \sqrt[3]{3^4} \\
 &= (2^4)^{\frac{1}{3}} \times (3^4)^{\frac{1}{3}} \\
 &= (2)^{\frac{4}{3}} \times (3)^{\frac{4}{3}} \quad \left[\because (a^m)^n = a^{mn}\right] \\
 &= (2 \times 3)^{\frac{4}{3}} \quad \left[\because a^m \times b^m = (ab)^m\right] \\
 &= 6^{\frac{4}{3}}
 \end{aligned}$$

Example 1: Find the values of x and y in the expression $3^{x+1} \times 7^{2y-1} = 189$.

Solution: It is given that $3^{x+1} \times 7^{2y-1} = 189$

$$\begin{aligned}
 \Rightarrow 3^{x+1} \times 7^{2y-1} &= 3 \times 3 \times 3 \times 7 \\
 \Rightarrow 3^{x+1} \times 7^{2y-1} &= 3^3 \times 7^1
 \end{aligned}$$

On equating the exponents of 3 and 7 on both sides of the above equation, we get:

$$x + 1 = 3 \text{ and } 2y - 1 = 1$$

$$\Rightarrow x = 3 - 1 = 2 \text{ and } 2y = 1 + 1 = 2$$

$$\Rightarrow x = 2 \text{ and } y = 1$$

Thus, the values of x and y are 2 and 1 respectively.

$$\left(\frac{243}{32}\right)^{-\frac{4}{5}} \times \left[\left(\frac{625}{81}\right)^{-\frac{3}{4}} \div \left(\frac{25}{4}\right)^{-\frac{3}{2}} \right]$$

Example 1: Simplify the expression

Solution:

$$\begin{aligned} & \left(\frac{243}{32}\right)^{-\frac{4}{5}} \times \left[\left(\frac{625}{81}\right)^{-\frac{3}{4}} \div \left(\frac{25}{4}\right)^{-\frac{3}{2}} \right] \\ &= \left[\left(\frac{3}{2}\right)^5 \right]^{-\frac{4}{5}} \times \left[\left\{ \left(\frac{5}{3}\right)^4 \right\}^{-\frac{3}{4}} \div \left\{ \left(\frac{5}{2}\right)^2 \right\}^{-\frac{3}{2}} \right] \\ &= \left(\frac{3}{2}\right)^{5 \times -\frac{4}{5}} \times \left\{ \left(\frac{5}{3}\right)^{4 \times -\frac{3}{4}} \div \left(\frac{5}{2}\right)^{2 \times -\frac{3}{2}} \right\} \quad \left[\because (a^m)^n = a^{mn} \right] \\ &= \left(\frac{3}{2}\right)^{-4} \times \left\{ \left(\frac{5}{3}\right)^{-3} \div \left(\frac{5}{2}\right)^{-3} \right\} \\ &= \left(\frac{2}{3}\right)^4 \times \left\{ \left(\frac{3}{5}\right)^3 \div \left(\frac{2}{5}\right)^3 \right\} \quad \left[\because \left(\frac{a}{b}\right)^{-m} = \left(\frac{b}{a}\right)^m \right] \\ &= \left(\frac{2}{3}\right)^4 \times \left(\frac{3}{5} \div \frac{2}{5}\right)^3 \quad \left[\because a^m \div b^m = (a \div b)^m \right] \\ &= \left(\frac{2}{3}\right)^4 \times \left(\frac{3}{5} \times \frac{5}{2}\right)^3 \\ &= \left(\frac{2}{3}\right)^4 \times \left(\frac{3}{2}\right)^3 \\ &= \left(\frac{2}{3}\right)^4 \times \left(\frac{2}{3}\right)^{-3} \quad \left[\because \left(\frac{b}{a}\right)^m = \left(\frac{a}{b}\right)^{-m} \right] \\ &= \left(\frac{2}{3}\right)^{4-3} \quad \left[\because a^m \times a^n = a^{m+n} \right] \\ &= \frac{2}{3} \end{aligned}$$

Example 2: Prove that $\frac{x^{a(b-c)}}{x^{b(a-c)}} \div \left(\frac{x^b}{x^a}\right)^c = 1$.

Solution:

$$\begin{aligned} & \frac{x^{a(b-c)}}{x^{b(a-c)}} \div \left(\frac{x^b}{x^a} \right)^c \\ &= \frac{x^{(ab-ac)}}{x^{(ab-bc)}} \div (x^{b-a})^c \quad \left[\because (a^m)^n = a^{mn} \text{ and } \frac{a^m}{a^n} = a^{m-n} \right] \\ &= x^{(ab-ac)-(ab-bc)} \times \frac{1}{x^{(b-a)c}} \quad \left[\because \frac{a^m}{a^n} = a^{m-n} \right] \\ &= x^{ab-ac-ab+bc} \times \frac{1}{x^{bc-ac}} \quad \left[\because (a^m)^n = a^{mn} \right] \\ &= x^{-ac+bc} \times x^{ac-bc} \quad \left[\because \frac{1}{a^m} = a^{-m} \right] \\ &= x^{-ac+bc+ac-bc} \quad \left[\because a^m \times a^n = a^{m+n} \right] \\ &= x^0 \\ &= 1 \end{aligned}$$

