

Relations and Functions

Short Answer Type Questions

Q. 1 Let $A = \{a, b, c\}$ and the relation R be defined on A as follows

$$R = \{(a, a), (b, c), (a, b)\}$$

Then, write minimum number of ordered pairs to be added in R to make R reflexive and transitive.

Sol. Given relation, $R = \{(a, a), (b, c), (a, b)\}$.

To make R is reflexive we must add (b, b) and (c, c) to R . Also, to make R is transitive we must add (a, c) to R .

So, minimum number of ordered pair is to be added are (b, b) , (c, c) , (a, c) .

Q. 2 Let D be the domain of the real valued function f defined by $f(x) = \sqrt{25 - x^2}$. Then, write D .

Sol. Given function is, $f(x) = \sqrt{25 - x^2}$

For real valued of $f(x)$ $25 - x^2 \geq 0$

$$x^2 \leq 25$$

$$-5 \leq x \leq +5$$

$$\therefore D = [-5, 5]$$

Q. 3 If $f, g : R \rightarrow R$ be defined by $f(x) = 2x + 1$ and $g(x) = x^2 - 2$, $\forall x \in R$, respectively. Then, find gof .

Thinking Process

If $f, g : R \rightarrow R$ be two functions, then $gof(x) = g\{f(x)\} \forall x \in R$.

Sol. Given that, $f(x) = 2x + 1$ and $g(x) = x^2 - 2$, $\forall x \in R$

$$\begin{aligned} \therefore gof &= g\{f(x)\} \\ &= g(2x + 1) = (2x + 1)^2 - 2 \\ &= 4x^2 + 4x + 1 - 2 \\ &= 4x^2 + 4x - 1 \end{aligned}$$

Q. 4 Let $f: R \rightarrow R$ be the function defined by $f(x) = 2x - 3, \forall x \in R$. Write f^{-1} .

Sol. Given that,
Now, let

$$\begin{aligned} f(x) &= 2x - 3, \forall x \in R \\ y &= 2x - 3 \\ 2x &= y + 3 \\ x &= \frac{y + 3}{2} \\ \therefore f^{-1}(x) &= \frac{x + 3}{2} \end{aligned}$$

Q. 5 If $A = \{a, b, c, d\}$ and the function $f = \{(a, b), (b, d), (c, a), (d, c)\}$, write f^{-1} .

Sol. Given that,
and

$$\begin{aligned} A &= \{a, b, c, d\} \\ f &= \{(a, b), (b, d), (c, a), (d, c)\} \\ f^{-1} &= \{(b, a), (d, b), (a, c), (c, d)\} \end{aligned}$$

Q. 6 If $f: R \rightarrow R$ is defined by $f(x) = x^2 - 3x + 2$, write $f\{f(x)\}$.

🔑 Thinking Process

To solve this problem use the formula i.e., $(a + b + c)^2 = (a^2 + b^2 + c^2 + 2ab + 2bc + 2ca)$

Sol. Given that,

$$\begin{aligned} \therefore f(x) &= x^2 - 3x + 2 \\ \therefore f\{f(x)\} &= f(x^2 - 3x + 2) \\ &= (x^2 - 3x + 2)^2 - 3(x^2 - 3x + 2) + 2 \\ &= x^4 + 9x^2 + 4 - 6x^3 - 12x + 4x^2 - 3x^2 + 9x - 6 + 2 \\ &= x^4 + 10x^2 - 6x^3 - 3x \\ f\{f(x)\} &= x^4 - 6x^3 + 10x^2 - 3x \end{aligned}$$

Q. 7 Is $g = \{(1, 1), (2, 3), (3, 5), (4, 7)\}$ a function? If g is described by $g(x) = \alpha x + \beta$, then what value should be assigned to α and β ?

Sol. Given that, $g = \{(1, 1), (2, 3), (3, 5), (4, 7)\}$.

Here, each element of domain has unique image. So, g is a function.

Now given that,

$$\begin{aligned} g(x) &= \alpha x + \beta \\ g(1) &= \alpha + \beta \\ \alpha + \beta &= 1 \quad \dots(i) \\ g(2) &= 2\alpha + \beta \\ 2\alpha + \beta &= 3 \quad \dots(ii) \end{aligned}$$

From Eqs. (i) and (ii),

$$\begin{aligned} \Rightarrow 2(1 - \beta) + \beta &= 3 \\ \Rightarrow 2 - 2\beta + \beta &= 3 \\ \Rightarrow 2 - \beta &= 3 \\ \beta &= -1 \\ \text{If } \beta &= -1, \text{ then } \alpha = 2 \\ \alpha = 2, \beta &= -1 \end{aligned}$$



Q. 8 Are the following set of ordered pairs functions? If so examine whether the mapping is injective or surjective.

(i) $\{(x, y) : x \text{ is a person, } y \text{ is the mother of } x\}$.

(ii) $\{(a, b) : a \text{ is a person, } b \text{ is an ancestor of } a\}$.

Sol. (i) Given set of ordered pair is $\{(x, y) : x \text{ is a person, } y \text{ is the mother of } x\}$.

It represent a function. Here, the image of distinct elements of x under f are not distinct, so it is not a injective but it is a surjective.

(ii) Set of ordered pairs = $\{(a, b) : a \text{ is a person, } b \text{ is an ancestor of } a\}$

Here, each element of domain does not have a unique image. So, it does not represent function.

Q. 9 If the mappings f and g are given by $f = \{(1, 2), (3, 5), (4, 1)\}$ and $g = \{(2, 3), (5, 1), (1, 3)\}$, write fog .

Sol. Given that,

$$f = \{(1, 2), (3, 5), (4, 1)\}$$

and

$$g = \{(2, 3), (5, 1), (1, 3)\}$$

Now,

$$fog(2) = f\{g(2)\} = f(3) = 5$$

$$fog(5) = f\{g(5)\} = f(1) = 2$$

$$fog(1) = f\{g(1)\} = f(3) = 5$$

$$fog = \{(2, 5), (5, 2), (1, 5)\}$$

Q. 10 Let C be the set of complex numbers. Prove that the mapping $f : C \rightarrow R$ given by $f(z) = |z|$, $\forall z \in C$, is neither one-one nor onto.

Sol. The mapping

$$f : C \rightarrow R$$

Given,

$$f(z) = |z|, \forall z \in C$$

$$f(1) = |1| = 1$$

$$f(-1) = |-1| = 1$$

$$f(1) = f(-1)$$

But

$$1 \neq -1$$

So, $f(z)$ is not one-one. Also, $f(z)$ is not onto as there is no pre-image for any negative element of R under the mapping $f(z)$.

Q. 11 Let the function $f : R \rightarrow R$ be defined by $f(x) = \cos x$, $\forall x \in R$. Show that f is neither one-one nor onto.

Sol. Given function, $f(x) = \cos x$, $\forall x \in R$

Now,

$$f\left(\frac{\pi}{2}\right) = \cos \frac{\pi}{2} = 0$$

\Rightarrow

$$f\left(\frac{-\pi}{2}\right) = \cos \frac{\pi}{2} = 0$$

\Rightarrow

$$f\left(\frac{\pi}{2}\right) = f\left(\frac{-\pi}{2}\right)$$

But

$$\frac{\pi}{2} \neq \frac{-\pi}{2}$$

So, $f(x)$ is not one-one.

Now, $f(x) = \cos x$, $\forall x \in R$ is not onto as there is no pre-image for any real number. Which does not belonging to the intervals $[-1, 1]$, the range of $\cos x$.

Q. 12 Let $X = \{1, 2, 3\}$ and $Y = \{4, 5\}$. Find whether the following subsets of $X \times Y$ are functions from X to Y or not.

- (i) $f = \{(1, 4), (1, 5), (2, 4), (3, 5)\}$ (ii) $g = \{(1, 4), (2, 4), (3, 4)\}$
 (iii) $h = \{(1, 4), (2, 5), (3, 5)\}$ (iv) $k = \{(1, 4), (2, 5)\}$

Sol. Given that, $X = \{1, 2, 3\}$ and $Y = \{4, 5\}$
 $X \times Y = \{(1, 4), (1, 5), (2, 4), (2, 5), (3, 4), (3, 5)\}$

(i) $f = \{(1, 4), (1, 5), (2, 4), (3, 5)\}$
 f is not a function because f has not unique image.

(ii) $g = \{(1, 4), (2, 4), (3, 4)\}$
 Since, g is a function as each element of the domain has unique image.

(iii) $h = \{(1, 4), (2, 5), (3, 5)\}$
 It is clear that h is a function.

(iv) $k = \{(1, 4), (2, 5)\}$
 k is not a function as 3 has not any image under the mapping.

Q. 13 If functions $f : A \rightarrow B$ and $g : B \rightarrow A$ satisfy $gof = I_A$, then show that f is one-one and g is onto.

Sol. Given that, $f : A \rightarrow B$ and $g : B \rightarrow A$ satisfy $gof = I_A$
 $\therefore gof = I_A$
 $\Rightarrow gof\{f(x_1)\} = gof\{f(x_2)\}$
 $\Rightarrow g(x_1) = g(x_2)$ [$\because gof = I_A$]
 $\therefore x_1 = x_2$
 Hence, f is one-one and g is onto.

Q. 14 Let $f : R \rightarrow R$ be the function defined by $f(x) = \frac{1}{2 - \cos x}$, $\forall x \in R$.

Then, find the range of f .

🔑 **Thinking Process**

Range of $f = \{y \in Y : y = f(x) \text{ for some } x\}$ and use range of $\cos x$ is $[-1, 1]$

Sol. Given function, $f(x) = \frac{1}{2 - \cos x}$, $\forall x \in R$

Let $y = \frac{1}{2 - \cos x}$

$$\Rightarrow 2y - y \cos x = 1$$

$$\Rightarrow y \cos x = 2y - 1$$

$$\Rightarrow \cos x = \frac{2y - 1}{y} = 2 - \frac{1}{y} \Rightarrow \cos x = 2 - \frac{1}{y}$$

$$\Rightarrow -1 \leq \cos x \leq 1 \Rightarrow -1 \leq 2 - \frac{1}{y} \leq 1$$

$$\Rightarrow -3 \leq -\frac{1}{y} \leq -1 \Rightarrow 1 \leq \frac{1}{y} \leq 3$$

$$\Rightarrow \frac{1}{3} \leq \frac{1}{y} \leq 1$$

So, y range is $\left[\frac{1}{3}, 1\right]$.

Q. 15 Let n be a fixed positive integer. Define a relation R in Z as follows $\forall a, b \in Z, aRb$ if and only if $a - b$ is divisible by n . Show that R is an equivalence relation.

Sol. Given that, $\forall a, b \in Z, aRb$ if and only if $a - b$ is divisible by n .

Now,

I. **Reflexive**

$aRa \Rightarrow (a - a)$ is divisible by n , which is true for any integer a as '0' is divisible by n .

Hence, R is reflexive.

II. **Symmetric**

	aRb
\Rightarrow	$a - b$ is divisible by n .
\Rightarrow	$-b + a$ is divisible by n .
\Rightarrow	$-(b - a)$ is divisible by n .
\Rightarrow	$(b - a)$ is divisible by n .
\Rightarrow	bRa

Hence, R is symmetric.

III. **Transitive**

Let aRb and bRc

\Rightarrow	$(a - b)$ is divisible by n and $(b - c)$ is divisible by n
\Rightarrow	$(a - b) + (b - c)$ is divisible by n
\Rightarrow	$(a - c)$ is divisible by n
\Rightarrow	aRc

Hence, R is transitive.

So, R is an equivalence relation.

Long Answer Type Questions

Q. 16 If $A = \{1, 2, 3, 4\}$, define relations on A which have properties of being

- (i) reflexive, transitive but not symmetric.
- (ii) symmetric but neither reflexive nor transitive.
- (iii) reflexive, symmetric and transitive.

Sol. Given that, $A = \{1, 2, 3, 4\}$

(i) Let $R_1 = \{(1, 1), (1, 2), (2, 3), (2, 2), (1, 3), (3, 3)\}$

R_1 is reflexive, since, $(1, 1)$ $(2, 2)$ $(3, 3)$ lie in R_1 .

Now, $(1, 2) \in R_1, (2, 3) \in R_1 \Rightarrow (1, 3) \in R_1$

Hence, R_1 is also transitive but $(1, 2) \in R_1 \Rightarrow (2, 1) \notin R_1$.

So, it is not symmetric.

(ii) Let $R_2 = \{(1, 2), (2, 1)\}$

Now, $(1, 2) \in R_2, (2, 1) \in R_2$

So, it is symmetric.

(iii) Let $R_3 = \{(1, 2), (2, 1), (1, 1), (2, 2), (3, 3), (1, 3), (3, 1), (2, 3)\}$

Hence, R_3 is reflexive, symmetric and transitive.

Q. 17 Let R be relation defined on the set of natural number N as follows, $R = \{(x, y) : x \in N, y \in N, 2x + y = 41\}$. Find the domain and range of the relation R . Also verify whether R is reflexive, symmetric and transitive.

Sol. Given that, $R = \{(x, y) : x \in N, y \in N, 2x + y = 41\}$.
 Domain = $\{1, 2, 3, \dots, 20\}$
 Range = $\{1, 3, 5, 7, \dots, 39\}$
 $R = \{(1, 39), (2, 37), (3, 35), \dots, (19, 3), (20, 1)\}$
 R is not reflexive as $(2, 2) \notin R$
 $2 \times 2 + 2 \neq 41$
 So, R is not symmetric.
 As $(1, 39) \in R$ but $(39, 1) \notin R$
 So, R is not transitive.
 As $(11, 19) \in R, (19, 3) \in R$
 But $(11, 3) \notin R$
 Hence, R is neither reflexive, nor symmetric and nor transitive.

Q. 18 Given, $A = \{2, 3, 4\}, B = \{2, 5, 6, 7\}$. Construct an example of each of the following

- an injective mapping from A to B .
- a mapping from A to B which is not injective.
- a mapping from B to A .

Sol. Given that, $A = \{2, 3, 4\}, B = \{2, 5, 6, 7\}$
 (i) Let $f : A \rightarrow B$ denote a mapping
 $f = \{(x, y) : y = x + 3\}$
 i.e., $f = \{(2, 5), (3, 6), (4, 7)\}$, which is an injective mapping.
 (ii) Let $g : A \rightarrow B$ denote a mapping such that $g = \{(2, 2), (3, 5), (4, 5)\}$, which is not an injective mapping.
 (iii) Let $h : B \rightarrow A$ denote a mapping such that $h = \{(2, 2), (5, 3), (6, 4), (7, 4)\}$, which is a mapping from B to A .

Q. 19 Give an example of a map

- which is one-one but not onto.
- which is not one-one but onto.
- which is neither one-one nor onto.

Sol. (i) Let $f : N \rightarrow N$, be a mapping defined by $f(x) = 2x$ which is one-one.
 For $f(x_1) = f(x_2)$
 $\Rightarrow 2x_1 = 2x_2$
 $x_1 = x_2$
 Further f is not onto, as for $1 \in N$, there does not exist any x in N such that $f(x) = 2x = 1$.
 (ii) Let $f : N \rightarrow N$, given by $f(1) = f(2) = 1$ and $f(x) = x - 1$ for every $x > 2$ is onto but not one-one. f is not one-one as $f(1) = f(2) = 1$. But f is onto.
 (iii) The mapping $f : R \rightarrow R$ defined as $f(x) = x^2$, is neither one-one nor onto.

Q. 20 Let $A = \mathbb{R} - \{3\}$, $B = \mathbb{R} - \{1\}$. If $f : A \rightarrow B$ be defined by $f(x) = \frac{x-2}{x-3}$,

$\forall x \in A$. Then, show that f is bijective.

Thinking Process

A function $f : x \rightarrow y$ is said to be bijective, if f is both one-one and onto.

Sol. Given that, $A = \mathbb{R} - \{3\}$, $B = \mathbb{R} - \{1\}$.

$f : A \rightarrow B$ is defined by $f(x) = \frac{x-2}{x-3}$, $\forall x \in A$

For injectivity

$$\text{Let } f(x_1) = f(x_2) \Rightarrow \frac{x_1-2}{x_1-3} = \frac{x_2-2}{x_2-3}$$

$$\begin{aligned} \Rightarrow & (x_1-2)(x_2-3) = (x_2-2)(x_1-3) \\ \Rightarrow & x_1x_2 - 3x_1 - 2x_2 + 6 = x_1x_2 - 3x_2 - 2x_1 + 6 \\ \Rightarrow & -3x_1 - 2x_2 = -3x_2 - 2x_1 \\ \Rightarrow & -x_1 = -x_2 \Rightarrow x_1 = x_2 \end{aligned}$$

So, $f(x)$ is an injective function.

For surjectivity

$$\text{Let } y = \frac{x-2}{x-3} \Rightarrow x-2 = xy-3y$$

$$\Rightarrow x(1-y) = 2-3y \Rightarrow x = \frac{2-3y}{1-y}$$

$$\Rightarrow x = \frac{3y-2}{y-1} \in A, \forall y \in B \quad [\text{codomain}]$$

So, $f(x)$ is surjective function.

Hence, $f(x)$ is a bijective function.

Q. 21 Let $A = [-1, 1]$, then, discuss whether the following functions defined on A are one-one onto or bijective.

(i) $f(x) = \frac{x}{2}$

(ii) $g(x) = |x|$

(iii) $h(x) = x|x|$

(iv) $k(x) = x^2$

Sol. Given that, $A = [-1, 1]$

(i) $f(x) = \frac{x}{2}$

$$\begin{aligned} \text{Let } f(x_1) &= f(x_2) \\ \Rightarrow \frac{x_1}{2} &= \frac{x_2}{2} \Rightarrow x_1 = x_2 \end{aligned}$$

So, $f(x)$ is one-one.

Now, let $y = \frac{x}{2}$

$$\Rightarrow x = 2y \notin A, \forall y \in A$$

As for $y = 1 \in A$, $x = 2 \notin A$

So, $f(x)$ is not onto.

Also, $f(x)$ is not bijective as it is not onto.

(ii) $g(x) = |x|$

Let

$$g(x_1) = g(x_2)$$

\Rightarrow

$$|x_1| = |x_2| \Rightarrow x_1 = \pm x_2$$

So, $g(x)$ is not one-one.

Now,

$$y = |x| \Rightarrow x = \pm y \notin A, \forall y \in A$$

So, $g(x)$ is not onto, also, $g(x)$ is not bijective.

(iii) $h(x) = x|x|$

Let

$$h(x_1) = h(x_2)$$

\Rightarrow

$$x_1|x_1| = x_2|x_2| \Rightarrow x_1 = x_2$$

So, $h(x)$ is one-one.

Now, let

$$y = x|x|$$

\Rightarrow

$$y = x^2 \in A, \forall x \in A$$

So, $h(x)$ is onto also, $h(x)$ is a bijective.

(iv) $k(x) = x^2$

Let

$$k(x_1) = k(x_2)$$

\Rightarrow

$$x_1^2 = x_2^2 \Rightarrow x_1 = \pm x_2$$

Thus, $k(x)$ is not one-one.

Now, let

$$y = x^2$$

\Rightarrow

$$x = \sqrt{y} \notin A, \forall y \in A$$

As for $y = -1$, $x = \sqrt{-1} \notin A$

Hence, $k(x)$ is neither one-one nor onto.

Q. 22 Each of the following defines a relation of N

(i) x is greater than y , $x, y \in N$.

(ii) $x + y = 10$, $x, y \in N$.

(iii) xy is square of an integer $x, y \in N$.

(iv) $x + 4y = 10$, $x, y \in N$

Determine which of the above relations are reflexive, symmetric and transitive.

Sol. (i) x is greater than y , $x, y \in N$

$$(x, x) \in R$$

For xRx

$x > x$ is not true for any $x \in N$.

Therefore, R is not reflexive.

Let

$$(x, y) \in R \Rightarrow xRy$$

$$x > y$$

but $y > x$ is not true for any $x, y \in N$

Thus, R is not symmetric.

Let

$$xRy \text{ and } yRz$$

$$x > y \text{ and } y > z \Rightarrow x > z$$

\Rightarrow

$$xRz$$

So, R is transitive.

(ii) $x + y = 10, x, y \in N$

$$R = \{(x, y) : x + y = 10, x, y \in N\}$$

$$R = \{(1, 9), (2, 8), (3, 7), (4, 6), (5, 5), (6, 4), (7, 3), (8, 2), (9, 1)\} \quad (1, 1) \notin R$$

So, R is not reflexive.

$$(x, y) \in R \Rightarrow (y, x) \in R$$

Therefore, R is symmetric.

$$(1, 9) \in R, (9, 1) \in R \Rightarrow (1, 1) \notin R$$

Hence, R is not transitive.

(iii) Given xy , is square of an integer $x, y \in N$.

$$\Rightarrow R = \{(x, y) : xy \text{ is a square of an integer } x, y \in N\}$$

$$(x, x) \in R, \forall x \in N$$

As x^2 is square of an integer for any $x \in N$.

Hence, R is reflexive.

$$\text{If } (x, y) \in R \Rightarrow (y, x) \in R$$

Therefore, R is symmetric.

$$\text{If } (x, y) \in R, (y, z) \in R$$

So, xy is square of an integer and yz is square of an integer.

Let $xy = m^2$ and $yz = n^2$ for some $m, n \in Z$

$$x = \frac{m^2}{y} \text{ and } z = \frac{x^2}{y}$$

$$xz = \frac{m^2 n^2}{y^2}, \text{ which is square of an integer.}$$

So, R is transitive.

(iv) $x + 4y = 10, x, y \in N$

$$R = \{(x, y) : x + 4y = 10, x, y \in N\}$$

$$R = \{(2, 2), (6, 1)\}$$

$$(1, 1), (3, 3), \dots, \notin R$$

Thus, R is not reflexive.

$$(6, 1) \in R \text{ but } (1, 6) \notin R$$

Hence, R is not symmetric.

$$(x, y) \in R \Rightarrow x + 4y = 10 \text{ but } (y, z) \in R$$

$$y + 4z = 10 \Rightarrow (x, z) \in R$$

So, R is transitive.

Q. 23 Let $A = \{1, 2, 3, \dots, 9\}$ and R be the relation in $A \times A$ defined by $(a, b) R (c, d)$ if $a + d = b + c$ for $(a, b), (c, d)$ in $A \times A$. Prove that R is an equivalence relation and also obtain the equivalent class $[(2, 5)]$.

Sol. Given that, $A = \{1, 2, 3, \dots, 9\}$ and $(a, b) R (c, d)$ if $a + d = b + c$ for $(a, b) \in A \times A$ and $(c, d) \in A \times A$.

Let $(a, b) R (a, b)$

\Rightarrow

$$a + b = b + a, \forall a, b \in A$$

which is true for any $a, b \in A$.

Hence, R is reflexive.

Let $(a, b) R (c, d)$

$$a + d = b + c$$

$$c + b = d + a \Rightarrow (c, d) R (a, b)$$

So, R is symmetric.

$$\begin{aligned}
 \text{Let } & (a, b) R (c, d) \text{ and } (c, d) R (e, f) \\
 & a + d = b + c \text{ and } c + f = d + e \\
 & a + d = b + c \text{ and } d + e = c + f \\
 & (a + d) - (d + e) = (b + c) - (c + f) \\
 & (a - e) = b - f \\
 & a + f = b + e \\
 & (a, b) R (e, f)
 \end{aligned}$$

So, R is transitive.

Hence, R is an equivalence relation.

Now, equivalence class containing $[(2, 5)]$ is $\{(1, 4), (2, 5), (3, 6), (4, 7), (5, 8), (6, 9)\}$.

Q. 24 Using the definition, prove that the function $f : A \rightarrow B$ is invertible if and only if f is both one-one and onto.

Sol. A function $f : X \rightarrow Y$ is defined to be invertible, if there exist a function $g : Y \rightarrow X$ such that $g \circ f = I_X$ and $f \circ g = I_Y$. The function is called the inverse of f and is denoted by f^{-1} .

A function $f : X \rightarrow Y$ is invertible iff f is a bijective function.

Q. 25 Functions $f, g : R \rightarrow R$ are defined, respectively, by $f(x) = x^2 + 3x + 1$, $g(x) = 2x - 3$, find

(i) $f \circ g$ (ii) $g \circ f$ (iii) $f \circ f$ (iv) $g \circ g$

Sol. Given that, $f(x) = x^2 + 3x + 1$, $g(x) = 2x - 3$

$$\begin{aligned}
 \text{(i)} \quad f \circ g &= f\{g(x)\} = f(2x - 3) \\
 &= (2x - 3)^2 + 3(2x - 3) + 1 \\
 &= 4x^2 + 9 - 12x + 6x - 9 + 1 = 4x^2 - 6x + 1
 \end{aligned}$$

$$\begin{aligned}
 \text{(ii)} \quad g \circ f &= g\{f(x)\} = g(x^2 + 3x + 1) \\
 &= 2(x^2 + 3x + 1) - 3 \\
 &= 2x^2 + 6x + 2 - 3 = 2x^2 + 6x - 1
 \end{aligned}$$

$$\begin{aligned}
 \text{(iii)} \quad f \circ f &= f\{f(x)\} = f(x^2 + 3x + 1) \\
 &= (x^2 + 3x + 1)^2 + 3(x^2 + 3x + 1) + 1 \\
 &= x^4 + 9x^2 + 1 + 6x^3 + 6x + 2x^2 + 3x^2 + 9x + 3 + 1 \\
 &= x^4 + 6x^3 + 14x^2 + 15x + 5
 \end{aligned}$$

$$\begin{aligned}
 \text{(iv)} \quad g \circ g &= g\{g(x)\} = g(2x - 3) \\
 &= 2(2x - 3) - 3 \\
 &= 4x - 6 - 3 = 4x - 9
 \end{aligned}$$

Q. 26 Let $*$ be the binary operation defined on Q . Find which of the following binary operations are commutative

(i) $a * b = a - b$, $\forall a, b \in Q$ (ii) $a * b = a^2 + b^2$, $\forall a, b \in Q$
 (iii) $a * b = a + ab$, $\forall a, b \in Q$ (iv) $a * b = (a - b)^2$, $\forall a, b \in Q$

Sol. Given that $*$ be the binary operation defined on Q .

(i) $a * b = a - b$, $\forall a, b \in Q$ and $b * a = b - a$

So, $a * b \neq b * a$ [$\because b - a \neq a - b$]

Hence, $*$ is not commutative.

$$(ii) \quad \begin{aligned} a * b &= a^2 + b^2 \\ b * a &= b^2 + a^2 \end{aligned}$$

So, $*$ is commutative.

[since, '+' is on rational is commutative]

$$(iii) \quad \begin{aligned} a * b &= a + ab \\ b * a &= b + ab \end{aligned}$$

Clearly, $a + ab \neq b + ab$

So, $*$ is not commutative.

$$(iv) \quad \begin{aligned} a * b &= (a - b)^2, \forall a, b \in Q \\ b * a &= (b - a)^2 \end{aligned}$$

$$\therefore (a - b)^2 = (b - a)^2$$

Hence, $*$ is commutative.

Q. 27 If $*$ be binary operation defined on R by $a * b = 1 + ab, \forall a, b \in R$.

Then, the operation $*$ is

- (i) commutative but not associative.
- (ii) associative but not commutative.
- (iii) neither commutative nor associative.
- (iv) both commutative and associative.

Sol. (i) Given that, $a * b = 1 + ab, \forall a, b \in R$
 $a * b = ab + 1 = b * a$

So, $*$ is a commutative binary operation.

Also, $a * (b * c) = a * (1 + bc) = 1 + a(1 + bc)$

$$a * (b * c) = 1 + a + abc \quad \dots(i)$$

$$\begin{aligned} (a * b) * c &= (1 + ab) * c \\ &= 1 + (1 + ab)c = 1 + c + abc \end{aligned} \quad \dots(ii)$$

From Eqs. (i) and (ii),

$$a * (b * c) \neq (a * b) * c$$

So, $*$ is not associative

Hence, $*$ is commutative but not associative.

Objective Type Questions

Q. 28 Let T be the set of all triangles in the Euclidean plane and let a relation R on T be defined as aRb , if a is congruent to $b, \forall a, b \in T$.

Then, R is

- (a) reflexive but not transitive
- (b) transitive but not symmetric
- (c) equivalence
- (d) None of these

Sol. (c) Consider that aRb , if a is congruent to $b, \forall a, b \in T$.

Then, $aRa \Rightarrow a \cong a$,

which is true for all $a \in T$

So, R is reflexive, ... (i)

Let $aRb \Rightarrow a \cong b$
 $\Rightarrow b \cong a \Rightarrow bRa$
 $\Rightarrow bRa$
 So, R is symmetric. ... (ii)
 Let aRb and bRc
 $\Rightarrow a \cong b$ and $b \cong c$
 $\Rightarrow a \cong c \Rightarrow aRc$
 So, R is transitive. ... (iii)
 Hence, R is equivalence relation.

Q. 29 Consider the non-empty set consisting of children in a family and a relation R defined as aRb , if a is brother of b . Then, R is

- (a) symmetric but not transitive
- (b) transitive but not symmetric
- (c) neither symmetric nor transitive
- (d) both symmetric and transitive

Sol. (b) Given, $aRb \Rightarrow a$ is brother of b
 $\therefore aRa \Rightarrow a$ is brother of a , which is not true.
 So, R is not reflexive.

$$aRb \Rightarrow a \text{ is brother of } b.$$

This does not mean b is also a brother of a and b can be a sister of a .

Hence, R is not symmetric.

$$aRb \Rightarrow a \text{ is brother of } b$$

and $bRc \Rightarrow b$ is a brother of c .

So, a is brother of c .

Hence, R is transitive.

Q. 30 The maximum number of equivalence relations on the set $A = \{1, 2, 3\}$ are

- (a) 1
- (b) 2
- (c) 3
- (d) 5

Sol. (d) Given that, $A = \{1, 2, 3\}$
 Now, number of equivalence relations as follows

$$R_1 = \{(1, 1), (2, 2), (3, 3)\}$$

$$R_2 = \{(1, 1), (2, 2), (3, 3), (1, 2), (2, 1)\}$$

$$R_3 = \{(1, 1), (2, 2), (3, 3), (1, 3), (3, 1)\}$$

$$R_4 = \{(1, 1), (2, 2), (3, 3), (2, 3), (3, 2)\}$$

$$R_5 = \{(1, 2, 3) \Leftrightarrow A \times A = A^2\}$$

\therefore Maximum number of equivalence relation on the set $A = \{1, 2, 3\} = 5$

Q. 31 If a relation R on the set $\{1, 2, 3\}$ be defined by $R = \{(1, 2)\}$, then R is

- (a) reflexive
- (b) transitive
- (c) symmetric
- (d) None of these

Sol. (b) R on the set $\{1, 2, 3\}$ be defined by $R = \{(1, 2)\}$
 It is clear that R is transitive.

Q. 32 Let us define a relation R in R as aRb if $a \geq b$. Then, R is

- (a) an equivalence relation
- (b) reflexive, transitive but not symmetric
- (c) symmetric, transitive but not reflexive
- (d) neither transitive nor reflexive but symmetric

Sol. (b) Given that, aRb if $a \geq b$
 $\Rightarrow aRa \Rightarrow a \geq a$ which is true.
 Let aRb , $a \geq b$, then $b \geq a$ which is not true R is not symmetric.
 But aRb and bRc
 $\Rightarrow a \geq b$ and $b \geq c$
 $\Rightarrow a \geq c$
 Hence, R is transitive.

Q. 33 If $A = \{1, 2, 3\}$ and consider the relation

$$R = \{(1, 1), (2, 2), (3, 3), (1, 2), (2, 3), (1, 3)\}$$

Then, R is

- (a) reflexive but not symmetric
- (b) reflexive but not transitive
- (c) symmetric and transitive
- (d) neither symmetric nor transitive

Sol. (a) Given that, $A = \{1, 2, 3\}$
 and $R = \{(1, 1), (2, 2), (3, 3), (1, 2), (2, 3), (1, 3)\}$
 $\because (1, 1), (2, 2), (3, 3) \in R$
 Hence, R is reflexive.
 $(1, 2) \in R$ but $(2, 1) \notin R$
 Hence, R is not symmetric.
 $(1, 2) \in R$ and $(2, 3) \in R$
 $\Rightarrow (1, 3) \in R$
 Hence, R is transitive.

Q. 34 The identity element for the binary operation $*$ defined on $Q - \{0\}$ as

$$a * b = \frac{ab}{2}, \forall a, b \in Q - \{0\} \text{ is}$$

- (a) 1
- (b) 0
- (c) 2
- (d) None of these

Thinking Process

For given binary operation $*$: $A \times A \rightarrow A$, an element $e \in A$, if it exists, is called identity for the operation $*$, if $a * e = a = e * a, \forall a \in A$.

Sol. (c) Given that, $a * b = \frac{ab}{2}, \forall a, b \in Q - \{0\}$.

Let e be the identity element for $*$.

$$\therefore a * e = \frac{ae}{2}$$

$$\Rightarrow a = \frac{ae}{2} \Rightarrow e = 2$$



Q. 35 If the set A contains 5 elements and the set B contains 6 elements, then the number of one-one and onto mappings from A to B is

- (a) 720 (b) 120 (c) 0 (d) None of these

Sol. (c) We know that, if A and B are two non-empty finite set containing m and n elements respectively, then the number of one-one and onto mapping from A to B is

$$n! \text{ if } m = n$$

$$0 \text{ if } m \neq n$$

Given that,

$$m = 5 \text{ and } n = 6$$

\therefore

$$m \neq n$$

Number of mapping = 0

Q. 36 If $A = \{1, 2, 3, \dots, n\}$ and $B = \{a, b\}$. Then, the number of surjections from A into B is

- (a) nP_2 (b) $2^n - 2$ (c) $2^n - 1$ (d) None of these

Sol. (d) Given that, $A = \{1, 2, 3, \dots, n\}$ and $B = \{a, b\}$.

We know that, if A and B are two non-empty finite sets containing m and n elements respectively, then the number of surjection from A into B is

$${}^nC_m \times m!, \text{ if } n \geq m$$

$$0 \text{ if } n < m$$

Here, $m = 2$

$$\begin{aligned} \therefore \text{Number of surjection from } A \text{ into } B &= {}^nC_2 \times 2! = \frac{n!}{2!(n-2)!} \times 2! \\ &= \frac{n(n-1)(n-2)!}{2 \times 1(n-2)!} \times 2! = n^2 - n \end{aligned}$$

Q. 37 If $f : R \rightarrow R$ be defined by $f(x) = \frac{1}{x}, \forall x \in R$. Then, f is

- (a) one-one (b) onto (c) bijective (d) f is not defined

Thinking Process

In the given function at $x=0$, $f(x) = \infty$. So, the function is not define.

Sol. (d) Given that, $f(x) = \frac{1}{x}, \forall x \in R$

For

$$x = 0,$$

$f(x)$ is not defined.

Hence, $f(x)$ is a not define function.

Q. 38 If $f : R \rightarrow R$ be defined by $f(x) = 3x^2 - 5$ and $g : R \rightarrow R$ by

$$g(x) = \frac{x}{x^2 + 1}. \text{ Then, } gof \text{ is}$$

(a) $\frac{3x^2 - 5}{9x^4 - 30x^2 + 26}$

(b) $\frac{3x^2 - 5}{9x^4 - 6x^2 + 26}$

(c) $\frac{3x^2}{x^4 + 2x^2 - 4}$

(d) $\frac{3x^2}{9x^4 + 30x^2 - 2}$



Sol. (a) Given that, $f(x) = 3x^2 - 5$ and $g(x) = \frac{x}{x^2 + 1}$

$$\begin{aligned}gof &= g\{f(x)\} = g(3x^2 - 5) \\&= \frac{3x^2 - 5}{(3x^2 - 5)^2 + 1} = \frac{3x^2 - 5}{9x^4 - 30x^2 + 25 + 1} \\&= \frac{3x^2 - 5}{9x^4 - 30x^2 + 26}\end{aligned}$$

Q. 39 Which of the following functions from Z into Z are bijections?

- (a) $f(x) = x^3$ (b) $f(x) = x + 2$ (c) $f(x) = 2x + 1$ (d) $f(x) = x^2 + 1$

Sol. (b) Here,

$$f(x) = x + 2 \Rightarrow f(x_1) = f(x_2)$$

$$x_1 + 2 = x_2 + 2 \Rightarrow x_1 = x_2$$

Let

$$y = x + 2$$

$$x = y - 2 \in Z, \forall y \in x$$

Hence, $f(x)$ is one-one and onto.

Q. 40 If $f : R \rightarrow R$ be the functions defined by $f(x) = x^3 + 5$, then $f^{-1}(x)$ is

- (a) $(x + 5)^{\frac{1}{3}}$ (b) $(x - 5)^{\frac{1}{3}}$ (c) $(5 - x)^{\frac{1}{3}}$ (d) $5 - x$

Sol. (b) Given that,

$$f(x) = x^3 + 5$$

Let

$$y = x^3 + 5 \Rightarrow x^3 = y - 5$$

$$x = (y - 5)^{\frac{1}{3}} \Rightarrow f(x)^{-1} = (x - 5)^{\frac{1}{3}}$$

Q. 41 If $f : A \rightarrow B$ and $g : B \rightarrow C$ be the bijective functions, then $(gof)^{-1}$ is

- (a) $f^{-1}og^{-1}$ (b) fog (c) $g^{-1}of^{-1}$ (d) gof

Sol. (a) Given that, $f : A \rightarrow B$ and $g : B \rightarrow C$ be the bijective functions.

$$(gof)^{-1} = f^{-1}og^{-1}$$

Q. 42 If $f : R - \left\{\frac{3}{5}\right\} \rightarrow R$ be defined by $f(x) = \frac{3x + 2}{5x - 3}$, then

- (a) $f^{-1}(x) = f(x)$ (b) $f^{-1}(x) = -f(x)$ (c) $(fof)x = -x$ (d) $f^{-1}(x) = \frac{1}{19}f(x)$

Sol. (a) Given that,

$$f(x) = \frac{3x + 2}{5x - 3}$$

Let

$$y = \frac{3x + 2}{5x - 3}$$

$$3x + 2 = 5xy - 3y \Rightarrow x(3 - 5y) = -3y - 2$$

$$x = \frac{3y + 2}{5y - 3} \Rightarrow f^{-1}(x) = \frac{3x + 2}{5x - 3}$$

\therefore

$$f^{-1}(x) = f(x)$$

Q. 43 If $f: [0, 1] \rightarrow [0, 1]$ be defined by $f(x) = \begin{cases} x, & \text{if } x \text{ is rational} \\ 1-x, & \text{if } x \text{ is irrational} \end{cases}$

then $(f \circ f)x$ is

- (a) constant (b) $1+x$ (c) x (d) None of these

Sol. (c) Given that, $f: [0, 1] \rightarrow [0, 1]$ be defined by

$$f(x) = \begin{cases} x, & \text{if } x \text{ is rational} \\ 1-x, & \text{if } x \text{ is irrational} \end{cases}$$

$$\therefore (f \circ f)x = f(f(x)) = x$$

Q. 44 If $f: [2, \infty) \rightarrow R$ be the function defined by $f(x) = x^2 - 4x + 5$, then the range of f is

- (a) R (b) $[1, \infty)$ (c) $[4, \infty)$ (d) $[5, \infty)$

💡 **Thinking Process**

$$\text{Range of } f = \{y \in Y : y = f(x) \text{ for some } x \in X\}$$

Sol. (b) Given that, $f(x) = x^2 - 4x + 5$

Let

$$y = x^2 - 4x + 5$$

\Rightarrow

$$y = x^2 - 4x + 4 + 1 = (x-2)^2 + 1$$

\Rightarrow

$$(x-2)^2 = y-1 \Rightarrow x-2 = \sqrt{y-1}$$

\Rightarrow

$$x = 2 + \sqrt{y-1}$$

\therefore

$$y-1 \geq 0, y \geq 1$$

$$\text{Range} = [1, \infty)$$

Q. 45 If $f: N \rightarrow R$ be the function defined by $f(x) = \frac{2x-1}{2}$ and $g: Q \rightarrow R$

be another function defined by $g(x) = x + 2$. Then, $(g \circ f)\frac{3}{2}$ is

- (a) 1 (b) 1 (c) $\frac{7}{2}$ (d) None of these

Sol. (d) Given that, $f(x) = \frac{2x-1}{2}$ and $g(x) = x + 2$

$$\begin{aligned} (g \circ f)\frac{3}{2} &= g\left[f\left(\frac{3}{2}\right)\right] = g\left(\frac{2 \times \frac{3}{2} - 1}{2}\right) \\ &= g(1) = 1 + 2 = 3 \end{aligned}$$

Q. 46 If $f : R \rightarrow R$ be defined by $f(x) = \begin{cases} 2x : x > 3 \\ x^2 : 1 < x \leq 3 \\ 3x : x \leq 1 \end{cases}$

Then, $f(-1) + f(2) + f(4)$ is

- (a) 9 (b) 14 (c) 5 (d) None of these

Sol. (a) Given that, $f(x) = \begin{cases} 2x : x > 3 \\ x^2 : 1 < x \leq 3 \\ 3x : x \leq 1 \end{cases}$

$$\begin{aligned} f(-1) + f(2) + f(4) &= 3(-1) + (2)^2 + 2 \times 4 \\ &= -3 + 4 + 8 = 9 \end{aligned}$$

Q. 47 If $f : R \rightarrow R$ be given by $f(x) = \tan x$, then $f^{-1}(1)$ is

- (a) $\frac{\pi}{4}$ (b) $\left\{n\pi + \frac{\pi}{4} : n \in Z\right\}$
(c) Does not exist (d) None of these

Sol. (a) Given that,

$$f(x) = \tan x$$

Let

$$y = \tan x \Rightarrow x = \tan^{-1} y$$

\Rightarrow

$$f^{-1}(x) = \tan^{-1} x \Rightarrow f^{-1}(1) = \tan^{-1} 1$$

\Rightarrow

$$= \tan^{-1} \tan \frac{\pi}{4} = \frac{\pi}{4}$$

$$\left[\because \tan \frac{\pi}{4} = 1 \right]$$

Fillers

Q. 48 Let the relation R be defined in N by aRb , if $2a + 3b = 30$. Then, $R = \dots$

Sol. Given that,

$$2a + 3b = 30$$

$$3b = 30 - 2a$$

$$b = \frac{30 - 2a}{3}$$

For

$$a = 3, b = 8$$

$$a = 6, b = 6$$

$$a = 9, b = 4$$

$$a = 12, b = 2$$

$$R = \{(3, 8), (6, 6), (9, 4), (12, 2)\}$$

Q. 49 If the relation R be defined on the set $A = \{1, 2, 3, 4, 5\}$ by $R = \{(a, b) : |a^2 - b^2| < 8\}$. Then, R is given by

Sol. Given,

$$A = \{1, 2, 3, 4, 5\},$$

$$R = \{(a, b) : |a^2 - b^2| < 8\}$$

$$R = \{(1, 1), (1, 2), (2, 1), (2, 2), (2, 3), (3, 2), (3, 3), (4, 3), (3, 4), (4, 4), (5, 5)\}$$



Q. 50 If $f = \{(1, 2), (3, 5), (4, 1)\}$ and $g = \{(2, 3), (5, 1), (1, 3)\}$, then
 $gof = \dots\dots\dots$ and $fog = \dots\dots\dots$.

Sol. Given that,
 $f = \{(1, 2), (3, 5), (4, 1)\}$ and $g = \{(2, 3), (5, 1), (1, 3)\}$
 $gof(1) = g\{f(1)\} = g(2) = 3$
 $gof(3) = g\{f(3)\} = g(5) = 1$
 $gof(4) = g\{f(4)\} = g(1) = 3$
 $gof = \{(1, 3), (3, 1), (4, 3)\}$
Now,
 $fog(2) = f\{g(2)\} = f(3) = 5$
 $fog(5) = f\{g(5)\} = f(1) = 2$
 $fog(1) = f\{g(1)\} = f(3) = 5$
 $fog = \{(2, 5), (5, 2), (1, 5)\}$

Q. 51 If $f : R \rightarrow R$ be defined by $f(x) = \frac{x}{\sqrt{1+x^2}}$, then $(fofof)(x) = \dots\dots\dots$.

Sol. Given that,
 $f(x) = \frac{x}{\sqrt{1+x^2}}$
 $(fofof)(x) = f[f\{f(x)\}]$
 $= f\left[f\left(\frac{x}{\sqrt{1+x^2}}\right)\right] = f\left(\frac{\frac{x}{\sqrt{1+x^2}}}{\sqrt{1+\frac{x^2}{1+x^2}}}\right)$
 $= f\left[\frac{x\sqrt{1+x^2}}{\sqrt{1+x^2}(\sqrt{2x^2+1})}\right] = f\left(\frac{x}{\sqrt{1+2x^2}}\right)$
 $= \frac{\frac{x}{\sqrt{1+2x^2}}}{\sqrt{1+\frac{x^2}{1+2x^2}}} = \frac{x\sqrt{1+2x^2}}{\sqrt{1+2x^2}\sqrt{1+3x^2}}$
 $= \frac{x}{\sqrt{1+3x^2}} = \frac{x}{\sqrt{3x^2+1}}$

Q. 52 If $f(x) = [4 - (x - 7)^3]$, then $f^{-1}(x) = \dots\dots\dots$.

Sol. Given that,
 $f(x) = [4 - (x - 7)^3]$
Let
 $y = [4 - (x - 7)^3]$
 $(x - 7)^3 = 4 - y$
 $(x - 7) = (4 - y)^{1/3}$
 $x = 7 + (4 - y)^{1/3}$
 \Rightarrow
 $f^{-1}(x) = 7 + (4 - x)^{1/3}$

True/False

Q. 53 Let $R = \{(3, 1), (1, 3), (3, 3)\}$ be a relation defined on the set $A = \{1, 2, 3\}$. Then, R is symmetric, transitive but not reflexive.

Sol. False

Given that, $R = \{(3, 1), (1, 3), (3, 3)\}$ be defined on the set $A = \{1, 2, 3\}$

$$(1, 1) \notin R$$

So, R is not reflexive.

$$(3, 1) \in R, (1, 3) \in R$$

Hence, R is symmetric.

Since,

$$(3, 1) \in R, (1, 3) \in R$$

But

$$(1, 1) \notin R$$

Hence, R is not transitive.

Q. 54 If $f : R \rightarrow R$ be the function defined by $f(x) = \sin(3x + 2) \forall x \in R$. Then, f is invertible.

Sol. False

Given that, $f(x) = \sin(3x + 2), \forall x \in R$ is not one-one function for all $x \in R$.

So, f is not invertible.

Q. 55 Every relation which is symmetric and transitive is also reflexive.

Sol. False

Let R be a relation defined by

$$R = \{(1, 2), (2, 1), (1, 1), (2, 2)\} \text{ on the set } A = \{1, 2, 3\}$$

It is clear that $(3, 3) \notin R$. So, it is not reflexive.

Q. 56 An integer m is said to be related to another integer n , if m is a integral multiple of n . This relation in Z is reflexive, symmetric and transitive.

Sol. False

The given relation is reflexive and transitive but not symmetric.

Q. 57 If $A = \{0, 1\}$ and N be the set of natural numbers. Then, the mapping $f : N \rightarrow A$ defined by $f(2n - 1) = 0, f(2n) = 1, \forall n \in N$, is onto.

Sol. True

Given,

$$A = \{0, 1\}$$

$$f(2n - 1) = 0, f(2n) = 1, \forall n \in N$$

So, the mapping $f : N \rightarrow A$ is onto.

Q. 58 The relation R on the set $A = \{1, 2, 3\}$ defined as $R = \{(1, 1), (1, 2), (2, 1), (3, 3)\}$ is reflexive, symmetric and transitive.

Sol. False

Given that,

$$R = \{(1, 1), (1, 2), (2, 1), (3, 3)\}$$

$$(2, 2) \notin R$$

So, R is not reflexive.

Q. 59 The composition of function is commutative.

Sol. *False*

Let

$$f(x) = x^2$$

and

$$g(x) = x + 1$$

$$fog(x) = f\{g(x)\} = f(x + 1)$$

$$= (x + 1)^2 = x^2 + 2x + 1$$

$$gof(x) = g\{f(x)\} = g(x^2) = x^2 + 1$$

\therefore

$$fog(x) \neq gof(x)$$

Q. 60 The composition of function is associative.

Sol. *True*

Let

$$f(x) = x, g(x) = x + 1$$

and

$$h(x) = 2x - 1$$

Then,

$$fo\{goh(x)\} = f\{g\{h(x)\}\}$$

$$= f\{g(2x - 1)\}$$

$$= f(2x - 1 + 1)$$

$$= f(2x) = 2x$$

\therefore

$$(fog)oh(x) = (fog)\{h(x)\}$$

$$= (fog)(2x - 1)$$

$$= f\{g(2x - 1)\}$$

$$= f(2x - 1 + 1)$$

$$= f(2x) = 2x$$

Q. 61 Every function is invertible.

Sol. *False*

Only bijective functions are invertible.

Q. 62 A binary operation on a set has always the identity element.

Sol. *False*

'+' is a binary operation on the set N but it has no identity element.