

20. Definite Integrals

Exercise 20.1

1. Question

Evaluate the following definite integrals:

$$\int_4^9 \frac{1}{\sqrt{x}} dx$$

Answer

Using the formula:

$$\begin{aligned} \int_a^b (x)^n dx &= \left[\frac{(x)^{n+1}}{n+1} \right]_a^b \\ \Rightarrow \int_4^9 (x)^{-\frac{1}{2}} dx &= \left[\frac{(x)^{-\frac{1}{2}+1}}{-\frac{1}{2}+1} \right]_4^9 \\ &= \left[\frac{(x)^{\frac{1}{2}}}{\frac{1}{2}} \right]_4^9 \\ &= \left[\frac{(9)^{\frac{1}{2}}}{\frac{1}{2}} \right] - \left[\frac{(4)^{\frac{1}{2}}}{\frac{1}{2}} \right] = \left[\frac{3}{\frac{1}{2}} \right] - \left[\frac{2}{\frac{1}{2}} \right] \\ &\Rightarrow 3 \times 2 - 2 \times 2 \\ &\Rightarrow 6 - 4 = 2 \end{aligned}$$

$$\Rightarrow \int_4^9 (x)^{-\frac{1}{2}} dx = 2$$

2. Question

Evaluate the following definite integrals:

$$\int_{-2}^3 \frac{1}{x+7} dx$$

Answer

Using the formula:

$$\begin{aligned} \int_a^b \left(\frac{1}{x} \right) dx &= [\log|x|]_a^b \\ \Rightarrow \int_{-2}^3 \left(\frac{1}{x+7} \right) dx &= [\log|x+7|]_{-2}^3 \\ &\Rightarrow \log 3 + 7 - \log(-2+7) \\ &\Rightarrow \log 10 - \log 5 \end{aligned}$$

$$\Rightarrow \log\left|\frac{10}{5}\right| = \log|2|$$

$$\int_a^b \left(\frac{1}{x}\right) dx = \log|2|$$

3. Question

Evaluate the following definite integrals:

$$\int_0^{1/2} \frac{1}{\sqrt{1-x^2}} dx$$

Answer

Using the formula:

$$\int_a^b \frac{1}{\sqrt{(1-x^2)}} dx = [-\cos^{-1}x]_a^b$$

$$\int_0^{1/2} \frac{1}{\sqrt{(1-x^2)}} = [-\cos^{-1}x]_0^{1/2}$$

$$\Rightarrow -[\cos^{-1} 1/2 - \cos^{-1} 0]$$

$$\Rightarrow -\left[\frac{\pi}{3} - \frac{\pi}{2}\right]$$

$$\Rightarrow \left[\frac{\pi}{6}\right]$$

$$\int_0^{1/2} \frac{1}{\sqrt{(1-x^2)}} = \left[\frac{\pi}{6}\right]$$

4. Question

Evaluate the following definite integrals:

$$\int_0^1 \frac{1}{1+x^2} dx$$

Answer

Using the formula:

$$\int_a^b \frac{1}{(1+x^2)} dx = [\tan^{-1}x]_a^b$$

$$\int_0^1 \frac{1}{(1+x^2)} dx = [\tan^{-1}x]_0^1$$

$$\Rightarrow [\tan^{-1} 1 - \tan^{-1} 0]$$

$$\Rightarrow \left[\frac{\pi}{4} - 0\right]$$

$$\Rightarrow \pi/4$$

$$\int_a^b \frac{1}{(1+x^2)} dx = \pi/4$$

5. Question

Evaluate the following definite integrals:

$$\int_2^3 \frac{x}{x^2 + 1} dx$$

Answer

Let $x^2 + 1 = t$

\Rightarrow On differentiation, we get

$$2x dx = dt$$

$$\Rightarrow x dx = \frac{dt}{2}$$

\Rightarrow Hence the question will become:

$$\int_2^3 \frac{dt/2}{t} = \int_2^3 \frac{dt}{2t}$$

$$\Rightarrow \frac{1}{2} \int_2^3 \frac{dt}{t}$$

Using the formula:

$$\int_a^b \left(\frac{1}{x}\right) dx = [\log|x|]_a^b$$

$$1/2 \int_2^3 \left(\frac{1}{t}\right) dt = [\log|t|]_2^3$$

$$\Rightarrow [\log|x^2 + 1|]_2^3$$

$$\Rightarrow \log|3^2 + 1| - \log|2^2 + 1|$$

$$\Rightarrow \log|10| - \log|5|$$

$$\Rightarrow \log|10/5|$$

$$\Rightarrow \log|2|$$

6. Question

Evaluate the following definite integrals:

$$\int_0^\infty \frac{1}{a^2 + b^2 x^2} dx$$

Answer

$$\Rightarrow \int_0^\infty \frac{1}{(a^2 + b^2 x^2)} dx = \int_0^\infty \frac{1}{b^2((\frac{a}{b})^2 + x^2)} dx$$



$$\Rightarrow \int_0^{\infty} \frac{1}{(a^2 + b^2 x^2)} dx = 1/b^2 \int_0^{\infty} \frac{1}{(\left(\frac{a}{b}\right)^2 + x^2)} dx$$

Now, Using the formula:

$$\int_a^b \frac{1}{(a^2 + x^2)} dx = \frac{1}{a} [\tan^{-1} x/a]_a^b$$

$$\Rightarrow \int_0^{\infty} \frac{1}{(a^2 + b^2 x^2)} dx = \frac{1}{b} [\tan^{-1} \frac{x}{b}]_0^{\infty}$$

$$\Rightarrow \int_0^{\infty} \frac{1}{(a^2 + b^2 x^2)} dx = \left(\frac{b}{a}\right) [\tan^{-1} \frac{bx}{a}]_0^{\infty}$$

$$\Rightarrow \int_0^{\infty} \frac{1}{(a^2 + b^2 x^2)} dx = \left(\frac{b}{a}\right) [\tan^{-1} \frac{b \times \infty}{a} - \tan^{-1} b \times \frac{0}{a}]$$

$$\Rightarrow \int_0^{\infty} \frac{1}{(a^2 + b^2 x^2)} dx = \left(\frac{b}{a}\right) [\tan^{-1} \infty - \tan^{-1} 0]$$

$$\Rightarrow \int_0^{\infty} \frac{1}{(a^2 + b^2 x^2)} dx = \left(\frac{b}{a}\right) \left[\frac{\pi}{2} - 0\right]$$

$$\Rightarrow \int_0^{\infty} \frac{1}{(a^2 + b^2 x^2)} dx = \left(\frac{b}{a}\right) \left(\frac{\pi}{2}\right)$$

7. Question

Evaluate the following definite integrals:

$$\int_{-1}^1 \frac{1}{1+x^2} dx$$

Answer

Using the formula:

$$\int_a^b \frac{1}{(1+x^2)} dx = [\tan^{-1} x]_a^b$$

$$\Rightarrow \int_{-1}^1 \frac{1}{(1+x^2)} dx = [\tan^{-1} x]_{-1}^1$$

$$\Rightarrow [\tan^{-1} (1) - \tan^{-1} (-1)]$$

$$\Rightarrow [\frac{\pi}{4} - \left(-\frac{\pi}{4}\right)]$$

$$\Rightarrow \frac{\pi}{4} + \frac{\pi}{4} = \frac{\pi}{2}$$

8. Question

Evaluate the following definite integrals:

$$\int_0^{\infty} e^{-x} dx$$

Answer

Using the formula:

$$\int_a^b e^x dx = [e^x]_a^b$$

$$\Rightarrow \int_0^\infty e^{-x} dx = -[e^{-x}]_0^\infty$$

$$\Rightarrow \int_0^\infty e^{-x} dx = -[e^{-\infty} - e^0]$$

$$\Rightarrow \int_0^\infty e^{-x} dx = -[0 - 1]$$

$$\Rightarrow \int_0^\infty e^{-x} dx = 1$$

9. Question

Evaluate the following definite integrals:

$$\int_0^1 \frac{x}{x+1} dx$$

Answer

$$\Rightarrow \int_0^1 \left(\frac{x}{x+1} \right) dx = \int_0^1 \left(\frac{x+1-1}{x+1} \right) dx$$

$$\Rightarrow \int_0^1 \left(\frac{x}{x+1} \right) dx = \int_0^1 \left(\frac{x+1}{x+1} \right) dx - \int_0^1 \left(\frac{1}{x+1} \right) dx$$

$$\Rightarrow \int_0^1 \left(\frac{x}{x+1} \right) dx = \int_0^1 1 dx - \int_0^1 \left(\frac{1}{x+1} \right) dx$$

Using the formula:

$$\int_a^b \left(\frac{1}{x} \right) dx = [\log|x|]_a^b$$

$$\Rightarrow \int_0^1 \left(\frac{x}{x+1} \right) dx = [x]_0^1 - [\log|x+1|]_0^1$$

$$\Rightarrow \int_0^1 \left(\frac{x}{x+1} \right) dx = [1 - 0] - [\log|1+1| - \log|0+1|]$$

$$\Rightarrow \int_0^1 \left(\frac{x}{x+1} \right) dx = [1] - [\log|2|] - 0$$

($\because \log 1 = 0$)

$$\Rightarrow \int_0^1 \left(\frac{x}{x+1} \right) dx = [1] - [\log|2|]$$

10. Question

Evaluate the following definite integrals:

$$\int_0^{\pi/2} (\sin x + \cos x) dx$$

Answer

$$\Rightarrow \int_0^{\pi/2} (\sin(x) + \cos(x)) dx = \int_0^{\pi/2} \sin(x) dx + \int_0^{\pi/2} \cos(x) dx$$

Using the formula:

$$\int_a^b \sin(x) dx = -[\cos(x)]_a^b$$

and

$$\int_a^b \cos(x) dx = [\sin(x)]_a^b$$

$$\Rightarrow \int_0^{\pi/2} (\sin(x) + \cos(x)) dx = -[\cos(x)]_0^{\pi/2} + [\sin(x)]_0^{\pi/2}$$

$$\Rightarrow \int_0^{\pi/2} (\sin(x) + \cos(x)) dx = -\left[\cos\left(\frac{\pi}{2}\right) - \cos(0) \right] + \left[\sin\left(\frac{\pi}{2}\right) - \sin(0) \right]$$

$$\Rightarrow \int_0^{\pi/2} (\sin(x) + \cos(x)) dx = -[0 - 1] + [1 - 0]$$

$$\Rightarrow \int_0^{\pi/2} (\sin(x) + \cos(x)) dx = -[-1] + [1]$$

$$\Rightarrow \int_0^{\pi/2} (\sin(x) + \cos(x)) dx = 2$$

11. Question

Evaluate the following definite integrals:

$$\int_{\pi/4}^{\pi/2} \cot x dx$$

Answer

Using the formula:

$$\int_a^b \cot(x) dx = [\log|\sin(x)|]_a^b$$

$$\Rightarrow \int_{\pi/4}^{\pi/2} \cot(x) dx = [\log|\sin(x)|]_{\pi/4}^{\pi/2}$$

$$\Rightarrow \int_{\pi/4}^{\pi/2} \cot(x) dx = \left[\log \left| \sin \left(\frac{\pi}{2} \right) \right| - \log \left| \sin \left(\frac{\pi}{4} \right) \right| \right]$$

$$\Rightarrow \int_{\pi/4}^{\pi/2} \cot(x) dx = [\log|1| - \log|1/\sqrt{2}|]$$

$$\Rightarrow \int_{\pi/4}^{\pi/2} \cot(x) dx = \left[\log \left| 1 / (\frac{1}{\sqrt{2}}) \right| \right]$$

$$\Rightarrow \int_{\pi/4}^{\pi/2} \cot(x) dx = [\log \sqrt{2}]$$

12. Question

Evaluate the following definite integrals:

$$\int_0^{\pi/4} \sec x dx$$

Answer

Using the formula:

$$\int_a^b \sec(x) dx = [\log|\sec(x) + \tan(x)|]_a^b$$

$$\Rightarrow \int_0^{\pi/4} \sec(x) dx = [\log|\sec(x) + \tan(x)|]_0^{\pi/4}$$

$$\Rightarrow \int_0^{\pi/4} \sec(x) dx = [\log \left| \sec \left(\frac{\pi}{4} \right) + \tan \left(\frac{\pi}{4} \right) \right| - \log|\sec(0) + \tan(0)|]$$

$$\Rightarrow \int_0^{\pi/4} \sec(x) dx = [\log|\sqrt{2} + 1|] - \log|1 + 0|$$

$$\Rightarrow \int_0^{\pi/4} \sec(x) dx = [\log|\sqrt{2} + 1|] - \log|1|$$

$$\Rightarrow \int_0^{\pi/4} \sec(x) dx = [\log|\sqrt{2} + 1|] - 0$$

($\because \log 1 = 0$)

$$\Rightarrow \int_0^{\pi/4} \sec(x) dx = [\log|\sqrt{2} + 1|]$$

13. Question

Evaluate the following definite integrals:

$$\int_{\pi/6}^{\pi/4} \csc(x) dx$$

Answer

Using the formula :

$$\int_a^b \csc(x) dx = [\log|\csc(x) - \cot(x)|]_a^b$$

$$\Rightarrow \int_{\pi/6}^{\pi/4} \csc(x) dx = [\log|\csc(x) - \cot(x)|]_{\pi/6}^{\pi/4}$$

$$\Rightarrow \int_{\pi/6}^{\pi/4} \csc(x) dx = [\log|\csc(\frac{\pi}{4}) - \cot(\frac{\pi}{4})| - \log|\csc(\frac{\pi}{6}) - \cot(\frac{\pi}{6})|]$$

$$\Rightarrow \int_{\pi/6}^{\pi/4} \csc(x) dx = [\log|\sqrt{2} - 1| - \log|2 - \sqrt{3}|]$$

$$(\because \csc(\frac{\pi}{4}) = \sqrt{2}, \cot(\frac{\pi}{4}) = 1, \csc(\frac{\pi}{6}) = 2, \cot(\frac{\pi}{6}) = \sqrt{3})$$

$$\Rightarrow \int_{\pi/6}^{\pi/4} \csc(x) dx = \left[\frac{\log|\sqrt{2} - 1|}{|2 - \sqrt{3}|} \right]$$

14. Question

Evaluate the following definite integrals:

$$\int_0^1 \frac{1-x}{1+x} dx$$

Answer

$$\int_0^1 \left(\frac{1-x}{1+x} \right) dx = \int_0^1 \left(\frac{1}{x+1} \right) dx - \int_0^1 \left(\frac{x}{x+1} \right) dx$$

$$\Rightarrow \int_0^1 \left(\frac{1-x}{1+x} \right) dx = \int_0^1 \left(\frac{1}{x+1} \right) dx - \int_0^1 \left(\frac{x+1-1}{x+1} \right) dx$$

$$\Rightarrow \int_0^1 \left(\frac{1-x}{1+x} \right) dx = \int_0^1 \left(\frac{1}{x+1} \right) dx - \int_0^1 \left(\frac{x+1}{x+1} \right) dx + \int_0^1 \left(\frac{1}{x+1} \right) dx$$

$$\Rightarrow \int_0^1 \left(\frac{1-x}{1+x} \right) dx = 2 \times \int_0^1 \left(\frac{1}{x+1} \right) dx - \int_0^1 \left(\frac{x+1}{x+1} \right) dx$$

$$\Rightarrow \int_0^1 \left(\frac{1-x}{1+x} \right) dx = 2 \times \int_0^1 \left(\frac{1}{x+1} \right) dx - \int_0^1 1 \times dx$$

Using the formula:

$$\int_a^b \left(\frac{1}{x} \right) dx = [\log|x|]_a^b$$

$$\Rightarrow \int_0^1 \left(\frac{1-x}{1+x} \right) dx = 2 \times [\log|x+1|]_0^1 - [x]_0^1$$

$$\Rightarrow \int_0^1 \left(\frac{1-x}{1+x} \right) dx = 2 \times [\log|1+1| - \log|0+1|] - [1-0]$$

$$\Rightarrow \int_0^1 \left(\frac{1-x}{1+x} \right) dx = 2 \times [\log|2| - \log|1|] - [1]$$

$$\Rightarrow \int_0^1 \left(\frac{1-x}{1+x} \right) dx = 2 \times [\log|2| - 0] - 1$$

$$\Rightarrow \int_0^1 \left(\frac{1-x}{1+x} \right) dx = 2 \times \log|2| - 1$$

15. Question

Evaluate the following definite integrals:

$$\int_0^\pi \frac{1}{1+\sin x} dx$$

Answer

$$\Rightarrow \int_0^\pi \left(\frac{1}{1+\sin x} \right) dx = \int_0^\pi \left(\frac{dx}{1+\sin x} \right) \times \left[\frac{1-\sin x}{1-\sin x} \right]$$

$$\Rightarrow \int_0^\pi \left(\frac{1}{1+\sin x} \right) dx = \int_0^\pi \frac{1-\sin x}{1-\sin^2 x} dx$$

$$\Rightarrow \int_0^\pi \left(\frac{1}{1+\sin x} \right) dx = \int_0^\pi \frac{1-\sin x}{\cos^2 x} dx$$

$$(\because 1 - \sin^2 x = \cos^2 x)$$

$$\Rightarrow \int_0^\pi \left(\frac{1}{1+\sin x} \right) dx = \int_0^\pi \frac{1}{\cos^2 x} dx - \int_0^\pi \frac{\sin x}{\cos^2 x} dx$$

$$\Rightarrow \int_0^\pi \left(\frac{1}{1+\sin x} \right) dx = \int_0^\pi \sec^2 x dx - \int_0^\pi \tan(x) \sec(x) dx$$

Now, we know,

$$\int_a^b \sec^2 x dx = [\tan(x)]_a^b$$

And,

$$\int_a^b \tan(x) \sec(x) dx = [\sec(x)]_a^b$$

$$\therefore \int_0^\pi \left(\frac{1}{1 + \sin x} \right) dx = [\tan(x)]_0^\pi - [\sec(x)]_0^\pi$$

$$\Rightarrow \int_0^\pi \left(\frac{1}{1 + \sin x} \right) dx = [\tan(\pi) - \tan(0)] - [\sec(\pi) - \sec(0)]$$

$$\Rightarrow \int_0^\pi \left(\frac{1}{1 + \sin x} \right) dx = [0 - 0] - [-1 - 1]$$

$$\Rightarrow \int_0^\pi \left(\frac{1}{1 + \sin x} \right) dx = -[-2]$$

$$\Rightarrow \int_0^\pi \left(\frac{1}{1 + \sin x} \right) dx = 2$$

16. Question

Evaluate the following definite integrals:

$$\int_{-\pi/4}^{\pi/4} \frac{1}{1 + \sin x} dx$$

Answer

$$\Rightarrow \int_{-\pi/4}^{\pi/4} \left(\frac{1}{1 + \sin x} \right) dx = \int_{-\pi/4}^{\pi/4} \left(\frac{dx}{1 + \sin x} \right) \times \left[\frac{1 - \sin x}{1 - \sin x} \right]$$

$$\Rightarrow \int_{-\pi/4}^{\pi/4} \left(\frac{1}{1 + \sin x} \right) dx = \int_{-\pi/4}^{\pi/4} \frac{1 - \sin x}{1 - \sin^2 x} dx$$

$$\Rightarrow \int_{-\pi/4}^{\pi/4} \left(\frac{1}{1 + \sin x} \right) dx = \int_{-\pi/4}^{\pi/4} \frac{1 - \sin x}{\cos^2 x} dx$$

$$(\because 1 - \sin^2 x = \cos^2 x)$$

$$\Rightarrow \int_{-\pi/4}^{\pi/4} \left(\frac{1}{1 + \sin x} \right) dx = \int_{-\pi/4}^{\pi/4} \frac{1}{\cos^2 x} dx - \int_{-\pi/4}^{\pi/4} \frac{\sin x}{\cos^2 x} dx$$

$$\Rightarrow \int_{-\pi/4}^{\pi/4} \left(\frac{1}{1 + \sin x} \right) dx = \int_{-\pi/4}^{\pi/4} \sec^2 x dx - \int_{-\pi/4}^{\pi/4} \tan(x) \sec(x) dx$$

Now, we know,

$$\int_a^b \sec^2 x dx = [\tan(x)]_a^b$$

And,

$$\int_a^b \tan(x) \sec(x) dx = [\sec(x)]_a^b$$

$$\therefore \int_{-\pi/4}^{\pi/4} \left(\frac{1}{1 + \sin x} \right) dx = [\tan(x)]_{-\pi/4}^{\pi/4} - [\sec(x)]_{-\pi/4}^{\pi/4}$$

$$\Rightarrow \int_{-\pi/4}^{\pi/4} \left(\frac{1}{1 + \sin x} \right) dx = [\tan(\pi/4) - \tan(-\pi/4)] - [\sec(\pi/4) - \sec(-\pi/4)]$$

$$\Rightarrow \int_{-\pi/4}^{\pi/4} \left(\frac{1}{1 + \sin x} \right) dx = [1 - (-1)] - [-\sqrt{2} - (\sqrt{2})]$$

$$(\because \sec(-\theta) = \sec \theta)$$

$$\Rightarrow \int_{-\pi/4}^{\pi/4} \left(\frac{1}{1 + \sin x} \right) dx = [2] - [-2\sqrt{2}]$$

$$\Rightarrow \int_{-\pi/4}^{\pi/4} \left(\frac{1}{1 + \sin x} \right) dx = 2 + 2\sqrt{2}$$

17. Question

Evaluate the following definite integrals:

$$\int_0^{\pi/2} \cos^2 x dx$$

Answer

$$\text{Let } I = \int_0^{\pi/2} \cos^2 x dx = \int_0^{\pi/2} \frac{1 + \cos(2x)}{2} dx$$

$$\Rightarrow \int_0^{\pi/2} \cos^2 x dx = \int_0^{\pi/2} \frac{1}{2} dx + \int_0^{\pi/2} \frac{\cos(2x)}{2} dx$$

Using the formula:

$$\int_a^b \cos(ax) dx = \left[\frac{\sin(ax)}{a} \right]_a^b$$

$$\Rightarrow \int_0^{\pi/2} \cos^2 x dx = \left(\frac{1}{2} \right) [x]_0^{\pi/2} + \left(\frac{1}{2} \right) \left[\frac{\sin(2x)}{2} \right]_0^{\pi/2}$$

$$\Rightarrow \int_0^{\pi/2} \cos^2 x dx = \left(\frac{1}{2} \right) \left[\frac{\pi}{2} - 0 \right] + \left(\frac{1}{4} \right) \left[\left(\sin 2 \times \frac{\pi}{2} \right) - \left(\sin 2 \times 0 \right) \right]$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} \cos^2 x dx = \frac{\pi}{4} + \left(\frac{1}{4}\right) [\sin \pi - \sin 0]$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} \cos^2 x dx = \frac{\pi}{4} + \left(\frac{1}{4}\right) [0 - 0]$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} \cos^2 x dx = \frac{\pi}{4}$$

18. Question

Evaluate the following definite integrals:

$$\int_0^{\pi/2} \cos^3 x dx$$

Answer

$$\Rightarrow \int_0^{\frac{\pi}{2}} \cos^3 x dx = \int_0^{\frac{\pi}{2}} \cos(x) \times \cos^2(x) dx$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} \cos^3 x dx = \int_0^{\frac{\pi}{2}} \cos(x) \times (1 - \sin^2(x)) dx$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} \cos^3 x dx = \int_0^{\frac{\pi}{2}} \cos(x) dx - \int_0^{\frac{\pi}{2}} \cos(x) \sin^2(x) dx$$

Let $\sin x = t$. Hence, $\cos x dx = dt$, for the second expression.

$$\Rightarrow \int_0^{\frac{\pi}{2}} \cos^3 x dx = [\sin x]_0^{\pi/2} - \int_0^{\frac{\pi}{2}} t^2 dt$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} \cos^3 x dx = [\sin x]_0^{\pi/2} - \left[\frac{t^3}{3}\right]_0^{\pi/2}$$

Put back $t = \sin(x)$

$$\Rightarrow \int_0^{\frac{\pi}{2}} \cos^3 x dx = [\sin \frac{\pi}{2} - \sin 0] - \left(\frac{1}{3}\right) [\sin^3 x]_0^{\pi/2}$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} \cos^3 x dx = [1 - 0] - \left(\frac{1}{3}\right) [\sin^3 \frac{\pi}{2} - \sin^3 0]$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} \cos^3 x dx = [1] - \left(\frac{1}{3}\right) [1^3 - 0^3]$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} \cos^3 x dx = [1] - \left(\frac{1}{3}\right) [1]$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} \cos^3 x dx = [1] - \left(\frac{1}{3}\right)$$

$$\Rightarrow \frac{\int_0^{\frac{\pi}{2}} \cos^3 x dx}{3} = 2$$

19. Question

Evaluate the following definite integrals:

$$\int_0^{\frac{\pi}{6}} \cos x \cos 2x dx$$

Answer

$$\int_0^{\frac{\pi}{6}} \cos x \times \cos(2x) dx = \int_0^{\frac{\pi}{6}} \cos x \times (2 \cos^2 x - 1) dx$$

$$\Rightarrow \int_0^{\frac{\pi}{6}} \cos x \times \cos(2x) dx = \int_0^{\frac{\pi}{6}} (2 \cos^3 x - \cos x) dx$$

$$\Rightarrow \int_0^{\frac{\pi}{6}} \cos x \times \cos(2x) dx = 2 \int_0^{\frac{\pi}{6}} \cos^3 x dx - \int_0^{\frac{\pi}{6}} \cos x dx$$

We know,

$$\Rightarrow \int_0^{\frac{\pi}{6}} \cos^3 x dx = \int_0^{\frac{\pi}{6}} \cos(x) \times \cos^2(x)$$

$$\Rightarrow \int_0^{\frac{\pi}{6}} \cos^3 x dx = \int_0^{\frac{\pi}{6}} \cos(x) \times (1 - \sin^2(x)) dx$$

$$\Rightarrow \int_0^{\frac{\pi}{6}} \cos^3 x dx = \int_0^{\frac{\pi}{6}} \cos(x) dx - \int_0^{\frac{\pi}{6}} \cos(x) \sin^2(x) dx$$

Let $\sin x = t$. Hence, $\cos x dx = dt$. For second expression,

$$\Rightarrow \int_0^{\frac{\pi}{6}} \cos^3 x dx = [\sin x]_0^{\frac{\pi}{6}} - \int_0^{\frac{\pi}{6}} t^2 dt$$

$$\Rightarrow \int_0^{\frac{\pi}{6}} \cos^3 x dx = [\sin x]_0^{\frac{\pi}{6}} - \left[\frac{t^3}{3} \right]_0^{\frac{\pi}{6}}$$

Put $t = \sin(x)$

$$\Rightarrow \int_0^{\frac{\pi}{6}} \cos^3 x dx = [\sin x]_0^{\frac{\pi}{6}} - \left[\frac{\sin^3 x}{3} \right]_0^{\frac{\pi}{6}}$$

$$\Rightarrow \int_0^{\frac{\pi}{6}} \cos^3 x dx = \left[\sin \frac{\pi}{6} - \sin 0 \right] - \left(\frac{1}{3} \right) \left[\sin^3 \frac{\pi}{6} - \sin^3 0 \right]$$

$$\Rightarrow \int_0^{\frac{\pi}{6}} \cos^3 x dx = \left[\frac{1}{2} \right] - \left(\frac{1}{3} \right) \left[\frac{1}{2} \right]^3$$

$$\Rightarrow \int_0^{\frac{\pi}{6}} \cos^3 x dx = \left[\frac{1}{2} \right] - \left[\frac{1}{24} \right]$$

$$\Rightarrow \int_0^{\frac{\pi}{6}} \cos^3 x dx = (12 - 1)/24$$

$$\Rightarrow \int_0^{\frac{\pi}{6}} \cos^3 x dx = (11)/24 \text{ (equation 2)}$$

From equation 2 put value of $\int_0^{\frac{\pi}{6}} \cos^3 x dx$ in equation 1.

$$\Rightarrow \int_0^{\frac{\pi}{6}} \cos x \times \cos(2x) dx = 2 \int_0^{\frac{\pi}{6}} (\cos^3 x) dx - \int_0^{\frac{\pi}{6}} \cos x dx$$

$$\Rightarrow \int_0^{\frac{\pi}{6}} \cos x \times \cos(2x) dx = 2 \times \left(\frac{11}{24} \right) - \int_0^{\frac{\pi}{6}} \cos x dx$$

$$\Rightarrow \int_0^{\frac{\pi}{6}} \cos x \times \cos(2x) dx = 2 \times \left(\frac{11}{24} \right) - [\sin x]_0^{\pi/6}$$

$$\Rightarrow \int_0^{\frac{\pi}{6}} \cos x \times \cos(2x) dx = 2 \times \left(\frac{11}{24} \right) - [\sin \frac{\pi}{6} - \sin 0]$$

$$\Rightarrow \int_0^{\frac{\pi}{6}} \cos x \times \cos(2x) dx = 2 \times \left(\frac{11}{24} \right) - \left[\frac{1}{2} - 0 \right]$$

$$\Rightarrow \int_0^{\frac{\pi}{6}} \cos x \times \cos(2x) dx = \left(\frac{11}{12} \right) - \left[\frac{1}{2} \right]$$

$$\Rightarrow \int_0^{\frac{\pi}{6}} \cos x \times \cos(2x) dx = \left(\frac{5}{12} \right)$$

20. Question

Evaluate the following definite integrals:

$$\int_0^{\pi/2} \sin x \sin 2x dx$$

Answer

$$\int_0^{\frac{\pi}{2}} \sin x \times \sin(2x) dx = \int_0^{\frac{\pi}{2}} \sin x \times 2 \times \sin x \cos x dx$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} \sin x \times \sin(2x) dx = 2 \int_0^{\frac{\pi}{2}} \sin^2 x \cos x dx$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} \sin x \times \sin(2x) dx = 2 \int_0^{\frac{\pi}{2}} (1 - \cos^2 x) \cos x dx$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} \sin x \times \sin(2x) dx = 2 \int_0^{\frac{\pi}{2}} \cos x dx - 2 \int_0^{\frac{\pi}{2}} \cos^3 x dx$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} \sin x \times \sin(2x) dx = 2[\sin x]_0^{\frac{\pi}{2}} - 2 \times \frac{2}{3}$$

First let us find,

$$\int_0^{\frac{\pi}{2}} \cos^3 x dx = \int_0^{\frac{\pi}{2}} \cos(x) \times \cos^2(x)$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} \cos^3 x dx = \int_0^{\pi/2} \cos(x) \times (1 - \sin^2(x)) dx$$

$$\Rightarrow \int_0^{\pi/2} \cos^3 x dx = \int_0^{\frac{\pi}{2}} \cos(x) dx - \int_0^{\frac{\pi}{2}} \cos(x) \sin^2(x) dx$$

Let $\sin x = t$. Hence, $\cos x dx = dt$. For second expression,

$$\Rightarrow \int_0^{\frac{\pi}{2}} \cos^3 x dx = [\sin x]_0^{\pi/2} - \int_0^{\frac{\pi}{2}} t^2 dt$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} \cos^3 x dx = [\sin x]_0^{\pi/2} - \left[\frac{t^3}{3} \right]_0^{\pi/2}$$

Put $t = \sin(x)$

$$\Rightarrow \int_0^{\frac{\pi}{2}} \cos^3 x dx = [\sin x]_0^{\pi/2} - \left[\frac{\sin^3 x}{3} \right]_0^{\pi/2}$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} \cos^3 x dx = \left[\sin \frac{\pi}{2} - \sin 0 \right] - \left(\frac{1}{3} \right) \left[\sin^3 \frac{\pi}{2} - \sin^3 0 \right]$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} \cos^3 x dx = 1 - \left(\frac{1}{3} \right)$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} \cos^3 x dx = \frac{2}{3}$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} \sin x \times \sin(2x) dx = 2 \times [1 - 0] - 2 \times \frac{2}{3}$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} \sin x \times \sin(2x) dx = 2 - \left(\frac{4}{3} \right)$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} \sin x \times \sin(2x) dx = \left(\frac{2}{3}\right)$$

21. Question

Evaluate the following definite integrals:

$$\int_{\pi/3}^{\pi/4} (\tan x + \cot x)^2 dx$$

Answer

$$\int_{\pi/3}^{\pi/4} (\tan x + \cot x)^2 dx = \int_{\pi/3}^{\pi/4} (\tan^2 x + \cot^2 x + 2 \times \tan x \times \cot x) dx$$

We know, $\tan x \times \cot x = 1$

$$\int_{\pi/3}^{\pi/4} (\tan x + \cot x)^2 dx = \int_{\pi/3}^{\pi/4} (\tan^2 x + \cot^2 x + 2) dx$$

We know, $\tan^2 x = \sec^2 x - 1$ and $\cot^2 x = \operatorname{cosec}^2 x - 1$

$$\Rightarrow \int_{\pi/3}^{\pi/4} (\tan x + \cot x)^2 dx = \int_{\pi/3}^{\pi/4} (\sec^2 x - 1 + \operatorname{cosec}^2 x - 1 + 2) dx$$

$$\Rightarrow \int_{\pi/3}^{\pi/4} (\tan x + \cot x)^2 dx = \int_{\pi/3}^{\pi/4} (\sec^2 x + \operatorname{cosec}^2 x) dx$$

$$\Rightarrow \int_{\pi/3}^{\pi/4} (\tan x + \cot x)^2 dx = \int_{\pi/3}^{\pi/4} (\sec^2 x) dx + \int_{\pi/3}^{\pi/4} (\operatorname{cosec}^2 x) dx$$

We know integration of $\sec^2 x$ is $\tan x$ and of $\operatorname{cosec}^2 x$ is $-\cot x$. Therefore,

$$\Rightarrow \int_{\pi/3}^{\pi/4} (\tan x + \cot x)^2 dx = [\tan x]_{\pi/3}^{\pi/4} + -[\cot x]_{\pi/3}^{\pi/4}$$

$$\Rightarrow \int_{\pi/3}^{\pi/4} (\tan x + \cot x)^2 dx = \left[\tan \frac{\pi}{4} - \tan \frac{\pi}{3} \right] - \left[\cot \frac{\pi}{4} - \cot \frac{\pi}{3} \right]$$

$$\Rightarrow \int_{\pi/3}^{\pi/4} (\tan x + \cot x)^2 dx = [1 - \sqrt{3}] - [1 - 1/\sqrt{3}]$$

$$\Rightarrow \int_{\pi/3}^{\pi/4} (\tan x + \cot x)^2 dx = \frac{1}{\sqrt{3}} - \sqrt{3}$$

22. Question

Evaluate the following definite integrals:

$$\int_0^{\pi/2} \cos^4 x dx$$

Answer

$$\int_0^{\frac{\pi}{2}} \cos^4 x dx = \int_0^{\frac{\pi}{2}} \cos^2 x \times \cos^2 x dx$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} \cos^4 x dx = \int_0^{\frac{\pi}{2}} \left(\frac{1 + \cos 2x}{2} \right) \times \left(\frac{1 + \cos 2x}{2} \right) dx$$

$$(\because 1 + \cos 2\theta = 2 \cos^2 \theta)$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} \cos^4 x dx = \left(\frac{1}{4} \right) \int_0^{\frac{\pi}{2}} (1 + \cos 2x)(1 + \cos 2x) dx$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} \cos^4 x dx = \left(\frac{1}{4} \right) \int_0^{\frac{\pi}{2}} (1 + \cos^2 2x + 2\cos 2x) dx$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} \cos^4 x dx = \left(\frac{1}{4} \right) \int_0^{\frac{\pi}{2}} \left(1 + \frac{(1 + \cos 4x)}{2} + 2\cos 2x \right) dx$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} \cos^4 x dx = \left(\frac{1}{8} \right) \int_0^{\frac{\pi}{2}} (2 + (1 + \cos 4x) + 4\cos 2x) dx$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} \cos^4 x dx = \left(\frac{1}{8} \right) \left[\int_0^{\frac{\pi}{2}} (2dx) + \int_0^{\frac{\pi}{2}} (1 + \cos 4x) dx + \int_0^{\frac{\pi}{2}} (4\cos 2x) dx \right]$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} \cos^4 x dx = \left(\frac{1}{8} \right) \left[[2x]_0^{\frac{\pi}{2}} + [x]_0^{\frac{\pi}{2}} + [(sin 4x)/4]_0^{\frac{\pi}{2}} + 4[(sin 2x)/2]_0^{\frac{\pi}{2}} \right]$$

$$\int_0^{\frac{\pi}{2}} \cos^4 x dx = \left(\frac{1}{8} \right) \left[\left[2 \times \frac{\pi}{2} - 0 \right] + \left[\frac{\pi}{2} - 0 \right] + \left(\frac{1}{4} \right) [\sin 4 \times \frac{\pi}{2} - \sin 0] + (4/2)[\sin 2 \times \frac{\pi}{2} - \sin 0] \right]$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} \cos^4 x dx = \left(\frac{1}{8} \right) \left[[\pi] + \left[\frac{\pi}{2} \right] + \left(\frac{1}{4} \right) [\sin 2 \times \pi] + (2)[\sin \pi] \right]$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} \cos^4 x dx = \left(\frac{1}{8} \right) \left[[\pi] + \left[\frac{\pi}{2} \right] + 0 + (2)[0] \right]$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} \cos^4 x dx = \left(\frac{1}{8} \right) \left[[\pi] + \left[\frac{\pi}{2} \right] \right]$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} \cos^4 x dx = \left[\frac{3\pi}{16} \right]$$

23. Question

Evaluate the following definite integrals:

$$\int_0^{\pi/2} (a^2 \cos^2 x + b^2 \sin^2 x) dx$$

Answer

$$\int_0^{\frac{\pi}{2}} (a^2 \cos^2 x + b^2 \sin^2 x) dx = \int_0^{\frac{\pi}{2}} [a^2 \cos^2 x + b^2(1 - \cos^2 x)] dx$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} (a^2 \cos^2 x + b^2 \sin^2 x) dx = \int_0^{\frac{\pi}{2}} [a^2 \cos^2 x + b^2(1 - b^2 \cos^2 x)] dx$$

$$\int_0^{\frac{\pi}{2}} (a^2 \cos^2 x + b^2 \sin^2 x) dx = \int_0^{\frac{\pi}{2}} [a^2 \cos^2 x] dx + b^2 \int_0^{\frac{\pi}{2}} 1 \times dx - b^2 \int_0^{\frac{\pi}{2}} [\cos^2 x] dx$$

$$\int_0^{\frac{\pi}{2}} (a^2 \cos^2 x + b^2 \sin^2 x) dx = \frac{a^2}{2} [1 + \cos 2x]_0^{\pi/2} + b^2 [x]_0^{\pi/2} - \frac{b^2}{2} [1 + \cos 2x]_0^{\frac{\pi}{2}}$$

$$\begin{aligned} \int_0^{\frac{\pi}{2}} (a^2 \cos^2 x + b^2 \sin^2 x) dx &= \frac{a^2}{2} \left[\left(1 + \cos 2 \times \frac{\pi}{2} \right) - (1 - \cos 0) \right] + b^2 \left[\frac{\pi}{2} - 0 \right] \\ &\quad - \frac{b^2}{2} \left[\left(1 + \cos 2 \times \frac{\pi}{2} \right) - (1 - \cos 0) \right] \end{aligned}$$

$$\begin{aligned} \Rightarrow \int_0^{\frac{\pi}{2}} (a^2 \cos^2 x + b^2 \sin^2 x) dx &= \frac{a^2}{2} [(1 + -1) - (1 - 1)] + b^2 \left[\frac{\pi}{2} \right] - \\ &\quad \frac{b^2}{2} [(1 + -1) - (1 - 1)] \end{aligned}$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} (a^2 \cos^2 x + b^2 \sin^2 x) dx = b^2 \left[\frac{\pi}{2} \right]$$

24. Question

Evaluate the following definite integrals:

$$\int_0^{\pi/2} \sqrt{1 + \sin x} dx$$

Answer

$$\int_0^{\frac{\pi}{2}} (\sqrt{1 + \sin x}) dx = \int_0^{\frac{\pi}{2}} (\sqrt{1 + \sin x}) \times \frac{\sqrt{1 - \sin x}}{\sqrt{1 - \sin x}} dx$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} (\sqrt{1 + \sin x}) dx = \int_0^{\frac{\pi}{2}} \left[\frac{(\sqrt{1 - \sin^2 x})}{\sqrt{1 - \sin x}} \right] dx$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} (\sqrt{1 + \sin x}) dx = \int_0^{\frac{\pi}{2}} \left[\frac{\sqrt{\cos^2 x}}{\sqrt{1 - \sin x}} \right] dx$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} (\sqrt{1 + \sin x}) dx = \int_0^{\frac{\pi}{2}} \left[\frac{\cos x}{\sqrt{1 - \sin x}} \right] dx$$

Let $1 - \sin x = t^2$. Hence, $-\cos x dx = 2t dt$ and $\cos x dx = -2t dt$.

$$\Rightarrow \int_0^{\frac{\pi}{2}} (\sqrt{1 + \sin x}) dx = \int_0^{\frac{\pi}{2}} \left[\frac{-2t}{\sqrt{t^2}} \right] dt$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} (\sqrt{1 + \sin x}) dx = \int_0^{\frac{\pi}{2}} \left[\frac{-2t}{t} \right] dt$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} (\sqrt{1 + \sin x}) dx = (-2) \int_0^{\frac{\pi}{2}} [1] dt$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} (\sqrt{1 + \sin x}) dx = (-2) [t]_0^{\frac{\pi}{2}}$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} (\sqrt{1 + \sin x}) dx = (-2) [\sqrt{1 - \sin x}]_0^{\frac{\pi}{2}}$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} (\sqrt{1 + \sin x}) dx = (-2) [\sqrt{1 - \sin \left(\frac{\pi}{2} \right)} - \sqrt{1 - \sin 0}]$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} (\sqrt{1 + \sin x}) dx = (-2) [\sqrt{1 - 1} - \sqrt{1 - 0}]$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} (\sqrt{1 + \sin x}) dx = (-2) [-1]$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} (\sqrt{1 + \sin x}) dx = 2$$

25. Question

Evaluate the following definite integrals:

$$\int_0^{\pi/2} \sqrt{1 + \cos x} dx$$

Answer

$$\Rightarrow \int_0^{\frac{\pi}{2}} (\sqrt{1 + \cos x}) dx = \int_0^{\frac{\pi}{2}} (\sqrt{1 + \cos x}) \times \frac{\sqrt{1 - \cos x}}{\sqrt{1 - \cos x}} dx$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} (\sqrt{1 + \cos x}) dx = \int_0^{\frac{\pi}{2}} \left[\frac{(\sqrt{1 - \cos^2 x})}{\sqrt{1 - \cos x}} \right] dx$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} (\sqrt{1 + \cos x}) dx = \int_0^{\frac{\pi}{2}} \left[\frac{\sqrt{\sin^2 x}}{\sqrt{1 - \cos x}} \right] dx$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} (\sqrt{1 + \cos x}) dx = \int_0^{\frac{\pi}{2}} \left[\frac{\sin x}{\sqrt{1 - \cos x}} \right] dx$$

Let $1 - \cos x = t^2$ hence $\sin x dx = 2 t dt$

$$\Rightarrow \int_0^{\frac{\pi}{2}} (\sqrt{1 + \cos x}) dx = \int_0^{\frac{\pi}{2}} \left[\frac{2t}{\sqrt{t^2}} \right] dt$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} (\sqrt{1 + \cos x}) dx = \int_0^{\frac{\pi}{2}} \left[\frac{2t}{t} \right] dt$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} (\sqrt{1 + \cos x}) dx = (2) \int_0^{\frac{\pi}{2}} [1] dt$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} (\sqrt{1 + \cos x}) dx = (2) [t]_0^{\pi/2}$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} (\sqrt{1 + \cos x}) dx = (2) [\sqrt{1 - \cos x}]_0^{\pi/2}$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} (\sqrt{1 + \cos x}) dx = (2) [\sqrt{1 - \cos(\frac{\pi}{2})} - \sqrt{1 - \cos 0}]$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} (\sqrt{1 + \cos x}) dx = (2) [\sqrt{1 - 0} - \sqrt{1 - 1}]$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} (\sqrt{1 + \cos x}) dx = (2)[1]$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} (\sqrt{1 + \cos x}) dx = 2$$

26. Question

Evaluate the following definite Integrals:

$$\int_0^{\pi/2} x^2 \sin x \, dx$$

Answer

We are asked to calculate $\int_0^{\frac{\pi}{2}} x \sin x \, dx$

For this we have to apply integration by parts

Let u and v be two functions then

$$\int u \, dv = uv - \int v \, du$$

To choose the first function u we use "ILATE" rule

That is

I=inverse trigonometric function

L=logarithmic function

A=algebraic function

T=trigonometric functions

E=exponential function

So in this preference, the first function is chosen to make the integration simpler.

Now, In the given question x is an algebraic function and it is chosen as u (A comes first in "ILATE" rule)

So first let us integrate the equation and then let us substitute the limits in it

$$\int x \sin x \, dx = x \int \sin x \, dx - \int \left(x \int \sin x \, dx \right) \, dx$$

$$= -x \cos x + \int \cos x \, dx$$

Therefore, now substitute the limits given:

Note that $\int \sin x \, dx = -\cos x$ and $\int \cos x \, dx = \sin x$

$$\int_0^{\pi/2} x \sin x \, dx = \left[(-x \cos x + \int \cos x \, dx) \right]_0^{\pi/2}$$

$$= [(-x \cos x + \sin x)]_0^{\pi/2}$$

First we have to substitute the upper limit and then subtract the second limit value from it

$$= -\left(\frac{\pi}{2}\right) \cos\left(\frac{\pi}{2}\right) + \sin\left(\frac{\pi}{2}\right) - (0 \cos 0 + \sin 0)$$

Note that $\sin 0 = 0$ and $\cos 0 = 1$

$$= 0 + 1 - 0$$

$$= 1$$

27. Question

Evaluate the following definite Integrals:

$$\int_0^{\pi/2} x \cos x \, dx$$

Answer

We are asked to calculate $\int_0^{\frac{\pi}{2}} x \cos x \, dx$

For this we have to apply integration by parts

Let u and v be two functions then

$$\int u \, dv = uv - \int v \, du$$

To choose the first function u we use "ILATE" rule

That is

I=inverse trigonometric function

L=logarithmic function

A=algebraic function

T=trigonometric functions

E=exponential function

So in this preference, the first function is chosen to make the integration simpler.

Now, In the given question x is an algebraic function and it is chosen as u(A comes first in "ILATE" rule)

So first let us integrate the equation and then let us substitute the limits in it

$$\begin{aligned}\int x \cos x \, dx &= x \int \cos x \, dx - \int \left(x \int \cos x \, dx \right) \, dx \\ &= x \sin x - \int \sin x \, dx\end{aligned}$$

Therefore, now substitute the limits given:

Note that $\int \sin x \, dx = -\cos x$ and $\int \cos x \, dx = \sin x$

$$\int_0^{\frac{\pi}{2}} x \cos x \, dx = \left[x \sin x - \int \sin x \, dx \right]_0^{\frac{\pi}{2}}$$

$$\int_0^{\frac{\pi}{2}} x \cos x \, dx = \left[x \sin x + \cos x \right]_0^{\frac{\pi}{2}}$$

First we have to substitute the upper limit and then subtract the second limit value from it

$$\begin{aligned}&= \left[\frac{\pi}{2} \right] \sin \frac{\pi}{2} + \cos \frac{\pi}{2} - [0 \sin 0 + \cos 0] \\ &= \left[\frac{\pi}{2} \right] 1 + 0 - [0 + 1] \\ &= \frac{\pi - 2}{2}\end{aligned}$$

28. Question

Evaluate the following definite Integrals:

$$\int_0^{\pi/2} x^2 \cos x \, dx$$

Answer

For this, we have to apply integration by parts

Let u and v be two functions then

$$\int u \, dv = uv - \int v \, du$$

To choose the first function u we use "ILATE" rule

That is

I=inverse trigonometric function

L=logarithmic function

A=algebraic function

T=trigonometric functions

E=exponential function

So in this preference,, the first function is chosen to make the integration simpler.

Now, In the given question x^2 is an algebraic function and it is chosen as u (A comes first in "ILATE" rule)

So first let us integrate the equation and then let us substitute the limits in it.

$$\int x^2 \cos x \, dx = x^2 \int \cos x \, dx - \int \left(x^2 \int \cos x \, dx \right) \, dx$$

$$= x^2 \sin x - \int 2x \sin x \, dx$$

$$= x^2 \sin x - 2 \left[x \int \sin x \, dx - \int \left(\int \sin x \, dx \right) \, dx \right]$$

$$= x^2 \sin x - 2 \left[-x \cos x - \int \cos x \, dx \right]$$

So now we have to substitute the limits in this equation.

And should subtract upper limit value from lower limit value

$$\int_0^{\pi/2} x^2 \cos x \, dx = [x^2 \sin x - 2 \left[-x \cos x - \int \cos x \, dx \right]]_0^{\pi/2}$$

$$\sin \frac{\pi}{2} = 1, \cos \frac{\pi}{2} = 0, \sin 0 = 0, \cos 0 = 1.$$

$$= [x^2 \sin x - 2 \left[-x \cos x + \sin x \right]]_0^{\pi/2}$$

$$= \frac{\pi^2}{4} \sin \frac{\pi}{2} - 2 \left[-\frac{\pi}{2} \cos \frac{\pi}{2} + \sin \frac{\pi}{2} \right] - \{0 \times \sin 0 - 2[-0 \cos 0 + \sin 0]\}$$

$$= \frac{\pi^2}{4} \times 1 - 2[0 + 1] - \{0 - 2[-0 \times 1 + 0]\}$$

$$= \frac{\pi^2}{4} - 2.$$

29. Question

Evaluate the following definite Integrals:

$$\int_0^{\pi/4} x^2 \sin x \, dx$$

Answer

For this, we have to apply integration by parts

Let u and v be two functions then

$$\int u \, dv = uv - \int v \, du$$

To choose the first function u we use "ILATE" rule

That is

I=inverse trigonometric function

L=logarithmic function

A=algebraic function

T=trigonometric functions

E=exponential function

So in this preference,, the first function is chosen to make the integration simpler.

Now, In the given question x^2 is an algebraic function and it is chosen as u (A comes first in "ILATE" rule)

So first let us integrate the equation and then let us substitute the limits in it.

$$\begin{aligned} \int x^2 \sin x \, dx &= x^2 \int \sin x \, dx - \int 2x \left(\int \sin x \, dx \right) \, dx \\ &= -x^2 \cos x + 2 \left[x \int \cos x \, dx - \int \int \cos x \, dx \, dx \right] \\ &= -x^2 \cos x + 2 \left[x \sin x - \int \sin x \, dx \right] \end{aligned}$$

So now we have to substitute the limits in this equation.

And should subtract upper limit value from lower limit value

$$\int_0^{\pi/4} x^2 \sin x \, dx = \left[-x^2 \cos x + 2 \left[x \sin x - \int \sin x \, dx \right] \right]_0^{\pi/4}$$

$$\sin \frac{\pi}{2} = 1, \cos \frac{\pi}{2} = 0, \sin 0 = 0, \cos 0 = 1$$

$$\cos \frac{\pi}{4} = \sin \frac{\pi}{4} = \frac{1}{\sqrt{2}}$$

$$\begin{aligned} &= \left[-x^2 \cos x + 2 \left[x \sin x + \cos x \right] \right]_0^{\pi/4} \\ &= \left[\frac{-\pi^2}{16} \times \cos \frac{\pi}{4} + 2 \left[\frac{\pi}{4} \sin \frac{\pi}{4} + \cos \frac{\pi}{4} - \{-0 \times \cos 0 + [0 \times \sin 0 + 2 \cos 0]\} \right] \right] \\ &= \left[\frac{-\pi^2}{16} \times \frac{1}{\sqrt{2}} + 2 \left[\frac{\pi}{4} \times \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}} \right] - \{0 + 2[0 + 1]\} \right] \\ &= \frac{1}{\sqrt{2}} \left[\frac{-\pi^2}{16} + \frac{\pi}{2} + 2 \right] - 2 \\ &= \sqrt{2} + \frac{\pi}{2\sqrt{2}} - \frac{\pi^2}{16\sqrt{2}} - 2 \end{aligned}$$

30. Question

Evaluate the following definite Integrals:

$$\int_0^{\pi/2} x^2 \cos 2x \, dx$$

Answer

For this we have to apply integration by parts

Let u and v be two functions then

$$\int u \, dv = uv - \int v \, du$$

To choose the first function u we use "ILATE" rule

That is

I=inverse trigonometric function

L=logarithmic function

A=algebraic function

T=trigonometric functions

E=exponential function

So in this preference, the first function is chosen to make the integration simpler.

Now, In the given question x^2 is an algebraic function and it is chosen as u(A comes first in "ILATE" rule)

So first let us integrate the equation and then let us substitute the limits in it.

Note that $\int \sin x = -\cos x$ and $\int \cos x = \sin x$

$$\begin{aligned}\int x^2 \cos 2x \, dx &= x^2 \int \cos 2x \, dx - \int \left(x^2 \int \cos 2x \, dx \right) \, dx \\&= \frac{x^2 \sin 2x}{2} - \int 2x \times \frac{\sin 2x}{2} \, dx \\&= \frac{x^2 \sin 2x}{2} - [x \int \sin 2x \, dx - \int \left(\int \sin 2x \, dx \right) \, dx] \\&= \frac{x^2 \sin 2x}{2} + \left[\frac{x \cos 2x}{2} - \int \frac{\cos 2x}{2} \, dx \right] \\&= \frac{x^2 \sin 2x}{2} + \left[\frac{x \cos 2x}{2} - \frac{\sin 2x}{4} \right] \\ \int_0^{\pi/2} x^2 \cos 2x \, dx &= \left[\frac{x^2 \sin 2x}{2} + \left[\frac{x \cos 2x}{2} - \frac{\sin 2x}{4} \right] \right]_0^{\pi/2} \\&= \frac{\pi^2}{4} \times \sin \left(2 \times \frac{\pi}{2} \right) \times \frac{1}{2} + \left[\frac{\pi}{2} \times \cos \left(2 \times \frac{\pi}{2} \right) \times \frac{1}{2} - \sin \left(2 \times \frac{\pi}{2} \right) \times \frac{1}{4} \right] \\&\quad - \left\{ \frac{\pi^2}{4} \times \sin(2 \times 0) \times \frac{1}{2} + \left[\frac{\pi}{2} \times \cos(2 \times 0) \times \frac{1}{2} - \sin(2 \times 0) \times \frac{1}{4} \right] \right\} \\&= \frac{\pi^2}{8} \times 0 - 1 \times \frac{1}{4} \times 0 - \left\{ 0 + \left[\frac{\pi}{4} \right] \right\} \\&= -\frac{\pi}{4}\end{aligned}$$

31. Question

Evaluate the following definite Integrals:

$$\int_0^{\pi/2} x^2 \cos^2 x \, dx$$

Answer

For this we have to apply integration by parts

Let u and v be two functions then

$$\int u \, dv = uv - \int v \, du$$

To choose the first function u we use "ILATE" rule

That is

I=inverse trigonometric function

L=logarithmic function

A=algebraic function

T=trigonometric functions

E=exponential function

So in this preference, the first function is chosen to make the integration simpler.

Now, In the given question x^2 is an algebraic function and it is chosen as u(A comes first in "ILATE" rule)

So first let us integrate the equation and then let us substitute the limits in it.

Let us recall a formula $\cos 2x = 2\cos^2 x - 1$

Now substitute it

$$\int x^2 \cos^2 x \, dx = \int x^2 \left(1 + \frac{\cos 2x}{2}\right) dx$$

$$= \frac{1}{2} \int (x^2 + x^2 \cos 2x) dx$$

$$= \frac{1}{2} \int (x^2 dx + \int x^2 \cos 2x dx)$$

Now let us recall other formula i.e. $\int x^n = \frac{x^{n+1}}{n+1}$

$$\int \sin x = -\cos x \text{ and } \int \cos x = \sin x$$

Using them we can write the equation as

$$\int x^2 dx = \left[\frac{x^3}{3} \right]_0^{\frac{\pi}{2}} = \frac{\pi^2}{24}$$

$$\int x^2 \cos 2x \, dx = x^2 \int \cos 2x dx - \int \int \cos 2x \times \left(\frac{d^2 x}{dx} \right) dx$$

$$= \frac{x^2 \sin 2x}{2} = \int 2x \times \frac{\sin 2x}{2} dx$$

$$= \frac{x^2 \sin 2x}{2} - [x \int \sin 2x \, dx - \int \int \sin 2x \, dx]$$

$$= \frac{x^2 \sin 2x}{2} + \left[\frac{x \cos 2x}{2} - \int \frac{\cos 2x}{2} \right]$$

$$\begin{aligned}
&= \frac{x^2 \sin 2x}{2} + \left[\frac{x \cos 2x}{2} - \frac{\sin 2x}{4} \right]_0^{\pi/2} \\
\int_0^{\pi/2} x^2 \cos 2x \, dx &= \left[\frac{x^2 \sin 2x}{2} + \left[\frac{x \cos 2x}{2} - \frac{\sin 2x}{4} \right] \right]_0^{\pi/2} \\
&= \frac{\pi^2}{4} \times \sin\left(2 \times \frac{\pi}{2}\right) \times \frac{1}{2} + \left[\frac{\pi}{2} \times \cos\left(2 \times \frac{\pi}{2}\right) \times \frac{1}{2} - \sin\left(2 \times \frac{\pi}{2}\right) \times \frac{1}{4} \right] \\
&\quad - \left\{ \frac{\pi^2}{4} \times \sin(2 \times 0) \times \frac{1}{2} + \left[\frac{\pi}{2} \times \cos(2 \times 0) \times \frac{1}{2} - \sin(2 \times 0) \times \frac{1}{4} \right] \right\} \\
&= \frac{\pi^2}{8} \times 0 - 1 \times \frac{1}{4} \times 0 - \left\{ 0 + \left[\frac{\pi}{4} \right] \right\} \\
&= \frac{\pi}{4}
\end{aligned}$$

On substituting these values we get

$$\begin{aligned}
\int_0^{\pi/2} x^2 \cos^2 x \, dx &= \frac{1}{2} \left[\frac{\pi^3}{24} - \frac{\pi}{4} \right] \\
&= \frac{\pi^3}{48} - \frac{\pi}{8}
\end{aligned}$$

32. Question

Evaluate the following definite Integrals:

$$\int_1^2 \log x \, dx$$

Answer

For this we have to apply integration by parts

Let u and v be two functions then

$$\int u \, dv = uv - \int v \, du$$

To choose the first function u we use "ILATE" rule

That is

I=inverse trigonometric function

L=logarithmic function

A=algebraic function

T=trigonometric functions

E=exponential function

So in this preference, the first function is chosen to make the integration simpler.

Now, In the given question 1 is an algebraic function and it is chosen as u(A comes first in "ILATE" rule)

So first let us integrate the equation and then let us substitute the limits in it.

$$\int \log x \, dx = \int 1 \times \log x \, dx = \log x \int x \, dx - \int \frac{1}{x} \left(\int x \, dx \right) \, dx$$

Let us recall that derivative of $\log x$ is $1/x$

$$= x \log x - \int x \times \frac{1}{x} \, dx$$

$$= x \log x - \int dx$$

$$= x \log x - x$$

Now let us substitute the limits

$$\int_1^2 \log x \, dx = [x \log x - x]_1^2$$

$$= 2 \log 2 - 2 - [1 \log 1 - 1]$$

$$= 2 \log 2 - 1$$

33. Question

Evaluate the following definite Integrals:

$$\int_1^3 \frac{\log x}{(x+1)^2} \, dx$$

Answer

For this we have to apply integration by parts

Let u and v be two functions then

$$\int u \, dv = uv - \int v \, du$$

To choose the first function u we use "ILATE" rule

That is

I=inverse trigonometric function

L=logarithmic function

A=algebraic function

T=trigonometric functions

E=exponential function

So in this preference, the first function is chosen to make the integration simpler.

$$\begin{aligned} \int \frac{\log x \, dx}{(x+1)^2} &= \int \frac{1}{(x+1)^2} \times \log x \, dx \\ &= \log x \times \int \frac{1}{(x+1)^2} \, dx - \int \left(\int \frac{1}{(x+1)^2} \, dx \right) \frac{1}{x} \, dx \\ &= -\frac{\log x}{x+1} + \int \frac{1}{x(x+1)} \, dx \\ &= -\frac{\log x}{x+1} + \int \frac{(x+1-x)}{x(x+1)} \, dx \\ &= -\frac{\log x}{x+1} + \int \left(\frac{1}{x} - \frac{1}{x+1} \right) dx \end{aligned}$$

Now we will substitute the limits

$$\int_1^3 \frac{\log x \, dx}{(x+1)^2} = \left[-\frac{\log x}{x+1} + \int \left(\frac{1}{x} - \frac{1}{x+1} \right) dx \right]_1^3$$

$$\int_1^3 \frac{\log x \, dx}{(x+1)^2} = \left[-\frac{\log x}{x+1} + \log x - \log(x+1) \right]_1^3$$

$$= -\frac{\log 3}{3+1} + \log 3 - \log(3+1) - \left\{ -\frac{\log 1}{1+1} + \log 1 - \log(1+1) \right\}$$

$\log 1 = 0$ and $\log 4 = 2\log 2$

$$= \frac{3\log 3}{4} - \log 2$$

34. Question

Evaluate the following definite Integrals:

$$\int_1^e \frac{e^x}{x} (1 + x \log x) dx$$

Answer

let us assume that the given equation is L

$$L = \int_1^e \frac{e^x}{x} (1 + x \log x) dx$$

$$l = \int_1^e \frac{e^x}{x} dx + \int_1^e e^x \log x dx$$

For this we have to apply integration by parts

Let u and v be two functions then

$$\int u dv = uv - \int v du$$

To choose the first function u we use "ILATE" rule

That is

I=inverse trigonometric function

L=logarithmic function

A=algebraic function

T=trigonometric functions

E=exponential function

So in this preference, the first function is chosen to make the integration simpler.

$$= [e^x \log x]_1^e - \int_1^e e^x \log x dx + \int_1^e e^x \log x dx$$

$$= [e^x \log x]_1^e$$

$$= e^e \log e - e \log 1$$

We know that $\log e = 1$

$$= e^e - 0$$

$$= e^e$$

35. Question

Evaluate the following definite Integrals:

$$\int_1^e \frac{\log x}{x} dx$$



Answer

Here in this question by observation we can notice that the derivative of $\log x$ is $1/x$ and the function integral is like

$$\int f(x) d(f(x)) dx = f(x) 2/2$$

Here to solve these kinds of question let us assume $\log x = t$

$$\text{Now } \frac{1}{x} dx = dt$$

Now let us change the limits

$$x=1 \text{ then } t=0$$

$$x=e \text{ then } t=1$$

$$\int_1^e \frac{\log x}{x} dx = \int_0^1 t dt$$

$$= [\frac{t^2}{2}]_0^1$$

$$= \frac{1}{2} - 0$$

$$= \frac{1}{2}$$

36. Question

Evaluate the following definite Integrals:

$$\int_e^2 \left\{ \frac{1}{\log x} - \frac{1}{(\log x)^2} \right\} dx$$

Answer

For this we have to apply integration by parts

Let u and v be two functions then

$$\int u dv = uv - \int v du$$

To choose the first function u we use "ILATE" rule

That is

I=inverse trigonometric function

L=logarithmic function

A=algebraic function

T=trigonometric functions

E=exponential function

So in this preference, the first function is chosen to make the integration simpler.

$$\int \frac{1}{\log x} dx - \frac{1}{\log x} \int dx - \int dx \frac{d}{dx} (\frac{1}{\log x}) dx$$

$$= \frac{x}{\log x} + \int \frac{1 dx}{(\log x)^2}$$

Now let us substitute in the given question equation

$$\begin{aligned} \int_e^{e^2} \frac{1}{\log x} - \frac{1}{(\log x)^2} dx &= \left[\frac{x}{\log x} \right]_e^{e^2} + \int_e^{e^2} \frac{1 dx}{(\log x)^2} - \int_e^{e^2} \frac{1 dx}{(\log x)^2} \\ &= \left[\frac{x}{\log x} \right]_e^{e^2} \\ &= \frac{e^2}{\log e^2} - \frac{e}{\log e} \\ &= \frac{e^2}{2} - e \end{aligned}$$

37. Question

Evaluate the following definite Integrals:

$$\int_1^2 \frac{x+3}{x(x+2)} dx$$

Answer

$$\begin{aligned} &= \int_1^2 \frac{x}{x(x+2)} dx + \int_1^2 \frac{3}{x(x+2)} dx \\ &= \int_1^2 \frac{1}{(x+2)} dx + \int_1^2 \frac{3}{x(x+2)} dx \end{aligned}$$

Here we are solving the equation, recall $\frac{1}{x+2}$ is the derivative of $\log(x+2)$ and splitting the second one

$$\begin{aligned} &= [\log(x+2)]_1^2 + \frac{3}{2} \int \left(\frac{1}{x} - \frac{1}{x+2} \right) dx \\ &= [\log(x+2)]_1^2 - \left[\frac{3}{2} \log(x) - \frac{3}{2} \log(x+2) \right]_1^2 \\ &= \left[\frac{3}{2} \log(x) - \frac{1}{2} \log(x+2) \right]_1^2 \\ &= \frac{3}{2} \log(2) - \frac{1}{2} \log(2+2) - \left\{ \frac{3}{2} \log(1) - \frac{1}{2} \log(1+2) \right\} \\ &= \frac{1}{2} [3\log 2 - \log 4 + \log 3] \end{aligned}$$

Note that $\log 4 = 2\log 2$ and $\log 1 = 0$

$$\begin{aligned} &= \frac{1}{2} [3\log 2 - 2\log 2 + \log 3] \\ &= \frac{1}{2} [\log 2 + \log 3] \\ &= \frac{1}{2} [\log 6] \end{aligned}$$

38. Question

Evaluate the following definite Integrals:

$$\int_0^1 \frac{2x+3}{5x^2+1} dx$$

Answer

If the equation is in this form then convert the numerator as sum of derivative of denominator and some constant

Here we know that denominator derivative is $10x$

So to get it in the numerator multiply and divide by 5

Now you get the equation as

$$\begin{aligned}\int_0^1 \frac{2x + 3}{5x^2 + 1} dx &= \int_0^1 \frac{5(2x + 3)}{5x^2 + 1} dx \times \frac{1}{5} \\ &= \int_0^1 \frac{10x + 15}{5x^2 + 1} dx \times \frac{1}{5} \\ &= \int_0^1 \frac{10x}{5x^2 + 1} dx \times \frac{1}{5} + \int_0^1 \frac{15}{5x^2 + 1} dx \times \frac{1}{5} \\ &= \int_0^1 \frac{10x}{5x^2 + 1} dx \times \frac{1}{5} + \int_0^1 \frac{3}{5x^2 + 1} dx \\ &= \int_0^1 \frac{10x}{5x^2 + 1} dx \times \frac{1}{5} + \int_0^1 \frac{3}{5(x^2 + \frac{1}{5})} dx\end{aligned}$$

We already know that derivative of $\log x$ is $1/x$

Using that here derivative of $\log(5x^2 + 1)$ = $\frac{10x}{5x^2 + 1}$

And derivative of $\tan^{-1} \frac{x}{a} = \frac{1}{a^2 + x^2}$

$$\begin{aligned}\text{So } &= \frac{1}{5} \log(5x^2 + 1) + \frac{3}{5} \times \frac{1}{\sqrt{5}} \times \tan^{-1} \frac{x}{\sqrt{5}} \\ &= \frac{1}{5} \log(5x^2 + 1) + \frac{3}{\sqrt{5}} \times \tan^{-1} \sqrt{5}x\end{aligned}$$

Now substitute limits 1 and 0

$$\begin{aligned}&= \frac{1}{5} \log(5 \times 1 + 1) + \frac{3}{\sqrt{5}} \times \tan^{-1} \sqrt{5 \times 1} - \left\{ \frac{1}{5} \log(5 \times 0 + 1) \right. \\ &\quad \left. + \frac{3}{\sqrt{5}} \times \tan^{-1} \sqrt{5 \times 0} \right\} \\ &= \frac{1}{5} \log(6) - \frac{3}{\sqrt{5}} \times \tan^{-1} \sqrt{5} - \left\{ \frac{1}{5} \log(1) + 0 \right\} \\ &= \frac{1}{5} \log(6) - \frac{3}{\sqrt{5}} \times \tan^{-1} \sqrt{5}\end{aligned}$$

39. Question

Evaluate the following definite Integrals:

$$\int_0^2 \frac{1}{4 + x - x^2} dx$$

Answer

$$= \int_0^2 \frac{1}{4 + x - x^2} dx$$

Since it is a quadratic equation we are trying to make it a complete square

$$\begin{aligned}
&= \int_0^2 \frac{1}{-(-4-x+x^2)} dx \\
&= \int_0^2 \frac{1}{-(-4-x+x^2 + \frac{1}{4} - \frac{1}{4})} dx \\
&= \int_0^2 \frac{1}{-((x-\frac{1}{2})^2 - \frac{17}{4})} dx \\
&= \int_0^2 \frac{1}{(\frac{\sqrt{17}}{2})^2 - ((x-\frac{1}{2})^2)} dx
\end{aligned}$$

=here the equation is in the form of integral of $\frac{1}{x^2+a^2}$ the integral is equal to $\frac{1}{2a} \times \log(\frac{a+x}{a-x})$

Here let us assume that $t=x-\frac{1}{2}$

So that $dx=dt$

When $x=0 t=-1/2$

And when $x=-1/2, t=3/2$

$$\begin{aligned}
\int_0^2 \frac{1}{(\frac{\sqrt{17}}{2})^2 - ((x-\frac{1}{2})^2)} dx &= \int_{-\frac{1}{2}}^{\frac{3}{2}} \frac{1}{(\frac{\sqrt{17}}{2})^2 - (t)^2} dt \\
&= \frac{1}{2\frac{\sqrt{17}}{2}} \times \log\left(\frac{\frac{\sqrt{17}}{2}+t}{\frac{\sqrt{17}}{2}-t}\right) \Big|_{-\frac{1}{2}}^{\frac{3}{2}} \\
&= \frac{1}{\sqrt{17}} \times [\log\left(\frac{\frac{\sqrt{17}}{2}+\frac{3}{2}}{\frac{\sqrt{17}}{2}-\frac{1}{2}}\right) - \log\left(\frac{\frac{\sqrt{17}}{2}-\frac{1}{2}}{\frac{\sqrt{17}}{2}+\frac{3}{2}}\right)] \\
&= \frac{1}{\sqrt{17}} \times [\log\left(\frac{\sqrt{17}+3}{\sqrt{17}-3}\right) - \log\left(\frac{\sqrt{17}-1}{\sqrt{17}+1}\right)] \\
&= \frac{1}{\sqrt{17}} \times [\log\left(\frac{\sqrt{17}+3}{\sqrt{17}-3}\right) \times \left(\frac{\sqrt{17}+1}{\sqrt{17}-1}\right)] \\
&= \frac{1}{\sqrt{17}} \times \log\left(\frac{4\sqrt{17}+17+3}{17-4\sqrt{17}+3}\right) \\
&= \frac{1}{\sqrt{17}} \times \log\left(\frac{\sqrt{17}+5}{-\sqrt{17}+5}\right)
\end{aligned}$$

Now rationalize the denominator

We get as

$$= \frac{1}{\sqrt{17}} \times \log\left(\frac{5\sqrt{17}+21}{4}\right)$$

40. Question

Evaluate the following definite Integrals:

$$\int_0^1 \frac{1}{2x^2+x+1} dx$$

Answer

Since the denominator is a quadratic equation let us make it in form of a perfect square

$$2x^2 + x + 1 = (x^2 + \frac{x}{2} + \frac{1}{2}) \times 2$$

$$= ((x + \frac{1}{4})^2 + \frac{1}{2} - \frac{1}{16}) \times 2$$

$$= ((x + \frac{1}{4})^2 + \frac{7}{16}) \times 2$$

$$= ((x + \frac{1}{4})^2 + (\frac{\sqrt{7}}{4})^2) \times 2$$

Now the equation

$$\int_0^1 \frac{1}{2x^2 + x + 1} = \int_0^1 \frac{1}{((x + \frac{1}{4})^2 + (\frac{\sqrt{7}}{4})^2) \times 2}$$

$$= \frac{1}{2} \int_0^1 \frac{1}{((x + \frac{1}{4})^2 + (\frac{\sqrt{7}}{4})^2)}$$

derivative of $\tan^{-1} \frac{x}{a} = \frac{1}{a^2+x^2}$ using this we can write it as

$$= \frac{4}{2\sqrt{7}} [\tan^{-1}(\frac{x + \frac{1}{4}}{\frac{\sqrt{7}}{4}})]_0^1$$

$$= \{\frac{2}{\sqrt{7}} \{ \tan^{-1} \frac{5}{\sqrt{7}} - \tan^{-1} \frac{1}{\sqrt{7}} \}\}$$

41. Question

Evaluate the following definite Integrals:

$$\int_0^1 \sqrt{x(1-x)} dx$$

Answer

To solve these kinds of equations we generally take $x = \sin^2 \theta, \cos^2 \theta$

So now here let $x = \sin^2 \theta$

So now $dx = 2\sin\theta\cos\theta d\theta$

Now change the limits

$x=0$ then $\theta=0$

$x=1$ then $\theta=\frac{\pi}{2}$

So it is equal to

$$= \int_0^{\frac{\pi}{2}} \sqrt{\sin^2 \theta (1 - \sin^2 \theta)} 2 \sin\theta\cos\theta d\theta$$

$$= \int_0^{\frac{\pi}{2}} 2 \sin^2 \theta \cos^2 \theta d\theta$$

$$= \int_0^{\frac{\pi}{2}} 4 \sin^2 \theta \cos^2 \theta d\theta \cdot \frac{1}{2}$$

now use formula $\sin 2x = 2 \sin x \cos x$

$$= \frac{1}{2} \int_0^{\frac{\pi}{2}} \sin^2 2\theta d\theta$$

Now use formula that $\cos 2x = 1 - 2 \sin^2 \theta$

$$= \frac{1}{2} \int_0^{\frac{\pi}{2}} \frac{1 - \cos 4\theta}{2} d\theta$$

$$= \frac{1}{4} \int_0^{\frac{\pi}{2}} (1 - \cos 4\theta) d\theta$$

$$= \frac{1}{4} \int_0^{\frac{\pi}{2}} d\theta - \int_0^{\frac{\pi}{2}} \cos 4\theta d\theta \frac{1}{4}$$

Now let us recall other formula :

$$\int \sin x = -\cos x \text{ and } \int \cos x = \sin x$$

$$= \frac{1}{4} [\theta]_0^{\frac{\pi}{2}} - \frac{1}{4} [\frac{\sin 4\theta}{4}]_0^{\frac{\pi}{2}}$$

Now recall that $\sin 0 = 0, \cos 0 = 1$

$$= \frac{1}{4} [\frac{\pi}{2} - \frac{1}{4} \times \sin 4 \times \frac{\pi}{2} - \{0 - \frac{1}{4} \times \sin 4 \times 0\}]$$

$$= \frac{\pi}{8} - 0 - \{0 - 0\}$$

$$= \frac{\pi}{8}$$

42. Question

Evaluate the following definite Integrals:

$$\int_0^2 \frac{1}{\sqrt{3+2x-x^2}} dx$$

Answer

Here the equation is of form that a quadratic equation is in the root so now to solve this make the equation in the root in the form of $a^2 - x^2, a^2 + x^2$

$$\text{Here } 3 + 2x - x^2 = 3 + 1 - (1 - 2x + x^2)$$

$$= 4 - (x-1)^2$$

$$= 2^2 - (x-1)^2$$

Now just recall a formula that is derivative of $\sin^{-1} \frac{x}{a} = \frac{1}{\sqrt{a^2 - x^2}}$

Here $a=2$ and $x=x-1$

Now we get

$$\int_0^2 \frac{1}{\sqrt{3+2x-x^2}} dx = [\sin^{-1} \frac{x-1}{2}]_0^2$$

$$= \sin^{-1} \frac{2-1}{2} - \sin^{-1} \frac{0-1}{2}$$

$$= \frac{\pi}{6} + \frac{\pi}{6}$$

$$= \frac{\pi}{3}$$

43. Question

Evaluate the following definite Integrals:

$$\int_0^4 \frac{1}{\sqrt{4x-x^2}} dx$$

Answer

Here first we are converting the quadratic equation in to a perfect square

$$4x - x^2 = 4 - 4 + 4x - x^2$$

$$= 4 - (4 - 4x + x^2)$$

$$= 4 - (x - 2)^2$$

$$= (2)^2 - (x - 2)^2$$

Now just recall a formula that is derivative of $\sin^{-1} \frac{x}{a} = \frac{1}{\sqrt{a^2-x^2}}$

Here $a=2$ and $x=x-2$

Now we get

$$\int_0^4 \frac{1}{\sqrt{4x-x^2}} dx = [\sin^{-1} \frac{x-2}{2}]_0^4$$

$$= \sin^{-1} \frac{4-2}{2} - \sin^{-1} \frac{0-2}{2}$$

$$= \sin^{-1} 1 - \sin^{-1} -1$$

$$= \frac{\pi}{2} - (-\frac{\pi}{2})$$

$$= \pi$$

44. Question

Evaluate the following definite Integrals:

$$\int_{-1}^1 \frac{1}{x^2+2x+5} dx$$

Answer

Now denominator is in a quadratic form so let us make it in other form

$$x^2 + 2x + 5 = x^2 + 2x + 1 + 4$$

$$= (x+1)^2 + 4$$

$$= (x+1)^2 + 2^2$$

Recall a formula $\tan^{-1} \left(\frac{x}{a} \right) = \frac{a}{a^2+x^2}$

$$\text{So now } \int_{-1}^1 \frac{1}{x^2+2x+5} = [\tan^{-1} \left(\frac{x+1}{2} \right)]_{-1}^1 \times \frac{1}{2}$$

$$= (\tan^{-1} \left(\frac{1+1}{2} \right) - \tan^{-1} \left(\frac{-1+1}{2} \right)) \times \frac{1}{2}$$

$$= (\tan^{-1}(1) - \tan^{-1}(0)) \times \frac{1}{2}$$

$$= \frac{\pi}{4} \times \frac{1}{2} = \frac{\pi}{8}$$

45. Question

Evaluate the following definite Integrals:

$$\int_1^4 \frac{x^2 + x}{\sqrt{2x+1}} dx$$

Answer

To solve this let us assume that $2x+1=t^2$

$$2 dx = 2t dt$$

$$\text{So now } x=1, t=\sqrt{3}$$

$$x=4, t=3$$

So now after substitution we get

$$\begin{aligned} \int_1^4 \frac{x^2 + x}{\sqrt{2x+1}} dx &= \int_{\sqrt{3}}^3 \frac{\frac{t^2 - 1}{2})^2 + \frac{t^2 - 1}{2}}{t} t dt \\ &= \frac{1}{4} \int_{\sqrt{3}}^3 (t^4 - 2t^2 + 1 + 2t^2 - 2) dt \\ &= \frac{1}{4} \int_{\sqrt{3}}^3 (t^4 - 1) dt \end{aligned}$$

$$\text{Now let us recall other formula i.e. } \int x^n = \frac{x^{n+1}}{n+1}$$

$$\begin{aligned} &= \frac{1}{4} \left[\frac{t^5}{5} - t \right]_{\sqrt{3}}^3 \\ &= \frac{1}{4} \left[\frac{3^5}{5} - 3 \right] - \frac{1}{4} \left[\frac{\sqrt{3}^5}{5} - \sqrt{3} \right] \\ &= \frac{57 - \sqrt{3}}{5} \end{aligned}$$

46. Question

Evaluate the following definite Integrals:

$$\int_0^1 x(1-x)^5 dx$$

Answer

Now to make it simpler problem let us expand $(1-x)^5$ using binomial theorem

$$\text{So } (1-x)^5 = 1 - 5x + 10x^2 - 10x^3 + 5x^4 - x^5$$

$$\text{Let us also recall other formula i.e., } \int x^n = \frac{x^{n+1}}{n+1}$$

So now

$$\begin{aligned}
\int_0^1 x(1-x)^5 dx &= \int_0^1 x(1-5x+10x^2-10x^3+5x^4-x^5) dx \\
&= \left[\frac{x^2}{2} - \frac{5x^3}{3} + \frac{10x^4}{4} - \frac{10x^5}{5} + \frac{5x^6}{6} - \frac{x^7}{7} \right]_0^1 \\
&= \frac{1}{2} - \frac{5}{3} + \frac{10}{4} - \frac{10}{5} + \frac{5}{6} - \frac{1}{7} - \left\{ \frac{0}{2} - \frac{0}{3} + \frac{0}{4} - \frac{0}{5} + \frac{0}{6} - \frac{0}{7} \right\} \\
&= \frac{1}{42}
\end{aligned}$$

47. Question

Evaluate the following definite Integrals:

$$\int_1^2 \left(\frac{x-1}{x^2} \right) e^x dx$$

Answer

$$\int_1^2 \frac{x-1}{x^2} e^x dx = \int_1^2 \frac{1}{x} e^x dx - \int_1^2 \frac{1}{x^2} e^x dx$$

For this we have to apply integration by parts

Let u and v be two functions then

$$\int u dv = uv - \int v du$$

To choose the first function u we use "ILATE" rule

That is

I=inverse trigonometric function

L=logarithmic function

A=algebraic function

T=trigonometric functions

E=exponential function

So in this preference, the first function is chosen to make the integration simpler.

Now, In the given question x^2 is an algebraic function and it is chosen as u(A comes first in "ILATE" rule)

So first let us integrate the equation and then let us substitute the limits in it.

Here we are expanding only first integral first

$$\begin{aligned}
\int_1^2 \frac{1}{x} e^x dx - \int_1^2 \frac{1}{x^2} e^x dx &= \frac{1}{x} \int_1^2 e^x dx - \int_1^2 \int e^x dx \times \frac{d \frac{1}{x}}{dx} - \int_1^2 \frac{1}{x^2} e^x dx \\
&= \left[\frac{e^x}{x} \right]_1^2 + \int_1^2 \frac{1}{x^2} e^x dx - \int_1^2 \frac{1}{x^2} e^x dx \\
&= \left[\frac{e^x}{x} \right]_1^2 \\
&= \frac{e^2}{2} - \frac{e^1}{1}
\end{aligned}$$

$$= \frac{e^2}{2} - e$$

48. Question

Evaluate the following definite Integrals:

$$\int_0^1 \left(xe^{2x} + \sin \frac{\pi x}{2} \right) dx$$

Answer

First split the integral $\int_0^1 (xe^{2x}) dx + \int_0^1 (\sin \frac{\pi x}{2}) dx$

Now integrate by parts the first one

For this we have to apply integration by parts

Let u and v be two functions then

$$\int u dv = uv - \int v du$$

To choose the first function u we use "ILATE" rule

That is

I=inverse trigonometric function

L=logarithmic function

A=algebraic function

T=trigonometric functions

E=exponential function

So in this preference, the first function is chosen to make the integration simpler.

Now, In the given question x^2 is an algebraic function and it is chosen as u(A comes first in "ILATE" rule)

So first let us integrate the equation and then let us substitute the limits in it.

Remember $\int \sin x = -\cos x$ and $\int \cos x = \sin x$

$$= \int_0^1 (xe^{2x}) dx + \int_0^1 (\sin \frac{\pi x}{2}) dx$$

$$= \frac{xe^x}{2} - \frac{1}{2} \int e^{2x} + \frac{2}{\pi} [1 - 0]$$

$$= \frac{2e^2}{2} - \frac{e^2}{2} + \frac{1}{4} + \frac{2}{\pi}$$

$$= \frac{2e^2}{2} + \frac{1}{4} + \frac{2}{\pi}$$

49. Question

Evaluate the following definite Integrals:

$$\int_0^1 \left(xe^x + \cos \frac{\pi x}{4} \right) dx$$

Answer

First split the integral $\int_0^1 (xe^x) dx + \int_0^1 (\cos \frac{\pi x}{4}) dx$

Now integrate by parts the first one

For this we have to apply integration by parts

Let u and v be two functions then

$$\int u dv = uv - \int v du$$

To choose the first function u we use "ILATE" rule

That is

I=inverse trigonometric function

L=logarithmic function

A=algebraic function

T=trigonometric functions

E=exponential function

So in this preference, the first function is chosen to make the integration simpler.

Now, In the given question x^2 is an algebraic function and it is chosen as u(A comes first in "ILATE" rule)

So first let us integrate the equation and then let us substitute the limits in it.

Remember $\int \sin x = -\cos x$ and $\int \cos x = \sin x$

$$\begin{aligned} &= \int_0^1 (xe^x) dx + \int_0^1 (\cos \frac{\pi x}{4}) dx \\ &= [xe^x]_0^1 - \int e^x + \frac{4}{\pi} [\cos \frac{\pi x}{4}]_0^1 \\ &= [xe^x]_0^1 - [e^x]_0^1 + \frac{4}{\pi} [\cos \frac{\pi x}{4}]_0^1 \\ &= [e - 0] - [e - 0] + \frac{4}{\pi} [\cos \frac{\pi}{4} - \cos \frac{\pi \times 0}{4}] \\ &= 1 + \frac{2\sqrt{2}}{\pi} \end{aligned}$$

50. Question

Evaluate the following definite Integrals:

$$\int_{\pi/2}^{\pi} e^x \left(\frac{1 - \sin x}{1 - \cos x} \right) dx$$

Answer

Now using the formula

$$\sin 2x = 2 \sin x \cos x$$

$$\cos 2x = 1 - 2 \sin^2 x$$

$$\begin{aligned} \int_{\pi/2}^{\pi} e^x \left(\frac{1 - \sin x}{1 - \cos x} \right) dx &= \int_{\pi/2}^{\pi} e^x \left(\frac{1 - 2 \sin \frac{x}{2} \cos \frac{x}{2}}{2 \sin^2 \frac{x}{2}} \right) dx \\ &= - \int_{\pi/2}^{\pi} e^x \left(-\frac{1}{2} \operatorname{cosec}^2 \frac{x}{2} + \cot \frac{x}{2} \right) dx \end{aligned}$$

Here we know that derivative of $\cot\frac{x}{2}$ is $= -\frac{1}{2}\operatorname{cosec}^2\frac{x}{2}$

And it is in the form of e^x (function + derivative of function) so the equation integral will be e^x function

$$\begin{aligned}&= -[e^x \cot\frac{x}{2}]_{\frac{\pi}{2}}^{\frac{\pi}{2}} \\&= -[e^{\frac{\pi}{2}} \cot\frac{\pi}{2} - e^{\frac{\pi}{2}} \cot\frac{\pi}{2}]\end{aligned}$$

$\cot 90 = 0, \cot 45 = 1$

$$= e^{\frac{\pi}{2}}$$

51. Question

Evaluate the following definite Integrals:

$$\int_0^{2\pi} e^{x/2} \sin\left(\frac{x}{2} + \frac{\pi}{4}\right) dx$$

Answer

We know that $\sin\left(\frac{x}{2} + \frac{\pi}{4}\right) = \sin\left(\frac{x}{2}\right)\cos\left(\frac{\pi}{4}\right) + \sin\left(\frac{\pi}{4}\right)\cos\left(\frac{x}{2}\right)$

So the equation will be

$$= \int_0^{2\pi} e^{\frac{x}{2}} \left(\sin\left(\frac{x}{2}\right) \cos\left(\frac{\pi}{4}\right) + \sin\left(\frac{\pi}{4}\right) \cos\left(\frac{x}{2}\right) \right) dx$$

We know that $\cos 45 = \sin 45 = \frac{1}{\sqrt{2}}$

Substitute it

$$\begin{aligned}&= \int_0^{2\pi} e^{\frac{x}{2}} \left(\sin\left(\frac{x}{2}\right) \frac{1}{\sqrt{2}} + \cos\left(\frac{x}{2}\right) \frac{1}{\sqrt{2}} \right) dx \\&= \frac{1}{\sqrt{2}} \left(\int_0^{2\pi} e^{\frac{x}{2}} \left(\sin\left(\frac{x}{2}\right) + \cos\left(\frac{x}{2}\right) \right) dx \right) \\&= \frac{1}{\sqrt{2}} \left(\int_0^{2\pi} e^{\frac{x}{2}} \sin\left(\frac{x}{2}\right) dx \right) + \frac{1}{\sqrt{2}} \left(\int_0^{2\pi} e^{\frac{x}{2}} \cos\left(\frac{x}{2}\right) dx \right)\end{aligned}$$

Remember that $\int \sin x = -\cos x$ and $\int \cos x = \sin x$

Now integrate by parts For this we have to apply integration by parts

Let u and v be two functions then

$$\int u dv = uv - \int v du$$

To choose the first function u we use "ILATE" rule

That is

I=inverse trigonometric function

L=logarithmic function

A=algebraic function

T=trigonometric functions

E=exponential function

So in this preference, the first function is chosen to make the integration simpler.

Now, In the given question x^2 is an algebraic function and it is chosen as u(A comes first in "ILATE" rule)

So first let us integrate the equation and then let us substitute the limits in it.

$$\begin{aligned} &= \frac{1}{\sqrt{2}} \left\{ \sin\left(\frac{x}{2}\right) \int_0^{2\pi} e^{\frac{x}{2}} dx - \int_0^{2\pi} \int_0^x e^{\frac{x}{2}} dx d \sin\left(\frac{x}{2}\right) + \frac{1}{\sqrt{2}} \left\{ \cos\left(\frac{x}{2}\right) \int_0^{2\pi} e^{\frac{x}{2}} dx - \right. \right. \\ &\quad \left. \left. \int_0^{2\pi} \int_0^x e^{\frac{x}{2}} dx d \cos\left(\frac{x}{2}\right) \right\} \right. \\ &= \left[\frac{1}{\sqrt{2}} \left\{ \sin\left(\frac{x}{2}\right) 2e^{\frac{x}{2}} \right\} \right]_0^{2\pi} \\ &= \frac{1}{\sqrt{2}} \left\{ \sin\left(\frac{2\pi}{2}\right) 2e^{\frac{2\pi}{2}} - \frac{1}{\sqrt{2}} \left\{ \sin\left(\frac{0}{2}\right) 2e^{\frac{2\times 0}{2}} \right\} \right\} \\ &= 0 - 0 = 0 \end{aligned}$$

52. Question

Evaluate the following definite Integrals:

$$\int_0^{2\pi} e^x \cos\left(\frac{\pi}{4} + \frac{x}{2}\right) dx$$

Answer

Now let us use integration by parts

For this we have to apply integration by parts

Let u and v be two functions then

$$\int u dv = uv - \int v du$$

To choose the first function u we use "ILATE" rule

That is

I=inverse trigonometric function

L=logarithmic function

A=algebraic function

T=trigonometric functions

E=exponential function

So in this preference, the first function is chosen to make the integration simpler.

Now, In the given question x^2 is an algebraic function and it is chosen as u(A comes first in "ILATE" rule)

So first let us integrate the equation and then let us substitute the limits in it.

Remember that $\int \sin x = -\cos x$ and $\int \cos x = \sin x$

$$\begin{aligned} &= [\cos\left(\frac{x}{2} + \frac{\pi}{4}\right) e^x]_0^{2\pi} + \frac{1}{2} \int_0^{2\pi} e^x \sin\left(\frac{x}{2} + \frac{\pi}{4}\right) dx \\ L &= [\cos\left(\frac{x}{2} + \frac{\pi}{4}\right) e^x]_0^{2\pi} + \frac{1}{2} \left\{ [\sin\left(\frac{x}{2} + \frac{\pi}{4}\right) e^x]_0^{2\pi} - \frac{1}{2} \int_0^{2\pi} e^x \cos\left(\frac{x}{2} + \frac{\pi}{4}\right) dx \right\} \\ &\quad \} \\ L &= \left[\left\{ \cos\left(\frac{5\pi}{4}\right) e^{2\pi} - \cos\left(\frac{\pi}{4}\right) \right\} + \frac{1}{2} \left[\sin\left(\frac{5\pi}{4}\right) e^{2\pi} - \sin\left(\frac{\pi}{4}\right) \right] \right] - \frac{L}{4} \end{aligned}$$

$$\frac{5L}{4} = -\frac{1}{\sqrt{2}}(e^{2\pi} + 1) - \frac{1}{2\sqrt{2}}(e^{2\pi} + 1)$$

$$L = -\frac{3\sqrt{2}}{5}(e^{2\pi} + 1)$$

53. Question

Evaluate the following definite Integrals:

$$\int_0^{\pi} e^{2x} \sin\left(\frac{\pi}{4} + x\right) dx$$

Answer

$$\begin{aligned} \text{let } I &= \int_0^{\pi} e^{2x} \sin\left(x + \frac{\pi}{4}\right) dx \\ &= \sin\left(x + \frac{\pi}{4}\right) \int_0^{\pi} e^{2x} dx - \int_0^{\pi} e^{2x} d(\sin\left(x + \frac{\pi}{4}\right)) dx \text{ limit 0 to pi} \\ &= \sin\left(x + \frac{\pi}{4}\right) \frac{e^{2x}}{2} - \cos\left(x + \frac{\pi}{4}\right) \frac{e^{2x}}{4} - \frac{\int_0^{\pi} e^{2x} \sin\left(x + \frac{\pi}{4}\right) dx}{4} \\ \frac{5I}{4} &= \sin\left(x + \frac{\pi}{4}\right) \frac{e^{2x}}{2} - \cos\left(x + \frac{\pi}{4}\right) \frac{e^{2x}}{4} \text{ with limits 0 to pi} \\ I &= \frac{4}{5} [\sin\left(x + \frac{\pi}{4}\right) \frac{e^{2x}}{2} - \cos\left(x + \frac{\pi}{4}\right) \frac{e^{2x}}{4}]_0^{\pi} \\ &= \sin\left(\pi + \frac{\pi}{4}\right) \frac{e^{2\pi}}{2} - \cos\left(\pi + \frac{\pi}{4}\right) \frac{e^{2\pi}}{4} - \{\sin\left(0 + \frac{\pi}{4}\right) \frac{e^0}{2} - \cos\left(0 + \frac{\pi}{4}\right) \frac{e^0}{4}\} \\ &= -\frac{1}{\sqrt{2}} \frac{e^{2\pi}}{2} - \frac{1}{\sqrt{2}} \frac{e^{2\pi}}{4} - \left\{ \frac{1}{2\sqrt{2}} - \frac{1}{4\sqrt{2}} \right\} \\ &= -\frac{3}{\sqrt{2}} \frac{e^{2\pi}}{4} - \frac{1}{4\sqrt{2}} \end{aligned}$$

54. Question

Evaluate the following definite Integrals:

$$\int_0^1 \frac{1}{\sqrt{1+x} - \sqrt{x}} dx$$

Answer

Now let it be taken as I

$$I = \int_0^1 \frac{1}{\sqrt{1+x} - \sqrt{x}} dx$$

Now rationalize the denominator

$$\begin{aligned} I &= \int_0^1 \frac{1}{\sqrt{1+x} - \sqrt{x}} \times \frac{\sqrt{1+x} + \sqrt{x}}{\sqrt{1+x} + \sqrt{x}} dx \\ &= \int_0^1 \frac{\sqrt{1+x} + \sqrt{x}}{1+x-x} dx \\ &= \int_0^1 \frac{\sqrt{1+x} + \sqrt{x}}{1} dx \end{aligned}$$

Let us also recall formula $\int x^n dx = \frac{x^{n+1}}{n+1}$

$$= \int_0^1 \sqrt{1+x} dx + \int_0^1 \sqrt{x} dx$$

$$= \frac{2}{3} [(1+x)^{\frac{3}{2}}]_0^1 + \frac{2}{3} [(x)^{\frac{3}{2}}]_0^1$$

$$= \frac{2}{3} ((2)^{\frac{3}{2}} - 1) + \frac{2}{3}$$

$$= \frac{2}{3} ((2)^{\frac{3}{2}})$$

$$= \frac{4\sqrt{2}}{3}$$

55. Question

Evaluate the following definite Integrals:

$$\int_1^2 \frac{x}{(x+1)(x+2)} dx$$

Answer

$$= \int_1^2 \frac{2x+2-(x+1)}{(x+1)(x+2)} dx$$

$$= \int_1^2 \frac{2}{(x+2)} dx - \int_1^2 \frac{1}{(x+1)} dx$$

$$\text{Remember derivative of } \log x = \frac{1}{x}$$

So using that

$$= [-\log(x+1) + 2\log(x+2)]_1^2$$

Substitute upper limit and then subtract the lower limit from it

$$= -(\log 3 - \log 2) + 2(\log 4 - \log 3)$$

$$= -3\log 3 + 5\log 2$$

$$= \log\left(\frac{32}{27}\right)$$

56. Question

Evaluate the following definite Integrals:

$$\int_0^{\pi/2} \sin^3 x dx$$

Answer

$$\text{Let } I = \int_0^{\frac{\pi}{2}} \sin^3 x dx$$

$$= \int_0^{\frac{\pi}{2}} \sin^2 x \sin x dx$$

$$= \int_0^{\frac{\pi}{2}} (1 - \cos^2 x) \sin x dx$$

$$= \int_0^{\frac{\pi}{2}} \sin x \, dx - \int_0^{\frac{\pi}{2}} (\cos^2 x) \sin x \, dx$$

Remember that $\int \sin x = -\cos x$ and $\int \cos x = \sin x$

$$= [-\cos x]_0^{\frac{\pi}{2}} + [\frac{\cos^3 x}{3}]_0^{\frac{\pi}{2}}$$

$$= -\cos \frac{\pi}{2} - -\cos 0 + \{\frac{\cos^3 \frac{\pi}{2}}{3} - \frac{\cos^3 0}{3}\}$$

$$\cos 90 = 0, \cos 0 = 1$$

$$= -0 + 1 - \frac{1}{3} = \frac{2}{3}$$

57. Question

Evaluate the following definite Integrals:

$$\int_0^{\pi} \left(\sin^2 \frac{x}{2} - \cos^2 \frac{x}{2} \right) dx$$

Answer

$$I = \int_0^{\pi} \sin^2 \frac{x}{2} - \cos^2 \frac{x}{2} dx$$

$$= - \int_0^{\pi} \cos^2 \frac{x}{2} - \sin^2 \frac{x}{2} dx$$

$$= - \int_0^{\pi} \cos x dx$$

Because we have a formula $\cos x = \cos^2 \frac{x}{2} - \sin^2 \frac{x}{2}$

Remember that $\int \sin x = -\cos x$ and $\int \cos x = \sin x$

$$= [-\sin x]_0^{\pi}$$

$$= [-(\sin \pi - \sin 0)]$$

$$\sin 180 = 0, \sin 0 = 0$$

$$= 0$$

58. Question

Evaluate the following definite Integrals:

$$\int_1^2 e^{2x} \left(\frac{1}{x} - \frac{1}{2x^2} \right) dx$$

Answer

Let $2x=t$ then $2dx=dt$

When $x=1 t=2$

And when $x=2, t=4$

$$\int_1^2 e^{2x} \left(\frac{1}{x} - \frac{1}{2x^2} \right) dx = \frac{1}{2} \int_2^4 e^t \left(\frac{2}{t} - \frac{2}{t^2} \right) dt$$

$$= \int_2^4 e^t \left(\frac{1}{t} - \frac{1}{t^2} \right) dt$$

We can observe here that

$$\text{Derivative of } \frac{1}{x} = -\frac{1}{x^2}$$

Now it is in the form $e^x(\text{function} + \text{derivative of function})$

So the integral will be $e^x(\text{function})$

$$\begin{aligned} \int_2^4 e^t \left(\frac{1}{t} - \frac{1}{t^2} \right) dt &= \left[\frac{e^t}{t} \right]_2^4 \\ &= \frac{e^4}{4} - \frac{e^2}{2} \\ &= \frac{(e^4 - 2e^2)}{4} \end{aligned}$$

59. Question

Evaluate the following definite Integrals:

$$\int_1^2 \frac{1}{(x-1)(2-x)} dx$$

Answer

Let us solve the denominator

$$(x-1)(2-x) = 2x - 2 - x^2 + x$$

$$= 3x - x^2 - 2$$

$$= -(x - \frac{2}{3})^2 + \frac{1}{4}$$

$$\int_1^2 \frac{1}{\sqrt{(x-1)(2-x)}} dx = \int_1^2 \frac{1}{\sqrt{-(x - \frac{2}{3})^2 + \frac{1}{4}}} dx$$

$$= \int_1^2 \frac{1}{\sqrt{-\left(x - \frac{2}{3}\right)^2 + \frac{1}{2}}} dx$$

$$= \int_1^2 \frac{1}{\sqrt{\frac{1}{2} - \left(x - \frac{2}{3}\right)^2}} dx$$

Now just recall a formula that is derivative of $\sin^{-1} \frac{x}{a} = \frac{1}{\sqrt{a^2 - x^2}}$

$$= [\sin^{-1}(2x - 3)]_1^2$$

$$= \sin^{-1} 1 - \sin^{-1} -1$$

$$= \pi$$

60. Question

$$\text{If } \int_0^k \frac{1}{2+8x^2} dx = \frac{\pi}{16}, \text{ find value of k}$$

Answer

Given that $\int_0^k \frac{1}{2+8x^2} dx = \frac{\pi}{16}$, $k=?$

$$\int_0^k \frac{1}{2+8x^2} dx = \frac{\pi}{16}$$

$$\frac{1}{8} \int_0^k \frac{1}{(1/2)^2 + x^2} dx = \frac{\pi}{16}$$

$$\text{derivative of } \tan^{-1} \frac{x}{a} = \frac{1}{a^2+x^2}$$

$$\frac{1}{8} [2 \tan^{-1} 2x]_0^k = \frac{\pi}{16}$$

$$\frac{1}{8} [\tan^{-1} 2k - \tan^{-1} 0] = \frac{\pi}{16}$$

$$\tan^{-1} 2x = \frac{\pi}{4}$$

$$2k=1$$

$$k = \frac{1}{2}$$

61. Question

If $\int_0^a 3x^2 dx = 8$, find the value of a.

Answer

$$\int_0^a 3x^2 dx = 8$$

Let us also recall formula $\int x^n dx = \frac{x^{n+1}}{n+1}$

$$[x^3]_0^a = 8$$

$$a^3 = 8$$

$$a=2$$

$$\text{hence } a=2.$$

62. Question

Evaluate the following Integrals:

$$\int_{\pi}^{3\pi/2} \sqrt{1 - \cos 2x} dx$$

Answer

$$\int_{\pi}^{3\pi/2} \sqrt{1 - \cos 2x} dx$$

We know that $\cos 2x = 1 - 2\sin^2 x$

Now substitute that in the equation

We get

$$\begin{aligned} \int_{\pi}^{\frac{3\pi}{2}} \sqrt{1 - \cos 2x} dx &= \int_{\pi}^{\frac{3\pi}{2}} \sqrt{1 - (1 - 2\sin^2 x)} dx \\ &= \int_{\pi}^{\frac{3\pi}{2}} \sqrt{2\sin^2 x} dx \\ &= \sqrt{2} \int_{\pi}^{\frac{3\pi}{2}} \sin x dx \end{aligned}$$

We already know that integral of $\sin x$ is $-\cos x$

$$\begin{aligned} &= \sqrt{2} (-\cos x) \Big|_{\pi}^{\frac{3\pi}{2}} \\ &= -\sqrt{2} (\cos \pi - \cos \pi) \\ &= -\sqrt{2} (0 - (-1)) \\ &= -\sqrt{2} \end{aligned}$$

63. Question

Evaluate the following Integrals:

$$\int_0^{2\pi} \sqrt{1 + \sin \frac{x}{2}} dx$$

Answer

$$\text{Let } I = \int_0^{2\pi} \sqrt{1 + \sin(\frac{x}{2})} dx$$

$$\text{Let us recall that } \sin^2(\frac{x}{2}) + \cos^2(\frac{x}{2}) = 1$$

$$\text{And } \sin(\frac{x}{2}) = 2\sin(\frac{x}{4})\sin(\frac{x}{4})$$

$$I = \int_0^{2\pi} \sqrt{\sin^2(\frac{x}{2}) + \cos^2(\frac{x}{2}) + 2\sin(\frac{x}{4})\cos(\frac{x}{4})} dx$$

$$I = \int_0^{2\pi} \sqrt{(\sin(\frac{x}{4}) + \cos(\frac{x}{4}))^2} dx$$

$$= \int_0^{2\pi} (\sin(\frac{x}{4}) + \cos(\frac{x}{4})) dx$$

Recall: $\int \sin x = -\cos x$ and $\int \cos x = \sin x$

$$\begin{aligned} I &= \left[-\frac{\cos(\frac{x}{4})}{4} + \frac{\sin(\frac{x}{4})}{4} \right]_0^{2\pi} \\ &= -\frac{\cos(\frac{2\pi}{4})}{4} + \frac{\sin(\frac{2\pi}{4})}{4} - \left\{ -\frac{\cos(\frac{0}{4})}{4} + \frac{\sin(\frac{0}{4})}{4} \right\} \\ &= 4(0+1+1-0) \\ &= 8 \end{aligned}$$

64. Question

Evaluate the following Integrals:

$$\int_0^{\pi/4} (\tan x + \cot x)^{-2} dx$$

Answer

$$I = \int_0^{\pi/4} (\tan x + \cot x)^{-2} dx$$

$$= \int_0^{\pi/4} \frac{1}{(\tan x + \cot x)^2} dx$$

We know that $\tan x = \frac{\sin x}{\cos x}$ and $\cot x = \frac{\cos x}{\sin x}$

Now substitute them in the equation.

$$= \int_0^{\pi/4} \frac{1}{\left(\frac{\sin x}{\cos x} + \frac{\cos x}{\sin x}\right)^2} dx$$

Let us recall that $\sin^2(x) + \cos^2(x) = 1$

$$= \int_0^{\pi/4} (\sin x \cos x)^2 dx$$

Again using $\cos^2(x) = 1 - \sin^2(x)$

$$= \int_0^{\pi/4} (\sin x)^2 (1 - \sin^2(x)) dx$$

$$= \int_0^{\pi/4} (\sin x)^2 dx - \int_0^{\pi/4} (\sin^4(x)) dx$$

Here we are using reduction formula of $\sin x$

$$\int (\sin x)^n dx = \frac{n-1}{n} \times \int (\sin x)^{n-2} dx - \frac{\cos x (\sin x)^{n-1}}{n}$$

For $n=2$

$$\int_0^{\pi/4} (\sin x)^2 dx = \frac{1}{2} \int_0^{\pi/4} dx - \frac{\cos x \sin x}{2}$$

$$= \frac{1}{2} \left[\frac{\pi}{4} \right] - \frac{1/2}{2}$$

$$= \frac{\pi}{8} - \frac{1}{4}$$

$$\int_0^{\pi/4} (\sin^4(x)) x dx = \frac{4-1}{4} \int_0^{\pi/4} (\sin^2(x)) x dx - \frac{\cos x \sin^3 x}{4} \quad [\text{limits } 0, \frac{\pi}{4}]$$

$$= \frac{3}{4} \left\{ \frac{x}{2} - \frac{\cos x \sin x}{2} \right\} - \frac{\cos x \sin^3 x}{4} \quad [\text{limits } 0, \frac{\pi}{4}]$$

Now substitute limits

$$\frac{3}{4} \left\{ \frac{\pi}{8} - \frac{\cos \frac{\pi}{4} \sin \frac{\pi}{4}}{2} \right\} - \frac{\cos \frac{\pi}{4} \sin^3 \frac{\pi}{4}}{4} - \frac{3}{4} \left\{ 0 - \frac{\cos 0 \sin 0}{2} \right\} + \frac{\cos 0 \sin^3 0}{4}$$

$$\sin 45^\circ = \cos 45^\circ = 0, \cos 0 = 1, \sin 0 = 0$$

$$= \frac{3}{4} \left\{ \frac{\pi}{8} - \frac{1}{4} \right\} - \frac{1}{16}$$

$$\text{Now } \int_0^{\frac{\pi}{4}} (\tan x + \cot x)^{-2} dx = \frac{\pi - 1}{8} - \left\{ \frac{3}{4} \left\{ \frac{\pi}{8} - \frac{1}{4} \right\} - \frac{1}{16} \right\}$$

$$= \frac{\pi}{32}$$

65. Question

Evaluate the following Integrals:

$$\int_0^1 x \log(1+2x) dx$$

Answer

Now let us use integration by parts

For this we have to apply integration by parts

Let u and v be two functions then

$$\int u dv = uv - \int v du$$

To choose the first function u we use "ILATE" rule

That is

I=inverse trigonometric function

L=logarithmic function

A=algebraic function

T=trigonometric functions

E=exponential function

So in this preference, the first function is chosen to make the integration simpler.

Now, In the given question x^2 is an algebraic function and it is chosen as u(A comes first in "ILATE" rule)

So first let us integrate the equation and then let us substitute the limits in it.

$$\begin{aligned} \int_0^1 x \log(1+2x) dx &= \left[\frac{x^2 \log(1+2x)}{2} \right]_0^1 - \int_0^1 \frac{2x^2}{2(2x+1)} dx \\ &= \left[\frac{\log(1+2)}{2} - 0 \right] - \int_0^1 \frac{x}{2} - \frac{1}{4} + \frac{1}{4(2x+1)} dx \\ &= \frac{\log 3}{2} - \left[\frac{x^2}{4} - \frac{x}{4} + \frac{1}{8} \log(|2x+1|) \right]_0^1 \\ &= \frac{\log 3}{2} - \left\{ \frac{1}{4} - \frac{1}{4} + \frac{1}{8} \log(|2+1|) - \left[\frac{0}{4} - \frac{0}{4} + \frac{1}{8} \log(|0+1|) \right] \right\} \\ &= \frac{\log 3}{2} - \frac{\log 3}{8} \\ &= \frac{3 \log 3}{8} \end{aligned}$$

66. Question

Evaluate the following Integrals:

$$\int_{\pi/6}^{\pi/3} (\tan x + \cot x)^2 dx$$

Answer

$$= \int_{\pi/6}^{\pi/3} \tan^2 x + 2\tan x \cot x + \cot^2 x dx$$

$$\text{recall : } \sec^2 x - \tan^2 x = 1, \cosec^2 x - \cot^2 x = 1$$

$$= \int_{\pi/6}^{\pi/3} \sec^2 x - 1 + 2 + \cosec^2 x - 1 dx$$

$$= \int_{\pi/6}^{\pi/3} \sec^2 x + \cosec^2 x dx$$

Integral $\sec^2 x$ is $\tan x$ and integral of $\cosec^2 x = -\cot x$

$$= [\tan x]_{\pi/6}^{\pi/3} - [\cot x]_{\pi/6}^{\pi/3}$$

$$\tan 30 = \cot 60 = \frac{1}{\sqrt{3}}$$

$$\tan 60 = \cot 30 = \sqrt{3}$$

$$= \tan 60 - \cot 60 - \{\tan 30 - \cot 30\}$$

$$= \sqrt{3} - \frac{1}{\sqrt{3}} - \left(\frac{1}{\sqrt{3}} - \sqrt{3}\right)$$

$$= \frac{4}{\sqrt{3}}$$

67. Question

Evaluate the following Integrals:

$$\int_0^{\pi/4} (a^2 \cos^2 x + b^2 \sin^2 x) dx$$

Answer

$$I = \int_0^{\pi/4} a^2 \cos^2(x) + b^2 \sin^2(x) dx$$

$$= \int_0^{\pi/4} a^2(1 - \sin^2(x)) + b^2 \sin^2(x) dx$$

$$= \int_0^{\pi/4} a^2 + (b^2 - a^2) \sin^2(x) dx$$

$$\text{Recall: } \sin^2(x) = \frac{1+\cos 2x}{2}$$

$$= \int_0^{\pi/4} a^2 + (b^2 - a^2) \frac{1 + \cos 2x}{2} dx$$

We know integral of $\cos x$ is $\sin x$

$$\begin{aligned}
&= \left[a^2 x + \frac{b^2 - a^2}{2} \times \left(x + \frac{\sin 2x}{2} \right) \right]_0^{\frac{\pi}{4}} \\
&= a^2 \frac{\pi}{4} + \frac{b^2 - a^2}{2} \times \left(\frac{\pi}{4} + \frac{\sin 2 \frac{\pi}{4}}{2} \right) - \{ a^2 \times 0 + \frac{b^2 - a^2}{2} \times (0 + \frac{\sin 0}{2}) \}
\end{aligned}$$

$$\sin 0 = 0, \sin 90 = 1$$

$$= (b^2 + a^2) \frac{\pi}{8} + \frac{(b^2 - a^2)}{4}$$

68. Question

Evaluate the following Integrals:

$$\int_0^1 \frac{1}{1+2x+2x^2+2x^3+x^4} dx$$

Answer

$$I = \int_0^1 \frac{1}{1+2x+2x^2+2x^3+x^4} dx$$

Now arranging denominator, we get as

$$1+2x+2x^2+2x^3+x^4 = (1+x)^2(x^2+1)$$

$$\begin{aligned}
&\int_0^1 \frac{1}{(1+x)^2(x^2+1)} dx \\
&= \int_0^1 \frac{-x}{2(x^2+1)} dx + \int_0^1 \frac{1}{2(x+1)} dx + \int_0^1 \frac{1}{2(1+x)^2} dx
\end{aligned}$$

Now recall integral $\log x = \frac{1}{x}$

$$\text{And, } \int x^n dx = \frac{x^{n+1}}{n+1}$$

$$\begin{aligned}
&= -\left[\frac{\log(x^2+1)}{4} \right]_0^1 + \left[\frac{\log(x+1)}{2} \right]_0^1 + \left[\frac{1}{2(x+1)} \right]_0^1 \\
&= -\left[\frac{\log((1+1))}{4} - \frac{\log((0+1))}{4} \right] + \left[\frac{\log((0+1))}{2} - \frac{\log((0+1))}{2} \right] + \left[\frac{1}{2(1+1)} - \frac{1}{2(0+1)} \right] \\
&= \frac{\log 2}{4} + \frac{1}{4}
\end{aligned}$$

Exercise 20.2

1. Question

Evaluate the following Integrals:

$$\int_2^4 \frac{x}{x^2+1} dx$$

Answer

Given definite integral is: $\int_2^4 \frac{x}{x^2+1} dx$

Let us assume $I(x) = \int_2^4 \frac{x}{x^2 + 1} dx \dots\dots(1)$

Assume $y = x^2 + 1$

Differentiating w.r.t x on both sides we get,

$$dy = d(x^2 + 1)$$

$$dy = 2x dx$$

$$x dx = \frac{dy}{2} \dots\dots(2)$$

The upper limit for Integral

$$x = 4 \Rightarrow y = 4^2 + 1$$

$$\text{Upper limit: } y = 17 \dots\dots(3)$$

The lower limit for Integral

$$x = 2 \Rightarrow y = 2^2 + 1$$

$$\text{Lower limit: } y = 5 \dots\dots(4)$$

Substituting (2),(3),(4) in the eq(1), we get,

$$\Rightarrow I(x) = \int_5^{17} \frac{1}{y} \frac{dy}{2}$$

$$\Rightarrow I(x) = \frac{1}{2} \int_5^{17} \frac{1}{y} dy$$

We know that: $\int \frac{1}{x} dx = \log(x) + c$

$$\Rightarrow I(x) = \frac{1}{2} \log y \Big|_5^{17}$$

We know that: $\int_a^b f'(x) dx = |f(x)|_b^a = f(b) - f(a)$

[here $f'(x)$ is derivative of $f(x)$]

$$\Rightarrow I(x) = \frac{1}{2} \times (\log(17) - \log(5))$$

We know that: $\log\left(\frac{x}{y}\right) = \log(x) - \log(y)$

$$\Rightarrow I(x) = \frac{1}{2} \log\left(\frac{17}{5}\right)$$

$$\therefore \int_2^4 \frac{x}{x^2 + 1} dx = \frac{1}{2} \log\left(\frac{17}{5}\right)$$

2. Question

Evaluate the following Integrals:

$$\int_1^2 \frac{1}{x(1 + \log x)^2} dx$$

Answer

Given Definite Integral can be assumed as:

$$\Rightarrow I(x) = \int_1^2 \frac{1}{x(1+\log x)^2} dx \dots\dots(1)$$

Let us assume $y = 1 + \log x$

Differentiating w.r.t x on both sides we get

$$\Rightarrow dy = d(1 + \log x)$$

$$\Rightarrow dy = \frac{1}{x} dx \dots\dots(2)$$

Lower limit for Definite Integral:

$$\Rightarrow x = 1 \Rightarrow y = 1 + \log 1$$

$$\Rightarrow y = 1 \dots\dots(3)$$

Upper limit for Definite Integral:

$$\Rightarrow x = 2 \Rightarrow y = 1 + \log 2$$

$$\Rightarrow y = 1 + \log 2 \dots\dots(4)$$

Substituting (2),(3),(4) in the eq(1) we get,

$$\Rightarrow I(x) = \int_1^{1+\log 2} \frac{1}{y^2} dy$$

$$\Rightarrow I(x) = \int_1^{1+\log 2} y^{-2} dy$$

We know that: $\int y^n dy = \frac{y^{n+1}}{n+1} + C (n \neq -1)$

$$\Rightarrow I(x) = \left[\frac{y^{-2+1}}{-2+1} \right]_1^{1+\log 2}$$

$$\Rightarrow I(x) = \left[\frac{y^{-1}}{-1} \right]_1^{1+\log 2}$$

$$\Rightarrow I(x) = -\left[\frac{1}{y} \right]_1^{1+\log 2}$$

We know that: $\int_a^b f'(x) dx = |f(x)|_b^a = f(b) - f(a)$

[here $f'(x)$ is derivative of $f(x)$)

$$\Rightarrow I(x) = \frac{-1}{1+\log 2} - \frac{-1}{1}$$

$$\Rightarrow I(x) = 1 - \frac{1}{1+\log 2}$$

$$\Rightarrow I(x) = \frac{1+\log 2 - 1}{1+\log 2}$$

$$\Rightarrow I(x) = \frac{\log 2}{1+\log 2}$$

We know that $\log e = 1$ and $\log a + \log b = \log ab$

$$\Rightarrow I(x) = \frac{\log 2}{\log 2e}$$

$$\therefore \int_1^2 \frac{1}{x(1+\log x)^2} dx = \frac{\log 2}{\log 2e}$$

3. Question

Evaluate the following Integrals:

$$\int_1^2 \frac{3x}{9x^2 - 1} dx$$

Answer

Given Definite Integral can be written as:

$$\Rightarrow I(x) = \int_1^2 \frac{3x}{9x^2 - 1} dx \dots\dots(1)$$

Let us assume $y = 9x^2 - 1$

Differentiating w.r.t x on both sides we get

$$\Rightarrow dy = d(9x^2 - 1)$$

$$\Rightarrow dy = 18x dx$$

$$\Rightarrow 3x dx = \frac{dy}{6} \dots\dots(2)$$

Upper limit for Definite Integral:

$$\Rightarrow x = 1 \Rightarrow y = (9 \times 1^2) - 1$$

$$\Rightarrow y = 8 \dots\dots(3)$$

Lower limit for Definite Integral:

$$\Rightarrow x = 2 \Rightarrow y = (9 \times 2^2) - 1$$

$$\Rightarrow y = 35 \dots\dots(4)$$

Substituting (2),(3),(4) in the eq(1), we get,

$$\Rightarrow I(x) = \int_8^{35} \frac{1}{y} \frac{dy}{6}$$

$$\Rightarrow I(x) = \frac{1}{6} \int_8^{35} \frac{1}{y} dy$$

We know that: $\int \frac{1}{x} dx = \log x + c$

$$\Rightarrow I(x) = \frac{1}{6} \log x \Big|_8^{35}$$

We know that: $\int_a^b f'(x) dx = |f(x)|_b^a = f(b) - f(a)$

[here $f'(x)$ is derivative of $f(x)$)

$$\Rightarrow I(x) = \frac{1}{6} (\log 35 - \log 8)$$

We know that: $\log(\frac{x}{y}) = \log(x) - \log(y)$

$$\Rightarrow I(x) = \frac{1}{6} \log\left(\frac{35}{8}\right)$$

$$\therefore \int_{-1}^2 \frac{3x}{9x^2 - 1} dx = \frac{1}{6} \log\left(\frac{35}{8}\right)$$

4. Question

Evaluate the following Integrals:

$$\int_0^{\pi/2} \frac{1}{5\cos x + 3\sin x} dx$$

Answer

Given Definite Integral can be written as:

$$\Rightarrow I(x) = \int_0^{\frac{\pi}{2}} \frac{1}{5\cos x + 3\sin x} dx \dots\dots(1)$$

$$\text{We know that: } \cos x = \frac{1 - \tan^2\left(\frac{x}{2}\right)}{1 + \tan^2\left(\frac{x}{2}\right)}$$

$$\text{And } \sin x = \frac{2\tan\left(\frac{x}{2}\right)}{1 + \tan^2\left(\frac{x}{2}\right)}$$

Let us find the value of $5\cos x + 3\sin x$

$$\Rightarrow 5\cos x + 3\sin x = 5 \times \left(\frac{1 - \tan^2\left(\frac{x}{2}\right)}{1 + \tan^2\left(\frac{x}{2}\right)} \right) + 3 \times \left(\frac{2\tan\left(\frac{x}{2}\right)}{1 + \tan^2\left(\frac{x}{2}\right)} \right)$$

$$\Rightarrow 5\cos x + 3\sin x = \frac{5 - 5\tan^2\left(\frac{x}{2}\right) + 6\tan\left(\frac{x}{2}\right)}{1 + \tan^2\left(\frac{x}{2}\right)}$$

$$\text{We know that: } 1 + \tan^2 x = \sec^2 x$$

$$\Rightarrow 5\cos x + 3\sin x = \frac{5 - 5\tan^2\left(\frac{x}{2}\right) + 6\tan\left(\frac{x}{2}\right)}{\sec^2\left(\frac{x}{2}\right)} \dots\dots(2)$$

Substituting (2) in (1) we get,

$$\Rightarrow I(x) = \int_0^{\frac{\pi}{2}} \frac{\sec^2\left(\frac{x}{2}\right)}{5 - 5\tan^2\left(\frac{x}{2}\right) + 6\tan\left(\frac{x}{2}\right)} dx$$

$$\text{Let us assume: } \tan\left(\frac{x}{2}\right) = t$$

Differentiating on both sides w.r.t x we get,

$$\Rightarrow \frac{1}{2} \sec^2\left(\frac{x}{2}\right) dx = dt$$

$$\Rightarrow \sec^2\left(\frac{x}{2}\right) dx = 2dt \dots\dots(3)$$

The upper limit for the Definite Integral:

$$\Rightarrow x = \frac{\pi}{2} \Rightarrow t = \tan\left(\frac{\pi}{4}\right)$$

$$\Rightarrow t = 1 \dots\dots(4)$$

The lower limit for the Definite Integral:

$$\Rightarrow x = 0 \Rightarrow t = \tan\left(\frac{0}{2}\right)$$

$$\Rightarrow t=0.....(5)$$

Substituting (3),(4),(5) in the eq(1) we get,

$$\Rightarrow I(x) = \int_0^1 \frac{2dt}{5 - 5t^2 + 6t}$$

$$\Rightarrow I(x) = \frac{2}{5} \int_0^1 \frac{dt}{1 - t^2 + \frac{6t}{5}}$$

We need to convert the denominator into standard forms

$$\Rightarrow I(x) = \frac{2}{5} \int_0^1 \frac{dt}{1 - ((t)^2 - (2 \times \frac{3}{5} \times t) + (\frac{3}{5})^2) + \frac{9}{25}}$$

$$\Rightarrow I(x) = \frac{2}{5} \int_0^1 \frac{dt}{\frac{34}{25} - (t - \frac{3}{5})^2}$$

$$\Rightarrow I(x) = \frac{2}{5} \int_0^1 \frac{dt}{\left(\sqrt{\frac{34}{25}}\right)^2 - (t - \frac{3}{5})^2}$$

$$\text{We know that: } \int \frac{dx}{a^2 - x^2} = \frac{1}{2a} \log\left(\frac{a+x}{a-x}\right)$$

$$\text{In this problem the values, } a = \sqrt{\frac{34}{25}} \text{ and } x = -\frac{3}{5}.$$

Using these values and the standard result, we get,

$$\Rightarrow I(x) = \frac{2}{5} \frac{1}{2 \times \sqrt{\frac{34}{25}}} \log\left(\frac{\sqrt{\frac{34}{25}} + t - \frac{3}{5}}{\sqrt{\frac{34}{25}} - t + \frac{3}{5}}\right) \Bigg|_0^1$$

$$\Rightarrow I(x) = \frac{1}{5} \frac{5}{\sqrt{34}} \log\left(\frac{\frac{-3 + \sqrt{34}}{5} + t}{\frac{3 + \sqrt{34}}{5} - t}\right) \Bigg|_0^1$$

$$\Rightarrow I(x) = \frac{1}{\sqrt{34}} \log\left(\frac{\frac{\sqrt{34} - 3 + 5t}{5}}{\frac{3 + \sqrt{34} - 5t}{5}}\right) \Bigg|_0^1$$

$$\Rightarrow I(x) = \frac{1}{\sqrt{34}} \log\left(\frac{\sqrt{34} - 3 + 5t}{3 + \sqrt{34} - 5t}\right) \Bigg|_0^1$$

$$\text{We know that: } \int_a^b f'(x) dx = |f(x)|_b^a = f(b) - f(a)$$

[here $f'(x)$ is derivative of $f(x)$)

$$\Rightarrow I(x) = \frac{1}{\sqrt{34}} \times \left(\log\left(\frac{\sqrt{34} - 3 + 5}{\sqrt{34} + 3 - 5}\right) - \log\left(\frac{\sqrt{34} - 3 + 0}{\sqrt{34} + 3 - 0}\right) \right)$$

$$\Rightarrow I(x) = \frac{1}{\sqrt{34}} \times \left(\log \left(\frac{\sqrt{34} + 2}{\sqrt{34} - 2} \right) - \log \left(\frac{\sqrt{34} - 3}{\sqrt{34} + 3} \right) \right)$$

We know that: $\log(\frac{x}{y}) = \log(x) - \log(y)$

$$\Rightarrow I(x) = \frac{1}{\sqrt{34}} \times \left(\log \left(\frac{\frac{\sqrt{34} + 2}{\sqrt{34} - 2}}{\frac{\sqrt{34} - 3}{\sqrt{34} + 3}} \right) \right)$$

$$\Rightarrow I(x) = \frac{1}{\sqrt{34}} \times \log \left(\frac{(\sqrt{34} + 2) \times (\sqrt{34} + 3)}{(\sqrt{34} - 2) \times (\sqrt{34} - 3)} \right)$$

$$\Rightarrow I(x) = \frac{1}{\sqrt{34}} \times \log \left(\frac{40 + 5\sqrt{34}}{40 - 5\sqrt{34}} \right)$$

$$\Rightarrow I(x) = \frac{1}{\sqrt{34}} \log \left(\frac{5 \times (8 + \sqrt{34})}{5 \times (8 - \sqrt{34})} \right)$$

$$\Rightarrow I(x) = \frac{1}{\sqrt{34}} \log \left(\frac{8 + \sqrt{34}}{8 - \sqrt{34}} \right)$$

$$\therefore \int_0^{\frac{\pi}{2}} \frac{1}{5\cos x + 3\sin x} dx = \frac{1}{\sqrt{34}} \log \left(\frac{8 + \sqrt{34}}{8 - \sqrt{34}} \right)$$

5. Question

Evaluate the following Integrals:

$$\int_0^a \frac{x}{\sqrt{a^2 + x^2}} dx$$

Answer

Given Definite integral can be written as:

$$\Rightarrow I(x) = \int_0^a \frac{x}{\sqrt{a^2 + x^2}} dx \quad (1)$$

Let us assume $y = a^2 + x^2$

Differentiating w.r.t x on both sides we get,

$$\Rightarrow dy = d(a^2 + x^2)$$

$$\Rightarrow dy = 2x dx$$

$$\Rightarrow x dx = \frac{dy}{2} \quad \dots \dots (2)$$

Upper limit for the Definite Integral:

$$\Rightarrow x=a \Rightarrow y = a^2 + a^2$$

$$\Rightarrow y=2a^2 \quad \dots \dots (3)$$

Lower limit for the Definite Integral:

$$\Rightarrow x=0 \Rightarrow y = a^2 + 0^2$$

$$\Rightarrow y = a^2 \quad \dots \dots (4)$$

Substituting (2),(3),(4) in the eq(1), we get,

$$\Rightarrow I(x) = \int_{a^2}^{2a^2} \frac{dy}{2\sqrt{y}}$$

$$\Rightarrow I(x) = \frac{1}{2} \int_{a^2}^{2a^2} y^{-\frac{1}{2}} dy$$

We know that: $\int y^n dy = \frac{y^{n+1}}{n+1} + C (n \neq -1)$

$$\Rightarrow I(x) = \frac{1}{2} \left[\frac{y^{-\frac{1}{2}+1}}{-\frac{1}{2}+1} \right]_{a^2}^{2a^2}$$

$$\Rightarrow I(x) = \frac{1}{2} \left[\frac{y^{\frac{1}{2}}}{\frac{1}{2}} \right]_{a^2}^{2a^2}$$

$$\Rightarrow I(x) = y^{\frac{1}{2}} \Big|_{a^2}^{2a^2}$$

We know that: $\int_a^b f'(x) dx = |f(x)|_a^b = f(b) - f(a)$

[here $f'(x)$ is derivative of $f(x)$)

$$\Rightarrow I(x) = (2a^2)^{1/2} - (a^2)^{1/2}$$

$$\Rightarrow I(x) = \sqrt{2}a - a$$

$$\Rightarrow I(x) = a(\sqrt{2}-1)$$

$$\therefore \int_0^a \frac{x}{\sqrt{a^2+x^2}} dx = a(\sqrt{2}-1)$$

6. Question

Evaluate the following Integrals:

$$\int_0^1 \frac{e^x}{1+e^{2x}} dx$$

Answer

Given Definite Integral can be written as:

$$\Rightarrow I(x) = \int_0^1 \frac{e^x}{1+e^{2x}} dx \dots\dots(1)$$

Let us assume $y = e^x$

Differentiating w.r.t x on both sides we get,

$$\Rightarrow dy = e^x dx$$

$$\Rightarrow dy = e^x dx \dots\dots(2)$$

Upper limit for the Definite Integral:

$$\Rightarrow x = 1 \Rightarrow y = e^1$$

$$\Rightarrow y = e(3)$$

Lower limit for the Definite Integral:

$$\Rightarrow x = 0 \Rightarrow y = e^0$$

$$\Rightarrow y = 1 \quad (4)$$

Substituting (2), (3), (4) in the eq(1) we get,

$$\Rightarrow I(x) = \int_1^e \frac{dy}{1+y^2}$$

We know that: $\int \frac{1}{1+x^2} dx = \tan^{-1}x + C$

$$\Rightarrow I(x) = \tan^{-1}(x)|_1^e$$

We know that: $\int_a^b f'(x) dx = |f(x)|_a^b = f(b) - f(a)$

[here $f'(x)$ is derivative of $f(x)$)

$$\Rightarrow I(x) = \tan^{-1}(e) - \tan^{-1}(1)$$

$$\Rightarrow I(x) = \tan^{-1}(e) - \frac{\pi}{4}$$

$$\therefore \int_0^1 \frac{e^x}{1+e^{2x}} dx = \tan^{-1}e - \frac{\pi}{4}$$

7. Question

Evaluate the following Integrals:

$$\int_0^1 x e^{x^2} dx$$

Answer

Given Definite Integral can be written as:

$$\Rightarrow I(x) = \int_0^1 x e^{x^2} dx \quad (1)$$

Let us assume $y = x^2$

Differentiating w.r.t x on both sides we get,

$$\Rightarrow dy = d(x^2)$$

$$\Rightarrow dy = 2x dx$$

$$\Rightarrow x dx = \frac{dy}{2} \dots\dots (2)$$

Upper limit for the Definite Integral:

$$\Rightarrow x = 1 \Rightarrow y = 1^2$$

$$\Rightarrow y = 1 \dots\dots (3)$$

Lower limit for the Definite Integral:

$$\Rightarrow x = 0 \Rightarrow y = 0^2$$

$$\Rightarrow y = 0 \dots\dots (4)$$

Substituting (2), (3), (4) in the eq(1), we get,

$$\Rightarrow I(x) = \int_0^1 \frac{e^y dy}{2}$$

$$\Rightarrow I(x) = \frac{1}{2} \int_0^1 e^y dy$$

We know that: $\int e^x dx = e^x + C$

$$\Rightarrow I(x) = \frac{1}{2} e^y \Big|_0^1$$

We know that: $\int_a^b f'(x) dx = |f(x)|_a^b = f(b) - f(a)$

[here $f'(x)$ is derivative of $f(x)$)

$$\Rightarrow I(x) = \frac{1}{2} (e - e^0)$$

$$\Rightarrow I(x) = \frac{1}{2} (e - 1)$$

$$\therefore \int_0^1 x e^{x^2} dx = \frac{1}{2} (e - 1)$$

8. Question

Evaluate the following Integrals:

$$\int_1^3 \frac{\cos(\log x)}{x} dx$$

Answer

Given Definite Integral can be written as:

$$\Rightarrow I(x) = \int_1^3 \frac{\cos(\log x)}{x} dx \dots\dots(1)$$

Let us assume $y = \log x$

Differentiating w.r.t x on both sides

$$\Rightarrow d(y) = d(\log x)$$

$$\Rightarrow dy = \frac{1}{x} dx \dots\dots(2)$$

Upper limit for the Definite Integral:

$$\Rightarrow x = 3 \Rightarrow y = \log(3)$$

$$\Rightarrow y = \log 3 \dots\dots(3)$$

Lower limit for the Definite Integral:

$$\Rightarrow x = 1 \Rightarrow y = \log(1)$$

$$\Rightarrow y = 0 \dots\dots(4)$$

Substituting (2),(3),(4) in the eq(1) we get,

$$\Rightarrow I(x) = \int_0^{\log 3} \cos y dy$$

We know that $\int \cos x dx = \sin x + C$

$$\Rightarrow I(x) = \sin x \Big|_0^{\log 3}$$

We know that: $\int_a^b f'(x) dx = |f(x)|_a^b = f(b) - f(a)$

here $f'(x)$ is derivative of $f(x)$)

$$\Rightarrow I(x) = \sin(\log 3) - \sin(0)$$

$$\Rightarrow I(x) = \sin(\log 3) - 0$$

$$\Rightarrow I(x) = \sin(\log 3)$$

$$\therefore \int_1^3 \frac{\cos(\log x)}{x} dx = \sin(\log 3)$$

9. Question

Evaluate the following Integrals:

$$\int_0^1 \frac{2x}{1+x^4} dx$$

Answer

Given Definite Integral can be written as:

$$\Rightarrow I(x) = \int_0^1 \frac{2x}{1+x^4} dx \dots\dots(1)$$

Let us assume $y = x^2$

Differentiating w.r.t x on both sides we get,

$$\Rightarrow dy = d(x^2)$$

$$\Rightarrow dy = 2x dx \dots\dots(2)$$

Lower limit for the Definite Integral:

$$\Rightarrow x = 0 \Rightarrow y = 0^2$$

$$\Rightarrow y = 0 \dots\dots(3)$$

Upper limit for the Definite Integral:

$$\Rightarrow x = 1 \Rightarrow y = 1^2$$

$$\Rightarrow y = 1 \dots\dots(4)$$

Substituting (2),(3),(4) in the eq(1), we get,

$$\Rightarrow I(x) = \int_0^1 \frac{dy}{1+y^2}$$

We know that: $\int \frac{1}{1+x^2} dx = \tan^{-1}(x) + C$

$$\Rightarrow I(x) = \tan^{-1}(y)|_0^1$$

We know that $\int_a^b f(x) dx = [f(x)]_a^b = f(b) - f(a)$

[here $f'(x)$ is derivative of $f(x)$)

$$\Rightarrow I(x) = \tan^{-1}(1) - \tan^{-1}(0)$$

$$\Rightarrow I(x) = \frac{\pi}{4} - 0$$

$$\Rightarrow I(x) = \frac{\pi}{4}$$

$$\therefore \int_0^1 \frac{2x}{1+x^4} dx = \frac{\pi}{4}$$

10. Question

Evaluate the following Integrals:

$$\int_0^a \sqrt{a^2 - x^2} dx$$

Answer

Given Definite Integral can be written as:

$$\Rightarrow I(x) = \int_0^a \sqrt{a^2 - x^2} dx \dots\dots (1)$$

Let us assume $x = a \sin\theta$

Differentiating w.r.t x on both sides we get,

$$\Rightarrow dx = d(a \sin \theta)$$

$$\Rightarrow dx = a \cos \theta d\theta \dots\dots (2)$$

Let us find the value of $\sqrt{a^2 - x^2}$

$$\Rightarrow \sqrt{a^2 - x^2} = \sqrt{a^2 - (a \sin \theta)^2}$$

$$\Rightarrow \sqrt{a^2 - x^2} = \sqrt{(a^2 \times (1 - \sin^2 \theta))}$$

$$\Rightarrow \sqrt{a^2 - x^2} = a \times \sqrt{\cos^2 \theta}$$

$$(\because 1 - \sin^2 \theta = \cos^2 \theta)$$

$$\Rightarrow \sqrt{a^2 - x^2} = a \cos \theta \dots\dots (3)$$

Lower limit for the Definite Integral:

$$\Rightarrow x = 0 \Rightarrow \theta = \sin^{-1}\left(\frac{0}{a}\right)$$

$$\Rightarrow \theta = \sin^{-1}(0)$$

$$\Rightarrow \theta = 0 \dots\dots (4)$$

Upper limit for the Definite Integral:

$$\Rightarrow x = a \Rightarrow \theta = \sin^{-1}\left(\frac{a}{a}\right)$$

$$\Rightarrow \theta = \sin^{-1}(1)$$

$$\Rightarrow \theta = \frac{\pi}{2} \dots\dots (5)$$

Substituting (2),(3),(4),(5) in eq(1) we get,

$$\Rightarrow I(x) = \int_0^{\frac{\pi}{2}} a \cos \theta \times a \cos \theta d\theta$$

$$\Rightarrow I(x) = a^2 \int_0^{\frac{\pi}{2}} \cos^2 \theta d\theta$$

We know that $\cos 2\theta = 2\cos^2 \theta - 1$

Then

$$\cos^2 \theta = \frac{1 + \cos 2\theta}{2}$$

Using these result for the integration, we get,

$$\Rightarrow I(x) = a^2 \int_0^{\frac{\pi}{2}} \frac{1 + \cos 2\theta}{2} d\theta$$

$$\Rightarrow I(x) = \frac{a^2}{2} \int_0^{\frac{\pi}{2}} 1 + \cos 2\theta d\theta$$

We know that:

$\int adx = ax + c$ and also

$$\int \cos ax = \frac{-\sin ax}{a} + c.$$

We know that:

$$\int_a^b f'(x) dx = |f(x)|_a^b = f(b) - f(a)$$

[here $f'(x)$ is derivative of $f(x)$].

$$\Rightarrow I(x) = \frac{a^2}{2} \times \left(\theta - \frac{\sin 2\theta}{2} \right) \Big|_0^{\frac{\pi}{2}}$$

$$\Rightarrow I(x) = \frac{a^2}{2} \times \left(\left(\frac{\pi}{2} - \frac{\sin(\frac{2\pi}{2})}{2} \right) - \left(0 - \frac{\sin(2 \times 0)}{2} \right) \right)$$

$$\text{We know that } \sin n\pi = 0 \quad (n \in \mathbb{N}) \Rightarrow I(x) = \frac{a^2}{2} \times \left(\left(\frac{\pi}{2} - 0 \right) - (0 - 0) \right)$$

$$\Rightarrow I(x) = \frac{a^2}{2} \times \frac{\pi}{2}$$

$$\Rightarrow I(x) = \frac{a^2 \pi}{4}$$

$$\therefore \int_0^a \sqrt{a^2 - x^2} dx = \frac{\pi a^2}{4}$$

11. Question

Evaluate the following Integrals:

$$\int_0^{\pi/2} \sqrt{\sin \phi} \cos^5 \phi d\phi$$

Answer

Given Definite Integral can be written as:

$$\Rightarrow I(x) = \int_0^{\frac{\pi}{2}} \sqrt{\sin \phi} \cos^5 \phi d\phi$$

$$\Rightarrow I(x) = \int_0^{\frac{\pi}{2}} \sqrt{\sin\phi} \cos^4 \phi \cos\theta d\phi$$

Let us assume $\sin\phi = t$,

Differentiating w.r.t ϕ on both sides we get,

$$\Rightarrow dt = \cos\phi d\phi \dots\dots(2)$$

Upper limit for the Definite Integral:

$$\Rightarrow \phi = \frac{\pi}{2} \Rightarrow t = \sin\left(\frac{\pi}{2}\right)$$

$$\Rightarrow t = 1 \dots\dots(3)$$

Lower limit for the Definite Integral:

$$\Rightarrow \phi=0 \Rightarrow t = \sin(0)$$

$$\Rightarrow t = 0 \dots\dots(4)$$

We know that $\cos^2\phi = 1 - \sin^2\phi$

$$\Rightarrow \cos^2\phi = 1 - t^2 \dots\dots(5)$$

Substituting (2),(3),(4),(5) in the eq(1), we get,

$$\Rightarrow I(x) = \int_0^1 \sqrt{t}(1-t^2)^2 dt$$

$$\Rightarrow I(x) = \int_0^1 t^{\frac{1}{2}} \times (1-2t^2+t^4) dt$$

$$\Rightarrow I(x) = \int_0^1 t^{\frac{1}{2}} - 2t^{\frac{5}{2}} + t^{\frac{9}{2}} dt$$

We know that:

$$\int x^n dx = \frac{x^{n+1}}{n+1} + C (n \neq -1)$$

We know that:

$$\int_a^b f(x) dx = |f(x)|_a^b = f(b) - f(a)$$

[here $f'(x)$ is derivative of $f(x)$]

$$\Rightarrow I(x) = \frac{\frac{1}{2}+1}{\frac{1}{2}+1} \left[t^{\frac{5}{2}+1} \right]_0^1 - 2 \times \left(\frac{\frac{5}{2}+1}{\frac{5}{2}+1} \right) \left[t^{\frac{9}{2}+1} \right]_0^1$$

$$\Rightarrow I(x) = \frac{\frac{3}{2}}{\frac{3}{2}} \left[t^{\frac{7}{2}} \right]_0^1 - 2 \times \left(\frac{\frac{7}{2}}{\frac{7}{2}} \right) \left[t^{\frac{11}{2}} \right]_0^1$$

$$\Rightarrow I(x) = \left(\frac{2}{3} \times \left(1^{\frac{3}{2}} - 0^{\frac{3}{2}} \right) \right) - \left(2 \times \frac{2}{7} \times \left(1^{\frac{7}{2}} - 0^{\frac{7}{2}} \right) \right) + \left(\frac{2}{11} \times \left(1^{\frac{11}{2}} - 0^{\frac{11}{2}} \right) \right)$$

$$\Rightarrow I(x) = \left(\frac{2}{3} \times 1 \right) - \left(\frac{4}{7} \times 1 \right) + \left(\frac{2}{11} \times 1 \right)$$

$$\Rightarrow I(x) = \frac{2}{3} - \frac{4}{7} + \frac{2}{11}$$

$$\Rightarrow I(x) = \frac{64}{231}$$

$$\therefore \int_0^{\frac{\pi}{2}} \sqrt{\sin \phi} \cos^5 \phi d\phi = \frac{64}{231}$$

12. Question

Evaluate the following Integrals:

$$\int_0^{\pi/2} \frac{\cos x}{1 + \sin^2 x} dx$$

Answer

Given Definite Integral can be written as:

$$\Rightarrow I(x) = \int_0^{\frac{\pi}{2}} \frac{\cos x}{1 + \sin^2 x} dx \dots \dots (1)$$

Let us assume $y = \sin x$,

Differentiating on both sides w.r.t x we get,

$$\Rightarrow dy = d(\sin x)$$

$$\Rightarrow dy = \cos x dx \dots \dots (2)$$

Upper limit for the Definite Integral:

$$x = \frac{\pi}{2}, y = \sin \frac{\pi}{2} = 1$$

$$\Rightarrow y = 1 \dots \dots (3)$$

Lower limit for the Definite Integral:

$$\Rightarrow x = 0 \Rightarrow y = \sin(0)$$

$$\Rightarrow y = 0 \dots \dots (4)$$

Substituting (2),(3),(4) in the eq(1) we get,

$$\Rightarrow I(x) = \int_0^1 \frac{dt}{1 + t^2}$$

We know that:

$$\int \frac{1}{1 + x^2} dx = \tan^{-1} x + c$$

We know that:

$$\int_a^b f(x) dx = |f(x)|_a^b = f(b) - f(a)$$

[here $f'(x)$ is derivative of $f(x)$]

$$\Rightarrow I(x) = \tan^{-1} t|_0^1$$

$$\Rightarrow I(x) = (\tan^{-1}(1) - \tan^{-1}(0))$$

$$\Rightarrow I(x) = \frac{\pi}{4} - 0$$

$$\Rightarrow I(x) = \frac{\pi}{4}$$

$$\therefore \int_0^{\frac{\pi}{2}} \frac{\cos x}{1 + \sin^2 x} dx = \frac{\pi}{4}$$

13. Question

Evaluate the following Integrals:

$$\int_0^{\pi/2} \frac{\sin \theta}{\sqrt{1 + \cos \theta}} d\theta$$

Answer

Given Definite Integral can be written as:

$$\Rightarrow I(x) = \int_0^{\frac{\pi}{2}} \frac{\sin \theta}{\sqrt{1 + \cos \theta}} d\theta \dots\dots (1)$$

Let us assume $1 + \cos \theta = y$

Differentiating w.r.t θ on both sides we get,

$$\Rightarrow d(1 + \cos \theta) = d(y)$$

$$\Rightarrow -\sin \theta d\theta = dy$$

$$\Rightarrow \sin \theta d\theta = -dy \dots\dots (2)$$

Upper limit for the Definite Integral

$$\Rightarrow \theta = \frac{\pi}{2} \Rightarrow y = 1 + \cos\left(\frac{\pi}{2}\right)$$

$$\Rightarrow y = 1 \dots\dots (3)$$

Lower limit for the Definite Integral:

$$\Rightarrow \theta = 0 \Rightarrow y = 1 + \cos(0)$$

$$\Rightarrow y = 2$$

$$\Rightarrow y = 2 \dots\dots (4)$$

Substituting (2),(3),(4) in the eq(1), we get,

$$\Rightarrow I(x) = \int_2^1 -\frac{dy}{\sqrt{y}}$$

We know that:

$$\int_a^b f(x) dx = - \int_b^a f(x) dx$$

$$\Rightarrow I(x) = \int_1^2 y^{-\frac{1}{2}} dy$$

We know that:

$$\int x^n dx = \frac{x^{n+1}}{n+1} + C$$

We know that:

$$\int_a^b f'(x) dx = |f(x)|_a^b = f(b) - f(a)$$

[here $f'(x)$ is derivative of $f(x)$]

$$\Rightarrow I(x) = \left. \frac{y^{-\frac{1}{2}+1}}{-\frac{1}{2}+1} \right|_1^2$$

$$\Rightarrow I(x) = \left. \frac{y^{\frac{1}{2}}}{\frac{1}{2}} \right|_1^2$$

$$\Rightarrow I(x) = 2 \times \left(2^{\frac{1}{2}} - 1^{\frac{1}{2}} \right)$$

$$\Rightarrow I(x) = 2 \times (\sqrt{2} - \sqrt{1})$$

$$\Rightarrow I(x) = 2(\sqrt{2} - 1)$$

$$\therefore \int_0^{\frac{\pi}{2}} \frac{\sin \theta}{\sqrt{1 + \cos \theta}} d\theta = 2(\sqrt{2} - 1)$$

14. Question

Evaluate the following Integrals:

$$\int_0^{\pi/3} \frac{\cos x}{3 + 4 \sin x} dx$$

Answer

Given Definite Integral can be written as:

$$\Rightarrow I(x) = \int_0^{\frac{\pi}{3}} \frac{\cos x}{3 + 4 \sin x} dx \dots \dots (1)$$

Let us assume $3 + 4 \sin x = y$

Differentiating w.r.t x on both sides we get,

$$\Rightarrow d(3 + 4 \sin x) = d(y)$$

$$\Rightarrow 4 \cos x dx = dy$$

$$\Rightarrow \cos x dx = \frac{dy}{4} \dots \dots (2)$$

Lower limit for the Definite Integral:

$$\Rightarrow x = 0 \Rightarrow y = 3 + 4 \sin(0)$$

$$\Rightarrow y = 3 + 0$$

$$\Rightarrow y = 3 \dots \dots (3)$$

Upper limit for the Definite Integral:

$$\Rightarrow x = \frac{\pi}{3} \Rightarrow y = 3 + 4 \sin\left(\frac{\pi}{3}\right)$$

$$\Rightarrow y = 3 + (4 \times \frac{\sqrt{3}}{2})$$

$$\Rightarrow y = 3 + 2\sqrt{3} \dots \dots (4)$$

Substituting (2),(3),(4) in the eq(1) we get,

$$\Rightarrow I(x) = \int_3^{3+2\sqrt{3}} \frac{dy}{4y}$$

$$\Rightarrow I(x) = \frac{1}{4} \int_3^{3+2\sqrt{3}} \frac{dy}{y}$$

We know that:

$$\int \frac{dx}{x} = \log x + C$$

We know that:

$$\int_a^b f(x) dx = |f(x)|_a^b = f(b) - f(a)$$

[here $f'(x)$ is derivative of $f(x)$)

$$\Rightarrow I(x) = \frac{1}{4} \times \log y |_3^{3+2\sqrt{3}}$$

$$\Rightarrow I(x) = \frac{1}{4} \times (\log(3 + 2\sqrt{3}) - \log(3))$$

$$\text{We know that } \log\left(\frac{a}{b}\right) = \log a - \log b$$

$$\Rightarrow I(x) = \frac{1}{4} \times \log\left(\frac{3 + 2\sqrt{3}}{3}\right)$$

$$\therefore \int_0^{\frac{\pi}{3}} \frac{\cos x}{3 + 4\sin x} dx = \frac{1}{4} \log\left(\frac{3 + 2\sqrt{3}}{3}\right)$$

15. Question

Evaluate the following Integrals:

$$\int_0^1 \frac{\sqrt{\tan^{-1} x}}{1+x^2} dx$$

Answer

Given Definite Integral can be written as:

$$\Rightarrow I(x) = \int_0^1 \frac{\sqrt{\tan^{-1} x}}{1+x^2} dx \dots \dots (1)$$

Let us assume $\tan^{-1} x = y$

Differentiating w.r.t x on both sides we get,

$$\Rightarrow d(\tan^{-1} x) = d(y)$$

$$\Rightarrow \frac{1}{1+x^2} dx = dy \dots \dots (2)$$

Upper limit of the Definite Integral:

$$\Rightarrow x = 1 \Rightarrow y = \tan^{-1}(1)$$

$$\Rightarrow y = \frac{\pi}{4} \dots \dots (3)$$

Lower limit of the Definite Integral:

$$\Rightarrow x = 0 \Rightarrow y = \tan^{-1}(0)$$

$$\Rightarrow y = 0 \dots \dots (4)$$

Substituting (2),(3),(4) in the eq(1) we get,

$$\Rightarrow I(x) = \int_0^{\frac{\pi}{4}} \sqrt{t} dt$$

$$\Rightarrow I(x) = \int_0^{\frac{\pi}{4}} t^{\frac{1}{2}} dt$$

We know that:

$$\int x^n dx = \frac{x^{n+1}}{n+1} + C$$

We know that:

$$\int_a^b f'(x) dx = |f(x)|_a^b = f(b) - f(a)$$

[here $f'(x)$ is derivative of $f(x)$)

$$\Rightarrow I(x) = \left[\frac{t^{\frac{1}{2}+1}}{\frac{1}{2}+1} \right]_0^{\frac{\pi}{4}}$$

$$\Rightarrow I(x) = \left[\frac{t^{\frac{3}{2}}}{\frac{3}{2}} \right]_0^{\frac{\pi}{4}}$$

$$\Rightarrow I(x) = \frac{2}{3} \times \left(\left(\frac{\pi}{4} \right)^{\frac{3}{2}} - 0^{\frac{3}{2}} \right)$$

$$\Rightarrow I(x) = \frac{2}{3} \times \left(\frac{\frac{3}{2}}{4^{\frac{3}{2}}} \right)$$

$$\Rightarrow I(x) = \frac{2}{3} \times \left(\frac{\frac{3}{2}}{8} \right)$$

$$\Rightarrow I(x) = \frac{\frac{3}{2}}{12}$$

$$\therefore \int_0^1 \frac{\sqrt{\tan^{-1} x}}{1+x^2} dx = \frac{\frac{3}{2}}{12}$$

16. Question

Evaluate the following Integrals:

$$\int_0^2 x\sqrt{x+2} dx$$

Answer

Given Definite Integral can be written as:

$$\Rightarrow I(x) = \int_0^2 x\sqrt{x+2} dx \dots\dots (1)$$

Let us assume $x+2 = y$

Then, $x = y-2 \dots\dots (2)$

Differentiating on both side w.r.t x we get,

$$\Rightarrow d(x+2) = d(y)$$

$$\Rightarrow dx = dy \dots\dots (3)$$

Upper limit for the Definite Integral:

$$\Rightarrow x = 2 \Rightarrow y = 2+2$$

$$\Rightarrow y = 4 \dots\dots (4)$$

Lower limit for the Definite Integral:

$$\Rightarrow x = 0 \Rightarrow y = 0+2$$

$$\Rightarrow y = 2 \dots\dots (5)$$

Substituting (2),(3),(4),(5) in the eq(1) we get,

$$\Rightarrow I(x) = \int_2^4 (y-2)y^{\frac{1}{2}} dy$$

$$\Rightarrow I(x) = \int_2^4 (y^{\frac{3}{2}} - 2y^{\frac{1}{2}}) dy$$

We know that:

$$\int x^n dx = \frac{x^{n+1}}{n+1} + C$$

We know that:

$$\int_a^b f(x) dx = |f(x)|_a^b = f(b) - f(a)$$

[here $f'(x)$ is derivative of $f(x)$]

$$\Rightarrow I(x) = \frac{y^{\frac{3}{2}+1}}{\frac{3}{2}+1} \Big|_2^4 - 2 \times \left(\frac{y^{\frac{1}{2}+1}}{\frac{1}{2}+1} \right) \Big|_2^4$$

$$\Rightarrow I(x) = \frac{y^{\frac{5}{2}}}{\frac{5}{2}} \Big|_2^4 - 2 \times \left(\frac{y^{\frac{3}{2}}}{\frac{3}{2}} \right) \Big|_2^4$$

$$\Rightarrow I(x) = \frac{2}{5}y^{\frac{5}{2}} \Big|_2^4 - 2 \times \frac{2}{3}y^{\frac{3}{2}} \Big|_2^4$$

$$\Rightarrow I(x) = \left(\frac{2}{5} \times \left(4^{\frac{5}{2}} - 2^{\frac{5}{2}} \right) \right) - \left(\frac{4}{3} \times \left(4^{\frac{3}{2}} - 2^{\frac{3}{2}} \right) \right)$$

$$\Rightarrow I(x) = \left(\frac{2}{5} \times (32 - 4\sqrt{2}) \right) - \left(\frac{4}{3} \times (8 - 2\sqrt{2}) \right)$$

$$\Rightarrow I(x) = \frac{64}{5} - \frac{8\sqrt{2}}{5} - \frac{32}{3} + \frac{8\sqrt{2}}{3}$$

$$\Rightarrow I(x) = \frac{32}{15} + \frac{16\sqrt{2}}{15}$$

$$\Rightarrow I(x) = \frac{1}{15} \times (16\sqrt{2} + 32)$$

$$\Rightarrow I(x) = \frac{16\sqrt{2}(\sqrt{2} + 1)}{15}$$

$$\therefore \int_0^2 x\sqrt{x+2} dx = \frac{16\sqrt{2}(\sqrt{2} + 1)}{15}$$

17. Question

Evaluate the following Integrals:

$$\int_0^1 \tan^{-1} \left(\frac{2x}{1-x^2} \right) dx$$

Answer

Given Definite Integral can be written as:

$$\Rightarrow I(x) = \int_0^1 \tan^{-1} \left(\frac{2x}{1-x^2} \right) dx \dots \dots (1)$$

Let us assume $x = \tan y$

Differentiating w.r.t x on both sides we get,

$$\Rightarrow d(x) = d(\tan y)$$

$$\Rightarrow dx = \sec^2 y dy \dots \dots (2)$$

Then

$$\frac{2x}{1-x^2} = \frac{2\tan y}{1-\tan^2 y}$$

We know that:

$$\tan 2\theta = \frac{2\tan\theta}{1-\tan^2\theta}$$

Now,

$$\frac{2x}{1-x^2} = \tan 2y \dots \dots (3)$$

Upper limit for the Definite Integral:

$$\Rightarrow x = 1 \Rightarrow y = \tan^{-1}(1)$$

$$\Rightarrow y = \frac{\pi}{4} \dots \dots (4)$$

Lower limit for the Definite Integral:

$$\Rightarrow x = 0 \Rightarrow y = \tan^{-1}(0)$$

$$\Rightarrow y = 0 \dots\dots (5)$$

Substituting (2),(3),(4),(5) in (1) we get,

$$\Rightarrow I(x) = \int_0^{\frac{\pi}{4}} \tan^{-1}(\tan 2y) \sec^2 y dy$$

$$\Rightarrow I(x) = \int_0^{\frac{\pi}{4}} 2y \sec^2 y dy$$

$$\Rightarrow I(x) = 2 \int_0^{\frac{\pi}{4}} y \sec^2 y dy$$

We know that the By-partss integration is:

$$\Rightarrow \int UV dx = U \int V dx - \int \left(\frac{d}{dx}(U) \int V dx \right) dx$$

Now applying by parts Integration:

$$\Rightarrow I(x) = 2 \times \left(y \int_0^{\frac{\pi}{4}} \sec^2 y dy - \int_0^{\frac{\pi}{4}} \left(\frac{d}{dy}(y) \int \sec^2 y dy \right) dy \right)$$

We know that: $\int \sec^2 x dx = \tan x + C$

We know that:

$$\int_a^b f(x) dx = |f(x)|_a^b = f(b) - f(a)$$

[here $f'(x)$ is derivative of $f(x)$)

$$\Rightarrow I(x) = 2 \times \left((yt \tan y) \Big|_0^{\frac{\pi}{4}} - \int_0^{\frac{\pi}{4}} t \tan y dy \right)$$

We know that: $\int \tan x dx = -\log(\cos x) + C$

$$\Rightarrow I(x) = 2 \times \left((yt \tan y) \Big|_0^{\frac{\pi}{4}} - (-\log(\cos x)) \Big|_0^{\frac{\pi}{4}} \right)$$

$$\Rightarrow I(x) = 2 \times \left((yt \tan y) \Big|_0^{\frac{\pi}{4}} + (\log(\cos x)) \Big|_0^{\frac{\pi}{4}} \right)$$

$$\Rightarrow I(x) = 2 \times \left(\left(\frac{\pi}{4} \tan \left(\frac{\pi}{4} \right) - 0 \tan \left(\frac{0}{4} \right) \right) + \left(\log \left(\cos \left(\frac{\pi}{4} \right) \right) - \log \left(\cos \left(\frac{0}{4} \right) \right) \right) \right)$$

$$\Rightarrow I(x) = 2 \times \left(\frac{\pi}{4} - 0 + \log \left(\frac{1}{\sqrt{2}} \right) - \log(1) \right)$$

$$\Rightarrow I(x) = 2 \times \left(\frac{\pi}{4} + \log \left(2^{-\frac{1}{2}} \right) - 0 \right)$$

We know that: $\log(a^b) = b \log a$

$$\Rightarrow I(x) = 2 \times \left(\frac{\pi}{4} - \frac{1}{2} \log 2 \right)$$

$$\Rightarrow I(x) = \frac{\pi}{2} - \log 2$$

$$\therefore \int_0^1 \tan^{-1} \left(\frac{2x}{1-x^2} \right) dx = \frac{\pi}{2} - \log 2.$$

18. Question

Evaluate the following Integrals:

$$\int_0^{\pi/2} \frac{\sin x \cos x}{1 + \sin^4 x} dx$$

Answer

Given Definite Integral can be written as:

$$\Rightarrow I(x) = \int_0^{\frac{\pi}{2}} \frac{\sin x \cos x}{1 + \sin^4 x} dx \dots \dots (1)$$

Let us assume, $y = \sin^2 x$

Differentiating w.r.t x on both sides we get,

$$\Rightarrow dy = d(\sin^2 x)$$

$$\Rightarrow dy = 2\sin x \cos x dx$$

$$\Rightarrow \sin x \cos x dx = \frac{dy}{2} \dots \dots (2)$$

Upper limit for the Definite Integral:

$$\Rightarrow x = \frac{\pi}{2} \Rightarrow y = \sin^2 \frac{\pi}{2}$$

$$\Rightarrow y = 1 \dots \dots (3)$$

Lower limit for the Definite Integral:

$$\Rightarrow x = 0 \Rightarrow y = \sin^2 0$$

$$\Rightarrow y = 0 \dots \dots (4)$$

Substituting (2),(3),(4) in the eq(1) we get,

$$\Rightarrow I(x) = \int_0^1 \frac{dy}{2(1+y^2)}$$

$$\Rightarrow I(x) = \frac{1}{2} \int_0^1 \frac{dy}{1+y^2}$$

We know that:

$$\int \frac{1}{1+x^2} dx = \tan^{-1} x + C$$

We know that:

$$\int_a^b f'(x) dx = |f(x)|_a^b = f(b) - f(a)$$

[here $f'(x)$ is derivative of $f(x)$)

$$\Rightarrow I(x) = \frac{1}{2} \times (\tan^{-1} y)|_0^1$$

$$\Rightarrow I(x) = \frac{1}{2} \times (\tan - 11 - \tan - 10)$$

$$\Rightarrow I(x) = \frac{1}{2} \times \left(\frac{\pi}{4}\right)$$

$$\Rightarrow I(x) = \frac{\pi}{8}$$

$$\therefore \int_0^{\frac{\pi}{2}} \frac{\sin x \cos x}{1 + \sin^4 x} dx = \frac{\pi}{8}$$

19. Question

Evaluate the following Integrals:

$$\int_0^{\pi/2} \frac{dx}{a \cos x + b \sin x} \quad a, b > 0$$

Answer

Given Definite Integral can be written as:

$$\Rightarrow I(x) = \int_0^{\frac{\pi}{2}} \frac{dx}{a \cos x + b \sin x} \quad a, b > 0 \dots \dots (1)$$

We know that:

$$\cos x = \frac{1 - \tan^2\left(\frac{x}{2}\right)}{1 + \tan^2\left(\frac{x}{2}\right)} \text{ and}$$

$$\sin x = \frac{2 \tan\left(\frac{x}{2}\right)}{1 + \tan^2\left(\frac{x}{2}\right)}$$

Substituting these value in (1) we get,

$$\Rightarrow I(x) = \int_0^{\frac{\pi}{2}} \frac{dx}{a \left(\frac{1 - \tan^2\left(\frac{x}{2}\right)}{1 + \tan^2\left(\frac{x}{2}\right)} \right) + b \left(\frac{2 \tan\left(\frac{x}{2}\right)}{1 + \tan^2\left(\frac{x}{2}\right)} \right)}$$

$$\Rightarrow I(x) = \int_0^{\frac{\pi}{2}} \frac{1 + \tan^2\left(\frac{x}{2}\right)}{a - a \tan^2\left(\frac{x}{2}\right) + 2b \tan\left(\frac{x}{2}\right)} dx$$

We know that: $1 + \tan^2 x = \sec^2 x$

$$\Rightarrow I(x) = \int_0^{\frac{\pi}{2}} \frac{\sec^2\left(\frac{x}{2}\right)}{a \left(1 - \tan^2\left(\frac{x}{2}\right) + \frac{2b}{a} \tan\left(\frac{x}{2}\right) \right)} dx$$

$$\Rightarrow I(x) = \frac{1}{a} \int_0^{\frac{\pi}{2}} \frac{\sec^2\left(\frac{x}{2}\right)}{1 - (\tan^2\left(\frac{x}{2}\right) - 2 \times \frac{b}{a} \times \tan\left(\frac{x}{2}\right) + \left(\frac{b}{a}\right)^2) + \left(\frac{b}{a}\right)^2} dx$$

$$\Rightarrow I(x) = \frac{1}{a} \int_0^{\frac{\pi}{2}} \frac{\sec^2\left(\frac{x}{2}\right)}{\frac{a^2 + b^2}{a^2} - \left(\tan\left(\frac{x}{2}\right) - \frac{b}{a}\right)^2} dx$$

$$\Rightarrow I(x) = \frac{1}{a} \int_0^{\frac{\pi}{2}} \frac{\sec^2\left(\frac{x}{2}\right)}{\left(\sqrt{\frac{a^2+b^2}{a^2}}\right)^2 - \left(\tan\left(\frac{x}{2}\right) - \frac{b}{a}\right)^2} dx \dots \dots (1)$$

Let us assume,

$$y = \tan\left(\frac{x}{2}\right) - \left(\frac{b}{a}\right)$$

Differentiating w.r.t x on both sides we get,

$$\Rightarrow dy = d\left(\tan\left(\frac{x}{2}\right) - \left(\frac{b}{a}\right)\right)$$

$$\Rightarrow dy = \frac{1}{2} \sec^2\left(\frac{x}{2}\right) dx$$

$$\Rightarrow \sec^2\left(\frac{x}{2}\right) dx = 2dy \dots \dots (2)$$

Upper limit for the Definite Integral:

$$\Rightarrow x = \frac{\pi}{2} \Rightarrow y = \tan\left(\frac{\pi}{4}\right) - \frac{b}{a}$$

$$\Rightarrow y = 1 - \frac{b}{a} \dots \dots (3)$$

Lower limit for the Definite Integral:

$$\Rightarrow x = 0 \Rightarrow y = \tan(0) - \frac{b}{a}$$

$$\Rightarrow y = -\frac{b}{a} \dots \dots (4)$$

Substituting (2),(3),(4) in eq(1), we get,

$$\Rightarrow I(x) = \frac{1}{a} \int_{\frac{b}{a}}^{1-\frac{b}{a}} \frac{2dy}{\left(\sqrt{\frac{a^2+b^2}{a^2}}\right)^2 - y^2}$$

$$\Rightarrow I(x) = \frac{2}{a} \int_{\frac{b}{a}}^{1-\frac{b}{a}} \frac{dy}{\left(\sqrt{\frac{a^2+b^2}{a^2}}\right)^2 - y^2}$$

We know that:

$$\int \frac{dx}{a^2-x^2} = \frac{1}{2a} \log\left(\frac{a+x}{a-x}\right) + C$$

We know that:

$$\int_a^b f'(x) dx = |f(x)|_a^b = f(b) - f(a)$$

[here $f'(x)$ is derivative of $f(x)$].

$$\Rightarrow I(x) = \frac{2}{a} \times \frac{1}{2 \times \sqrt{\frac{a^2 + b^2}{a^2}}} \times \log \left(\left| \frac{\sqrt{\frac{a^2 + b^2}{a^2}} + y}{\sqrt{\frac{a^2 + b^2}{a^2}} - y} \right| \right)^{1-\frac{b}{a}}$$

$$\Rightarrow I(x) = \frac{1}{\sqrt{a^2 + b^2}} \times \log \left(\left| \frac{\sqrt{a^2 + b^2} + ay}{\sqrt{a^2 + b^2} - ay} \right| \right)^{1-\frac{b}{a}}$$

$$\Rightarrow I(x) = \frac{1}{\sqrt{a^2 + b^2}} \times \left(\log \left(\frac{\sqrt{a^2 + b^2} + a \left(1 - \frac{b}{a} \right)}{\sqrt{a^2 + b^2} - a \left(1 - \frac{b}{a} \right)} \right) - \log \left(\frac{\sqrt{a^2 + b^2} + a \left(-\frac{b}{a} \right)}{\sqrt{a^2 + b^2} - a \left(-\frac{b}{a} \right)} \right) \right)$$

$$\Rightarrow I(x) = \frac{1}{\sqrt{(a^2 + b^2)}} \times \left(\log \left(\frac{\sqrt{a^2 + b^2} + a - b}{\sqrt{a^2 + b^2} - a + b} \right) - \log \left(\frac{\sqrt{a^2 + b^2} - b}{\sqrt{a^2 + b^2} + b} \right) \right)$$

We know that: $\text{Log}\left(\frac{a}{b}\right) = \log(a) - \log(b)$

$$\Rightarrow I(x) = \frac{1}{\sqrt{a^2 + b^2}} \times \left(\log \left(\frac{\left(\frac{\sqrt{a^2 + b^2} + a - b}{\sqrt{a^2 + b^2} - a + b} \right)}{\left(\frac{\sqrt{a^2 + b^2} - b}{\sqrt{a^2 + b^2} + b} \right)} \right) \right)$$

$$\Rightarrow I(x) = \frac{1}{\sqrt{a^2 + b^2}} \times \left(\log \left(\frac{(\sqrt{a^2 + b^2} + a - b) \times (\sqrt{a^2 + b^2} + b)}{(\sqrt{a^2 + b^2} - a + b) \times (\sqrt{a^2 + b^2} - b)} \right) \right)$$

$$\Rightarrow I(x) = \frac{1}{\sqrt{a^2 + b^2}} \times \left(\log \left(\frac{a^2 + b^2 + b\sqrt{a^2 + b^2} + a\sqrt{a^2 + b^2} - b\sqrt{a^2 + b^2} + ab - b^2}{a^2 + b^2 - b\sqrt{a^2 + b^2} - a\sqrt{a^2 + b^2} + b\sqrt{a^2 + b^2} + ab - b^2} \right) \right)$$

$$\Rightarrow I(x) = \frac{1}{\sqrt{a^2 + b^2}} \times \left(\log \left(\frac{a^2 + a\sqrt{a^2 + b^2} + ab}{a^2 + ab - a\sqrt{a^2 + b^2}} \right) \right)$$

$$\Rightarrow I(x) = \frac{1}{\sqrt{a^2 + b^2}} \times \left(\log \left(\frac{a \times (a + \sqrt{a^2 + b^2} + b)}{a \times (a - \sqrt{a^2 + b^2} + b)} \right) \right)$$

$$\Rightarrow I(x) = \frac{1}{\sqrt{a^2 + b^2}} \times \left(\log \left(\frac{a + b + \sqrt{a^2 + b^2}}{a + b - \sqrt{a^2 + b^2}} \right) \right)$$

$$\therefore \int_0^{\frac{\pi}{2}} \frac{dx}{a \cos x + b \sin x} \quad a, b > 0 = \frac{1}{\sqrt{a^2 + b^2}} \log \left(\frac{a + b + \sqrt{a^2 + b^2}}{a + b - \sqrt{a^2 + b^2}} \right)$$

20. Question

Evaluate the following Integrals:

$$\int_0^{\pi/2} \frac{1}{5 + 4 \sin x} dx$$

Answer

Given Definite can be written as:

$$\Rightarrow I(x) = \int_0^{\frac{\pi}{2}} \frac{dx}{5 + 4\sin x}$$

We know that:

$$\sin x = \frac{2 \tan\left(\frac{x}{2}\right)}{1 + \tan^2\left(\frac{x}{2}\right)}$$

$$\Rightarrow I(x) = \int_0^{\frac{\pi}{2}} \frac{dx}{5 + 4 \left(\frac{2 \tan\left(\frac{x}{2}\right)}{1 + \tan^2\left(\frac{x}{2}\right)} \right)}$$

$$\Rightarrow I(x) = \int_0^{\frac{\pi}{2}} \frac{1 + \tan^2\left(\frac{x}{2}\right)}{5 \left(1 + \tan^2\left(\frac{x}{2}\right) \right) + 8 \tan\left(\frac{x}{2}\right)} dx$$

$$\Rightarrow I(x) = \int_0^{\frac{\pi}{2}} \frac{1 + \tan^2\left(\frac{x}{2}\right)}{5 + 5\tan^2\left(\frac{x}{2}\right) + 8 \tan\left(\frac{x}{2}\right)} dx$$

We know that: $1 + \tan^2 x = \sec^2 x$

$$\Rightarrow I(x) = \int_0^{\frac{\pi}{2}} \frac{\sec^2\left(\frac{x}{2}\right)}{5 \left(1 + \tan^2\left(\frac{x}{2}\right) + \frac{8}{5} \tan\left(\frac{x}{2}\right) \right)} dx$$

$$\Rightarrow I(x) = \frac{1}{5} \int_0^{\frac{\pi}{2}} \frac{\sec^2\left(\frac{x}{2}\right)}{1 + \left(\tan^2\left(\frac{x}{2}\right) + 2 \times \frac{4}{5} \times \tan\left(\frac{x}{2}\right) + \left(\frac{4}{5}\right)^2 \right) - \left(\frac{4}{5}\right)^2} dx$$

$$\Rightarrow I(x) = \frac{1}{5} \int_0^{\frac{\pi}{2}} \frac{\sec^2\left(\frac{x}{2}\right)}{\frac{9}{25} + \left(\tan\left(\frac{x}{2}\right) + \frac{4}{5} \right)^2} dx$$

$$\Rightarrow I(x) = \frac{1}{5} \int_0^{\frac{\pi}{2}} \frac{\sec^2\left(\frac{x}{2}\right)}{\left(\frac{3}{5}\right)^2 + \left(\tan\left(\frac{x}{2}\right) + \left(\frac{4}{5}\right) \right)^2} dx \dots\dots (1)$$

Let us assume,

$$y = \tan\left(\frac{x}{2}\right) + \left(\frac{4}{5}\right)$$

Differentiating w.r.t x on both sides we get,

$$\Rightarrow dy = d\left(\tan\left(\frac{x}{2}\right) + \left(\frac{4}{5}\right)\right)$$

$$\Rightarrow dy = \frac{1}{2} \sec^2\left(\frac{x}{2}\right) dx$$

$$\Rightarrow \sec^2\left(\frac{x}{2}\right) dx = 2dy \dots\dots (2)$$

Upper limit for the Definite Integral:

$$\Rightarrow x = \frac{\pi}{2} \Rightarrow y = \tan\left(\frac{\pi}{4}\right) + \left(\frac{4}{5}\right)$$



$$\Rightarrow y = 1 + \frac{4}{5}$$

$$\Rightarrow y = \frac{9}{5} \dots \dots (3)$$

Lower limit for the Definite Integral:

$$\Rightarrow x = 0 \Rightarrow y = \tan(0) + \frac{4}{5}$$

$$\Rightarrow y = 0 + \frac{4}{5}$$

$$\Rightarrow y = \frac{4}{5} \dots \dots (4)$$

Substituting (2),(3),(4) in the eq(1) we get,

$$\Rightarrow I(x) = \frac{1}{5} \int_{\frac{4}{5}}^{\frac{9}{5}} \frac{2dy}{\left(\frac{3}{5}\right)^2 + y^2}$$

$$\Rightarrow I(x) = \frac{2}{5} \int_{\frac{4}{5}}^{\frac{9}{5}} \frac{dy}{\left(\frac{3}{5}\right)^2 + y^2}$$

We know that:

$$\int \frac{dx}{x^2 + a^2} = \frac{1}{a} \tan^{-1}\left(\frac{x}{a}\right) + C$$

We know that:

$$\int_a^b f'(x) dx = |f(x)|_a^b = f(b) - f(a)$$

[here $f'(x)$ is derivative of $f(x)$].

$$\Rightarrow I(x) = \frac{2}{5} \times \frac{1}{3} \times \left(\tan^{-1}\left(\frac{5y}{3}\right) \Big|_{\frac{4}{5}}^{\frac{9}{5}} \right)$$

$$\Rightarrow I(x) = \frac{2}{3} \times (\tan^{-1}(3) - \tan^{-1}\left(\frac{4}{3}\right))$$

We know that:

$$\tan^{-1}(A) - \tan^{-1}(B) = \tan^{-1}\left(\frac{A-B}{1+AB}\right) \text{ (if } AB > -1\text{)}$$

$$\Rightarrow I(x) = \frac{2}{3} \times \left(\tan^{-1}\left(\frac{3 - \frac{4}{3}}{1 + \left(3 \times \frac{4}{3}\right)}\right) \right)$$

$$\Rightarrow I(x) = \frac{2}{3} \times \left(\tan^{-1}\left(\frac{\frac{5}{3}}{1+4}\right) \right)$$

$$\Rightarrow I(x) = \frac{2}{3} \times \left(\tan^{-1}\left(\frac{\frac{5}{3}}{5}\right) \right)$$

$$\Rightarrow I(x) = \frac{2}{3} \tan^{-1} \left(\frac{1}{3} \right)$$

$$\therefore \int_0^{\frac{\pi}{2}} \frac{1}{5 + 4\sin x} dx = \frac{2}{3} \tan^{-1} \left(\frac{1}{3} \right)$$

21. Question

Evaluate the following Integrals:

$$\int_0^{\frac{\pi}{2}} \frac{\sin x}{\sin x + \cos x} dx$$

Answer

Given Definite Integral can be written as:

$$\Rightarrow I(x) = \int_0^{\frac{\pi}{2}} \frac{\sin x}{\sin x + \cos x} dx \dots \dots (1)$$

Let us write numerator in terms of the denominator for easy calculation,

$$\Rightarrow \sin x = K(\sin x + \cos x) + L \times \frac{d}{dx}(\sin x + \cos x)$$

$$\Rightarrow \sin x = K(\sin x + \cos x) + L(\cos x - \sin x)$$

$$\Rightarrow \sin x = \sin x(K-L) + \cos x(K+L)$$

Comparing coefficients of corresponding terms on both sides we get,

$$\Rightarrow K + L = 0$$

$$\Rightarrow K - L = 1$$

On solving these two equations we get,

$$L = -\frac{1}{2} \text{ and } K = \frac{1}{2}$$

So numerator can be written as:

$$\frac{1}{2}(\sin x + \cos x) - \frac{1}{2}(\cos x - \sin x)$$

Substituting these values in (1) we get,

$$\Rightarrow I(x) = \int_0^{\frac{\pi}{2}} \frac{\frac{1}{2}(\sin x + \cos x) - \frac{1}{2}(\cos x - \sin x)}{\sin x + \cos x} dx$$

$$\Rightarrow I(x) = \int_0^{\frac{\pi}{2}} \frac{\frac{1}{2}(\sin x + \cos x)}{\sin x + \cos x} dx - \int_0^{\frac{\pi}{2}} \frac{\frac{1}{2}(\cos x - \sin x)}{\sin x + \cos x} dx$$

$$\Rightarrow I(x) = \frac{1}{2} \int_0^{\frac{\pi}{2}} dx - \frac{1}{2} \int_0^{\frac{\pi}{2}} \frac{d(\sin x + \cos x)}{\sin x + \cos x}$$

We know that:

$$\int dy = y + C \text{ and } \int \frac{dy}{y} = \log|y| + C$$

We know that:

$$\int_a^b f'(x) dx = |f(x)|_a^b = f(b) - f(a)$$

[here $f'(x)$ is derivative of $f(x)$].

$$\Rightarrow I(x) = \frac{1}{2}(x|_0^\pi) - \frac{1}{2}(\log(|\sin x + \cos x|)|_0^\pi)$$

$$\Rightarrow I(x) = \frac{1}{2} \times (\pi - 0) - \frac{1}{2}((\log(|\sin \pi + \cos \pi|)) - (\log(|\sin 0 + \cos 0|)))$$

$$\Rightarrow I(x) = \frac{\pi}{2} - \frac{1}{2} \times (\log(|0 - 1|) - \log(|0 + 1|))$$

$$\Rightarrow I(x) = \frac{\pi}{2} - \frac{1}{2} \times (\log 1 - \log 1)$$

$$\Rightarrow I(x) = \frac{\pi}{2} - \frac{1}{2} \times (0 - 0)$$

$$\Rightarrow I(x) = \frac{\pi}{2}$$

$$\therefore \int_0^\pi \frac{\sin x}{\sin x + \cos x} dx = \frac{\pi}{2}$$

22. Question

Evaluate the following Integrals:

$$\int_0^\pi \frac{1}{3 + 2 \sin x + \cos x} dx$$

Answer

Given Definite Integral can be written as:

$$\Rightarrow I(x) = \int_0^\pi \frac{1}{3 + 2 \sin x + \cos x} dx$$

We know that:

$$\sin x = \frac{2 \tan\left(\frac{x}{2}\right)}{1 + \tan^2\left(\frac{x}{2}\right)} \text{ and}$$

$$\cos x = \frac{1 - \tan^2\left(\frac{x}{2}\right)}{1 + \tan^2\left(\frac{x}{2}\right)}$$

$$\Rightarrow I(x) = \int_0^\pi \frac{1}{3 + 2 \left(\frac{2 \tan\left(\frac{x}{2}\right)}{1 + \tan^2\left(\frac{x}{2}\right)} \right) + \frac{1 - \tan^2\left(\frac{x}{2}\right)}{1 + \tan^2\left(\frac{x}{2}\right)}} dx$$

$$\Rightarrow I(x) = \int_0^\pi \frac{1 + \tan^2\left(\frac{x}{2}\right)}{3 \left(1 + \tan^2\left(\frac{x}{2}\right) \right) + 4 \tan\left(\frac{x}{2}\right) + 1 - \tan^2\left(\frac{x}{2}\right)} dx$$

$$\Rightarrow I(x) = \int_0^\pi \frac{1 + \tan^2\left(\frac{x}{2}\right)}{3 + 3 \tan^2\left(\frac{x}{2}\right) + 4 \tan\left(\frac{x}{2}\right) + 1 - \tan^2\left(\frac{x}{2}\right)} dx$$

$$\Rightarrow I(x) = \int_0^{\pi} \frac{1 + \tan^2\left(\frac{x}{2}\right)}{4 + 2\tan^2\left(\frac{x}{2}\right) + 4\tan\left(\frac{x}{2}\right)} dx$$

We know that: $1 + \tan^2 x = \sec^2 x$

$$\Rightarrow I(x) = \frac{1}{2} \times \int_0^{\pi} \frac{\sec^2\left(\frac{x}{2}\right)}{2 + \left(\tan^2\left(\frac{x}{2}\right) + 2\tan\left(\frac{x}{2}\right) + 1\right) - 1} dx$$

$$\Rightarrow I(x) = \frac{1}{2} \times \int_0^{\pi} \frac{\sec^2\left(\frac{x}{2}\right)}{1^2 + \left(\tan\left(\frac{x}{2}\right) + 1\right)^2} dx \dots \dots (1)$$

Let us assume, $y = 1 + \tan\left(\frac{x}{2}\right)$

Differentiating w.r.t x on both sides we get,

$$\Rightarrow dy = d(1 + \tan\left(\frac{x}{2}\right))$$

$$\Rightarrow dy = \frac{1}{2} \sec^2\left(\frac{x}{2}\right) dx$$

$$\Rightarrow \sec^2\left(\frac{x}{2}\right) = 2dy \dots \dots (2)$$

Upper limit for the Definite Integral:

$$\Rightarrow x = \pi \Rightarrow y = 1 + \tan\left(\frac{\pi}{2}\right)$$

$$\Rightarrow y = 1 + \infty$$

$$\Rightarrow y = \infty \dots \dots (3)$$

Lower limit for the Definite Integral:

$$\Rightarrow x = 0 \Rightarrow y = 1 + \tan(0)$$

$$\Rightarrow y = 1 + 0$$

$$\Rightarrow y = 1 \dots \dots (4)$$

Substituting (2),(3),(4) in the eq(1), we get,

$$\Rightarrow I(x) = \frac{1}{2} \times \int_0^{\infty} \frac{2dy}{1 + y^2}$$

$$\Rightarrow I(x) = \int_0^{\infty} \frac{dy}{1 + y^2}$$

We know that:

$$\int \frac{1}{1 + x^2} dx = \tan^{-1} x + C$$

We know that:

$$\int_a^b f'(x) dx = |f(x)|_a^b = f(b) - f(a)$$

[here $f'(x)$ is derivative of $f(x)$].

$$\Rightarrow I(x) = \tan^{-1} y|_0^{\infty}$$

$$\Rightarrow I(x) = \tan^{-1}(\infty) - \tan^{-1}(0)$$

$$\Rightarrow I(x) = \frac{\pi}{2} - \frac{\pi}{4}$$

$$\Rightarrow I(x) = \frac{\pi}{4}$$

$$\therefore \int_0^{\pi} \frac{1}{3 + 2\sin x + \cos x} dx = \frac{\pi}{4}$$

23. Question

Evaluate the following Integrals:

$$\int_0^1 \tan^{-1} x \, dx$$

Answer

Given Definite Integral can be written as:

$$\Rightarrow I(x) = \int_0^1 \tan^{-1} x \, dx$$

We will find the value of $\int \tan^{-1} x \, dx$ using by parts rule

Let us find the value of $\int \tan^{-1} x \, dx$

$$\Rightarrow \int \tan^{-1} x \, dx = \int 1 \cdot \tan^{-1} x \, dx$$

$$\Rightarrow \tan^{-1} x \int 1 \, dx - \int \left(\frac{d}{dx} (\tan^{-1} x) \int 1 \, dx \right) dx$$

$$\Rightarrow x \tan^{-1} x - \int \frac{1}{1+x^2} \cdot x \, dx$$

$$\Rightarrow x \tan^{-1} x - \frac{1}{2} \int \frac{2x}{1+x^2} \, dx$$

$$\Rightarrow x \tan^{-1} x - \frac{1}{2} \int \frac{d(1+x^2)}{1+x^2}$$

$$\Rightarrow x \tan^{-1} x - \frac{1}{2} \log(1+x^2)$$

We substitute this result in the Definite Integral:

We know that:

$$\int_a^b f'(x) \, dx = |f(x)|_a^b = f(b) - f(a)$$

[here $f'(x)$ is derivative of $f(x)$].

$$\Rightarrow I(x) = \left(x \tan^{-1} x - \frac{1}{2} \log(1+x^2) \right) \Big|_0^1$$

$$\Rightarrow I(x) = (1 \tan^{-1}(1) - \frac{1}{2} \log(1+1)) - (0 \tan^{-1}(0) - \frac{1}{2} \log(1+0))$$

$$\Rightarrow I(x) = \left(\frac{\pi}{4} - \frac{1}{2} \log 2 \right) - (0 - 0)$$

$$\Rightarrow I(x) = \frac{\pi}{4} - \frac{1}{2} \log 2$$

$$\therefore \int_0^1 \tan^{-1} x \, dx = \frac{\pi}{4} - \frac{1}{2} \log 2$$

24. Question

Evaluate the following Integrals:

$$\int_0^{1/2} \frac{x \sin^{-1} x}{\sqrt{1-x^2}} \, dx$$

Answer

Given Definite Integral can be written as:

$$\Rightarrow I(x) = \int_0^{\frac{1}{2}} \frac{x \sin^{-1} x}{\sqrt{1-x^2}} \, dx$$

Let us find the value of $\int \frac{x \sin^{-1} x}{\sqrt{1-x^2}} \, dx$ using by parts integration,

$$\begin{aligned} \Rightarrow \int \frac{x \sin^{-1} x}{\sqrt{1-x^2}} \, dx &= \sin^{-1} x \int \frac{x}{\sqrt{1-x^2}} \, dx - \int \left(\frac{d}{dx}(\sin^{-1} x) \int \frac{x}{\sqrt{1-x^2}} \, dx \right) dx \\ \Rightarrow (\sin^{-1} x \times -\frac{1}{2} \times \int -\frac{2x}{\sqrt{1-x^2}} \, dx) &- \int \left(\frac{1}{\sqrt{1-x^2}} \times -\frac{1}{2} \times \int \frac{-2x}{\sqrt{1-x^2}} \, dx \right) dx \\ \Rightarrow (\sin^{-1} x \times -\frac{1}{2} \times \int \frac{d(1-x^2)}{\sqrt{1-x^2}}) &- \int \left(\frac{1}{\sqrt{1-x^2}} \times -\frac{1}{2} \times \int \frac{d(1-x^2)}{\sqrt{1-x^2}} \right) dx \\ \Rightarrow (\sin^{-1} x \times -\frac{1}{2} \times 2 \times \sqrt{1-x^2}) &- \int \left(\frac{1}{\sqrt{1-x^2}} \times -\frac{1}{2} \times 2 \times \sqrt{1-x^2} \right) dx \\ \Rightarrow -\sqrt{1-x^2} \sin^{-1} x &- \int -1 \, dx \\ \Rightarrow -\sqrt{1-x^2} \sin^{-1} x + x & \end{aligned}$$

Now we substitute this result in the Definite Integral:

We know that:

$$\int_a^b f'(x) \, dx = [f(x)]_a^b = f(b) - f(a)$$

[here $f'(x)$ is derivative of $f(x)$].

$$\Rightarrow I(x) = \left(x - \sqrt{1-x^2} \sin^{-1} x \right) \Big|_0^{\frac{1}{2}}$$

$$\Rightarrow I(x) = \left(\frac{1}{2} - \sqrt{1 - \left(\frac{1}{2} \right)^2} \sin^{-1} \left(\frac{1}{2} \right) \right) - (0 - \sqrt{1-0^2} \sin^{-1}(0))$$

$$\Rightarrow I(x) = \left(\frac{1}{2} - \sqrt{1 - \frac{1}{4} \times \frac{\pi}{6}} \right) - (0 - 0)$$

$$\Rightarrow I(x) = \left(\frac{1}{2} - \sqrt{\frac{3\pi}{46}} \right)$$

$$\Rightarrow I(x) = \frac{1}{2} - \frac{\sqrt{3}\pi}{12}$$

$$\therefore \int_0^{\frac{1}{2}} \frac{x \sin^{-1} x}{\sqrt{1-x^2}} dx = \frac{1}{2} - \frac{\sqrt{3}\pi}{12}$$

25. Question

Evaluate the following Integrals:

$$\int_0^{\pi/4} (\sqrt{\tan x} + \sqrt{\cot x}) dx$$

Answer

Given Definite Integral can be written as:

$$\Rightarrow I(x) = \int_0^{\pi/4} (\sqrt{\tan x} + \sqrt{\cot x}) dx$$

We know that:

$$\tan x = \frac{\sin x}{\cos x} \text{ and}$$

$$\cot x = \frac{\cos x}{\sin x}$$

Substituting in the Definite Integral we get,

$$\Rightarrow I(x) = \int_0^{\pi/4} \left(\sqrt{\frac{\sin x}{\cos x}} + \sqrt{\frac{\cos x}{\sin x}} \right) dx$$

$$\Rightarrow I(x) = \int_0^{\pi/4} \left(\frac{(\sqrt{\sin x})^2 + (\sqrt{\cos x})^2}{\sqrt{\sin x \cos x}} \right) dx$$

$$\Rightarrow I(x) = \sqrt{2} \int_0^{\pi/4} \frac{\sin x + \cos x}{\sqrt{2 \sin x \cos x}} dx$$

$$\Rightarrow I(x) = \sqrt{2} \times \int_0^{\pi/4} \frac{\sin x + \cos x}{\sqrt{(1 - (-2 \sin x \cos x + 1))}} dx$$

$$\Rightarrow I(x) = \sqrt{2} \times \int_0^{\pi/4} \frac{\sin x + \cos x}{\sqrt{1 - (\sin^2 x + \cos^2 x - 2 \sin x \cos x)}} dx$$

$$\Rightarrow I(x) = \sqrt{2} \times \int_0^{\pi/4} \frac{\sin x + \cos x}{\sqrt{1 - (\sin x - \cos x)^2}} dx \dots\dots (1)$$

Let us assume, $y = \sin x - \cos x$

Differentiating w.r.t x on both sides we get,

$$\Rightarrow d(y) = d(\sin x - \cos x)$$

$$\Rightarrow dy = (\cos x + \sin x)dx \dots\dots (2)$$

Upper limit for the Definite Integral:

$$\Rightarrow x = \frac{\pi}{4} \Rightarrow y = \sin\left(\frac{\pi}{4}\right) - \cos\left(\frac{\pi}{4}\right)$$

$$\Rightarrow y = \frac{1}{\sqrt{2}} - \frac{1}{\sqrt{2}}$$

$$\Rightarrow y = 0 \dots (3)$$

Lower limit for the Definite Integral:

$$\Rightarrow x = 0 \Rightarrow y = \sin(0) - \cos(0)$$

$$\Rightarrow y = 0 - 1$$

$$\Rightarrow y = -1 \dots (4)$$

Substituting (2),(3),(4) in the eq(1) we get,

$$\Rightarrow I(x) = \sqrt{2} \times \int_{-1}^0 \frac{dy}{\sqrt{1-y^2}}$$

We know that:

$$\int \frac{1}{\sqrt{1-x^2}} dx = \sin^{-1}x + C$$

We know that:

$$\int_a^b f'(x) dx = |f(x)|_a^b = f(b) - f(a)$$

[here $f'(x)$ is derivative of $f(x)$)

$$\Rightarrow I(x) = \sqrt{2} \times (\sin^{-1}y|_{-1}^0)$$

$$\Rightarrow I(x) = \sqrt{2} \times (\sin^{-1}(0) - \sin^{-1}(-1))$$

$$\Rightarrow I(x) = \sqrt{2} \times (0 - \left(-\frac{\pi}{2}\right))$$

$$\Rightarrow I(x) = \sqrt{2} \times \frac{\pi}{2}$$

$$\Rightarrow I(x) = \frac{\pi}{\sqrt{2}}$$

$$\therefore \int_0^{\frac{\pi}{4}} (\sqrt{\tan x} + \sqrt{\cot x}) dx = \frac{\pi}{\sqrt{2}}$$

26. Question

Evaluate the following Integrals:

$$\int_0^{\pi/4} \frac{\tan^3 x}{1 + \cos 2x} dx$$

Answer

Given Definite Integral can be written as:

$$\Rightarrow I(x) = \int_0^{\frac{\pi}{4}} \frac{\tan^3 x}{1 + \cos 2x} dx$$

We know that: $1 + \cos 2x = 2\cos^2 x$ and

$$\frac{1}{\cos^2 x} = \sec^2 x$$

$$\Rightarrow I(x) = \int_0^{\frac{\pi}{4}} \frac{\tan^3 x}{2\cos^2 x} dx$$

$$\Rightarrow I(x) = \frac{1}{2} \times \int_0^{\frac{\pi}{4}} \tan^3 x \sec^2 x dx \dots\dots (1)$$

Let us assume, $y = \tan x$

Differentiating w.r.t x on both sides we get,

$$\Rightarrow dy = d(\tan x)$$

$$\Rightarrow dy = \sec^2 x dx \dots\dots (2)$$

Upper limit for the Definite Integral:

$$\Rightarrow x = \frac{\pi}{4} \Rightarrow y = \tan\left(\frac{\pi}{4}\right)$$

$$\Rightarrow y = 1 \dots\dots (3)$$

The lower limit for the Definite Integral:

$$\Rightarrow x = 0 \Rightarrow y = \tan\left(\frac{0}{4}\right)$$

$$\Rightarrow y = 0 \dots\dots (4)$$

Substituting (2),(3),(4) in eq(1) we get,

$$\Rightarrow I(x) = \frac{1}{2} \int_0^1 y^3 dy$$

We know that:

$$\int x^n dx = \frac{x^{n+1}}{n+1} + C$$

We know that:

$$\int_a^b f'(x) dx = |f(x)|_a^b = f(b) - f(a)$$

[here $f'(x)$ is derivative of $f(x)$].

$$\Rightarrow I(x) = \frac{1}{2} \times \left(\frac{y^4}{4} \Big|_0^1 \right)$$

$$\Rightarrow I(x) = \frac{1}{2} \times \left(\frac{y^4}{4} \Big|_0^1 \right)$$

$$\Rightarrow I(x) = \frac{1}{8} \times (14 - 04)$$

$$\Rightarrow I(x) = \frac{1}{8}$$

$$\therefore \int_0^{\frac{\pi}{4}} \frac{\tan^3 x}{1 + \cos 2x} dx = \frac{1}{8}$$

27. Question

Evaluate the following Integrals:

$$\int_0^{\pi} \frac{1}{5 + 3\cos x} dx$$

Answer

Given Definite Integral can be written as:

$$\Rightarrow I(x) = \int_0^{\pi} \frac{1}{5 + 3\cos x} dx$$

We know that:

$$\cos x = \frac{1 - \tan^2(\frac{x}{2})}{1 + \tan^2(\frac{x}{2})}$$

$$\Rightarrow I(x) = \int_0^{\pi} \frac{1}{5 + 3\left(\frac{1 - \tan^2(\frac{x}{2})}{1 + \tan^2(\frac{x}{2})}\right)} dx$$

$$\Rightarrow I(x) = \int_0^{\pi} \frac{1 + \tan^2(\frac{x}{2})}{5(1 + \tan^2(\frac{x}{2})) + 3(1 - \tan^2(\frac{x}{2}))} dx$$

$$\Rightarrow I(x) = \int_0^{\pi} \frac{1 + \tan^2(\frac{x}{2})}{5 + 5\tan^2(\frac{x}{2}) + 3 - 3\tan^2(\frac{x}{2})} dx$$

$$\Rightarrow I(x) = \int_0^{\pi} \frac{1 + \tan^2(\frac{x}{2})}{8 + 2\tan^2(\frac{x}{2})} dx$$

We know that: $1 + \tan^2 x = \sec^2 x$

$$\Rightarrow I(x) = \int_0^{\pi} \frac{\frac{1}{2}\sec^2(\frac{x}{2})}{2^2 + (\tan(\frac{x}{2}))^2} dx \dots\dots (1)$$

Let us assume, $y = \tan(\frac{x}{2})$

Differentiating w.r.t x on both the sides we get,

$$\Rightarrow dy = d(\tan(\frac{x}{2}))$$

$$\Rightarrow dy = \frac{1}{2}\sec^2(\frac{x}{2}) dx \dots\dots (2)$$

The upper limit for the Definite Integral:

$$\Rightarrow x = \pi \Rightarrow y = \tan(\frac{\pi}{2})$$

$$\Rightarrow y = \infty \dots\dots (3)$$

Lower limit for the Definite Integral:

$$\Rightarrow x = 0 \Rightarrow y = \tan\left(\frac{0}{2}\right)$$

$$\Rightarrow y = 0 \dots\dots(4)$$

Substituting (2),(3),(4) in the eq(1) we get,

$$\Rightarrow I(x) = \int_0^\infty \frac{dy}{2^2 + y^2}$$

We know that:

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1}\left(\frac{x}{a}\right) + C$$

We know that:

$$\int_a^b f(x)dx = |f(x)|_a^b = f(b) - f(a)$$

[here $f'(x)$ is derivative of $f(x)$].

$$\Rightarrow I(x) = \frac{1}{2} \times \left(\tan^{-1}\left(\frac{y}{2}\right) \Big|_0^\infty \right)$$

$$\Rightarrow I(x) = \frac{1}{2} \times (\tan^{-1}(\infty) - \tan^{-1}(0))$$

$$\Rightarrow I(x) = \frac{1}{2} \times \left(\frac{\pi}{2} - 0\right)$$

$$\Rightarrow I(x) = \frac{\pi}{4}$$

$$\therefore \int_0^\pi \frac{dx}{5 + 3\cos x} = \frac{\pi}{4}$$

28. Question

Evaluate the following Integrals:

$$\int_0^{\pi/2} \frac{1}{a^2 \sin^2 x + b^2 \cos^2 x} dx$$

Answer

Given Definite Integral can be written as:

$$\Rightarrow I(x) = \int_0^{\pi/2} \frac{1}{a^2 \sin^2 x + b^2 \cos^2 x} dx$$

$$\Rightarrow I(x) = \int_0^{\pi/2} \frac{1}{\cos^2 x \times \left(\left(\frac{a^2 \sin^2 x}{\cos^2 x} \right) + b^2 \right)} dx$$

$$\Rightarrow I(x) = \int_0^{\pi/2} \frac{1}{a^2 \tan^2 x + b^2} dx$$

$$\Rightarrow I(x) = \int_0^{\pi/2} \frac{\sec^2 x}{a^2 \tan^2 x + b^2} dx$$

$$\Rightarrow I(x) = \frac{1}{a^2} \times \int_0^{\frac{\pi}{2}} \frac{\sec^2 x}{(\tan x)^2 + \left(\frac{b}{a}\right)^2} dx \dots \dots (1)$$

Let us assume $y = \tan x$

Differentiating w.r.t x on both sides we get,

$$\Rightarrow dy = d(\tan x)$$

$$\Rightarrow dy = \sec^2 x dx \dots \dots (2)$$

Upper limit for the Definite Integral:

$$\Rightarrow x = \frac{\pi}{2} \Rightarrow y = \tan\left(\frac{\pi}{2}\right)$$

$$\Rightarrow y = \infty \dots \dots (3)$$

Lower limit for the Definite Integral:

$$\Rightarrow x = 0 \Rightarrow y = \tan(0)$$

$$\Rightarrow y = 0 \dots \dots (4)$$

Substituting (2),(3),(4) in the eq(1) we get,

$$\Rightarrow I(x) = \frac{1}{a^2} \int_0^{\infty} \frac{dt}{t^2 + \left(\frac{b}{a}\right)^2}$$

We know that:

$$\int \frac{1}{a^2 + x^2} dx = \tan^{-1} x + C$$

We know that:

$$\int_a^b f(x) dx = |f(x)|_a^b = f(b) - f(a)$$

[here $f'(x)$ is derivative of $f(x)$].

$$\Rightarrow I(x) = \frac{1}{a^2} \times \frac{1}{b} \times \left(\tan^{-1} \left(\frac{x}{b} \right) \right) \Big|_0^\infty$$

$$\Rightarrow I(x) = \frac{1}{ab} \times \left(\tan^{-1} \left(\frac{ax}{b} \right) \right) \Big|_0^\infty$$

$$\Rightarrow I(x) = \frac{1}{ab} \times (\tan^{-1}(\infty) - \tan^{-1}(0))$$

$$\Rightarrow I(x) = \frac{1}{ab} \times \left(\frac{\pi}{2} - 0 \right)$$

$$\Rightarrow I(x) = \frac{\pi}{2ab}$$

$$\therefore \int_0^{\frac{\pi}{2}} \frac{1}{a^2 \sin^2 x + b^2 \cos^2 x} dx = \frac{\pi}{2ab}$$

29. Question

Evaluate the following Integrals:

$$\int_0^{\pi/2} \frac{x + \sin x}{1 + \cos x} dx$$

Answer

Given Definite Integral can be written as:

$$\Rightarrow I(x) = \int_0^{\frac{\pi}{2}} \frac{\frac{1}{2}x + \sin x}{1 + \cos x} dx$$

We know that $\sin 2x = 2 \sin x \cos x$ and $1 + \cos 2x = 2 \cos^2 x$

$$\Rightarrow I(x) = \int_0^{\frac{\pi}{2}} \frac{\frac{1}{2}x + 2 \sin\left(\frac{x}{2}\right) \cos\left(\frac{x}{2}\right)}{2 \cos^2\left(\frac{x}{2}\right)} dx$$

$$\Rightarrow I(x) = \int_0^{\frac{\pi}{2}} \frac{\frac{1}{2}x \sec^2\left(\frac{x}{2}\right)}{2} dx + \int_0^{\frac{\pi}{2}} \tan\left(\frac{x}{2}\right) dx$$

Applying by-parts integration for 1st term only

$$\Rightarrow I(x) = x \int_0^{\frac{\pi}{2}} \frac{\sec^2\left(\frac{x}{2}\right)}{2} dx - \int_0^{\frac{\pi}{2}} \left(\frac{d}{dx}(x) \int \frac{\sec^2\left(\frac{x}{2}\right)}{2} dx \right) dx + \int_0^{\frac{\pi}{2}} \tan\left(\frac{x}{2}\right) dx$$

We know that:

$$\int_a^b f(x) dx = [f(x)]_a^b = f(b) - f(a)$$

[here $f'(x)$ is derivative of $f(x)$].

$$\Rightarrow I(x) = \left(x \tan\left(\frac{x}{2}\right) \right) \Big|_0^{\frac{\pi}{2}} - \int_0^{\frac{\pi}{2}} \tan\left(\frac{x}{2}\right) dx + \int_0^{\frac{\pi}{2}} \tan\left(\frac{x}{2}\right) dx$$

$$\Rightarrow I(x) = \left(\frac{\pi}{2} \tan\left(\frac{\pi}{4}\right) - 0 \tan(0) \right)$$

$$\Rightarrow I(x) = \frac{\pi}{2}$$

$$\therefore \int_0^{\frac{\pi}{2}} \frac{\frac{1}{2}x + \sin x}{1 + \cos x} dx = \frac{\pi}{2}$$

30. Question

Evaluate the following Integrals:

$$\int_0^1 \frac{\tan^{-1} x}{1+x^2} dx$$

Answer

Given Definite Integral can be written as:

$$\Rightarrow I(x) = \int_0^1 \frac{\tan^{-1} x}{1+x^2} dx \dots\dots(1)$$

Let us assume $y = \tan^{-1} x$

Differentiating w.r.t x on both sides we get,

$$\Rightarrow d(y) = d(\tan^{-1}x)$$

$$\Rightarrow dy = \frac{1}{1+x^2} dx \dots\dots(2)$$

Upper limit for the Definite Integral:

$$\Rightarrow x = 1 \Rightarrow y = \tan^{-1}(1)$$

$$\Rightarrow y = \frac{\pi}{4} \dots\dots(3)$$

Lower limit for the Definite Integral:

$$\Rightarrow x = 0 \Rightarrow y = \tan^{-1}(0)$$

$$\Rightarrow y = 0 \dots\dots(4)$$

Substitute (2),(3),(4) in the eq(1) we get,

$$\Rightarrow I(x) = \int_0^{\frac{\pi}{4}} y dy$$

We know that:

$$\int x^n dx = \frac{x^{n+1}}{n+1} + C$$

We know that:

$$\int_a^b f'(x) dx = |f(x)|_a^b = f(b) - f(a)$$

[here $f'(x)$ is derivative of $f(x)$].

$$\Rightarrow I(x) = \frac{y^2}{2} \Big|_0^{\frac{\pi}{4}}$$

$$\Rightarrow I(x) = \frac{1}{2} \times \left(\left(\frac{\pi}{4} \right)^2 - 0^2 \right)$$

$$\Rightarrow I(x) = \frac{1}{2} \times \frac{\pi^2}{16}$$

$$\Rightarrow I(x) = \frac{\pi^2}{32}$$

$$\therefore \int_0^1 \frac{\tan^{-1}x}{1+x^2} dx = \frac{\pi^2}{32}$$

31. Question

Evaluate the following Integrals:

$$\int_0^{\pi/4} \frac{\sin x + \cos x}{3 + \sin 2x} dx$$

Answer

$$\text{Let } I = \int_0^{\pi/4} \frac{\sin x + \cos x}{3 + \sin 2x} dx$$

In the denominator, we have $\sin 2x = 2 \sin x \cos x$

Note that we can write $2 \sin x \cos x = 1 - (1 - 2 \sin x \cos x)$



We also have $\sin^2 x + \cos^2 x = 1$

$$\Rightarrow 1 - 2 \sin x \cos x = \sin^2 x + \cos^2 x - 2 \sin x \cos x$$

$$\Rightarrow \sin 2x = 1 - (\sin x - \cos x)^2$$

So, using this, we can write our integral as

$$I = \int_0^{\frac{\pi}{4}} \frac{\sin x + \cos x}{3 + [1 - (\sin x - \cos x)^2]} dx$$

$$\Rightarrow I = \int_0^{\frac{\pi}{4}} \frac{\sin x + \cos x}{4 - (\sin x - \cos x)^2} dx$$

Now, put $\sin x - \cos x = t$

$$\Rightarrow (\cos x + \sin x) dx = dt \text{ (Differentiating both sides)}$$

When $x = 0, t = \sin 0 - \cos 0 = 0 - 1 = -1$

When,

$$x = \frac{\pi}{4}, t = \sin \frac{\pi}{4} - \cos \frac{\pi}{4} = \frac{1}{\sqrt{2}} - \frac{1}{\sqrt{2}} = 0$$

So, the new limits are -1 and 0.

Substituting this in the original integral,

$$I = \int_{-1}^0 \frac{1}{4 - t^2} dt$$

$$\Rightarrow I = \int_{-1}^0 \frac{1}{2^2 - t^2} dt$$

Recall,

$$\int \frac{1}{a^2 - x^2} dx = \frac{1}{2a} \ln \left| \frac{a+x}{a-x} \right| + C$$

$$\Rightarrow I = \left[\frac{1}{2(2)} \ln \left| \frac{2+t}{2-t} \right| \right]_{-1}^0$$

$$\Rightarrow I = \frac{1}{4} \left[\ln \left| \frac{2+0}{2-0} \right| - \ln \left| \frac{2+(-1)}{2-(-1)} \right| \right]$$

$$\Rightarrow I = \frac{1}{4} \left[\ln|1| - \ln \left| \frac{1}{3} \right| \right]$$

$$\Rightarrow I = \frac{1}{4} \ln 3 [\because \ln 1 = 0 \text{ & } \ln \left(\frac{1}{x} \right) = -\ln x]$$

$$\therefore \int_0^{\frac{\pi}{4}} \frac{\sin x + \cos x}{3 + \sin 2x} dx = \frac{1}{4} \ln 3$$

32. Question

Evaluate the following Integrals:

$$\int_0^1 x \tan^{-1} x \, dx$$

Answer

$$\text{Let } I = \int_0^1 x \tan^{-1} x \, dx$$

We will use integration by parts.

$$\text{Recall, } \int f(x)g(x)dx = f(x)[\int g(x)dx] - \int [f'(x)\int g(x)dx]dx + c$$

Here, take $f(x) = \tan^{-1} x$ and $g(x) = x$

$$\Rightarrow \int g(x)dx = \int x \, dx$$

We have,

$$\int x^n dx = \frac{1}{n+1} x^{n+1} + c$$

$$\Rightarrow \int g(x)dx = \frac{x^2}{2}$$

Now,

$$f'(x) = \frac{df(x)}{dx} = \frac{d}{dx}(\tan^{-1} x)$$

$$\Rightarrow f'(x) = \frac{1}{1+x^2}$$

Substituting these values, we evaluate the integral.

$$\Rightarrow I = \left[(\tan^{-1} x) \frac{x^2}{2} \right]_0^1 - \int_0^1 \frac{1}{1+x^2} \left(\frac{x^2}{2} \right) dx$$

$$\Rightarrow I = \left[\frac{x^2}{2} \tan^{-1} x \right]_0^1 - \frac{1}{2} \int_0^1 \frac{x^2}{1+x^2} dx$$

We can write,

$$\frac{x^2}{1+x^2} = 1 - \frac{1}{1+x^2}$$

$$\Rightarrow I = \left[\frac{x^2}{2} \tan^{-1} x \right]_0^1 - \frac{1}{2} \int_0^1 \left[1 - \frac{1}{1+x^2} \right] dx$$

$$\Rightarrow I = \left[\frac{1^2}{2} \tan^{-1}(1) - \frac{0^2}{2} \tan^{-1}(0) \right] - \left(\frac{1}{2} \int_0^1 dx - \frac{1}{2} \int_0^1 \frac{1}{1+x^2} dx \right)$$

Recall,

$$\int \frac{1}{1+x^2} dx = \tan^{-1} x + c$$

$$\Rightarrow I = \frac{1}{2} \times \frac{\pi}{4} - \left(\frac{1}{2} [x]_0^1 - \frac{1}{2} [\tan^{-1} x]_0^1 \right)$$

$$\Rightarrow I = \frac{\pi}{8} - \left(\frac{1}{2} [1 - 0] - \frac{1}{2} [\tan^{-1}(1) - \tan^{-1}(0)] \right)$$

$$\Rightarrow I = \frac{\pi}{8} - \left(\frac{1}{2} - \frac{1}{2} \left[\frac{\pi}{4} - 0 \right] \right)$$

$$\Rightarrow I = \frac{\pi}{8} - \left(\frac{1}{2} - \frac{\pi}{8} \right)$$

$$\Rightarrow I = \frac{\pi}{4} - \frac{1}{2}$$

$$\therefore \int_0^1 x \tan^{-1} x dx = \frac{\pi}{4} - \frac{1}{2}$$

33. Question

Evaluate the following Integrals:

$$\int_0^1 \frac{1-x^2}{x^4+x^2+1} dx$$

Answer

$$\text{Let } I = \int_0^1 \frac{1-x^2}{x^4+x^2+1} dx$$

In the denominator, we have $x^4 + x^2 + 1$

Note that we can write $x^4 + x^2 + 1 = (x^4 + 2x^2 + 1) - x^2$

We have $x^4 + 2x^2 + 1 = (1 + x^2)^2$

$$\Rightarrow x^4 + x^2 + 1 = (1 + x^2)^2 - x^2$$

So, using this, we can write our integral as

$$I = \int_0^1 \frac{1-x^2}{(1+x^2)^2 - x^2} dx$$

Dividing numerator and denominator with x^2 , we have

$$I = \int_0^1 \frac{\frac{1-x^2}{x^2}}{\left(\frac{x^2+1}{x}\right)^2 - \frac{x^2}{x^2}} dx$$

$$\Rightarrow I = \int_0^1 \frac{\frac{1}{x^2} - 1}{\left(x + \frac{1}{x}\right)^2 - 1} dx$$

$$\text{Put, } x + \frac{1}{x} = t$$

$$\Rightarrow \left(1 - \frac{1}{x^2}\right) dx = dt$$

(Differentiating both sides)

$$\Rightarrow \left(\frac{1}{x^2} - 1\right) dx = -dt$$

When $x = 0, t = 0 + \frac{1}{0} = \infty$

When $x = 1, t = 1 + \frac{1}{1} = 2$

So, the new limits are ∞ and 2.

Substituting this in the original integral,

$$I = \int_{\infty}^2 \frac{1}{t^2 - 1} (-dt)$$

$$\Rightarrow I = - \int_{\infty}^2 \frac{1}{t^2 - 1} dt$$

Recall, $\int \frac{1}{x^2 - a^2} dx = \frac{1}{2a} \ln \left| \frac{x-a}{x+a} \right| + c$

$$\Rightarrow I = - \left[\frac{1}{2(1)} \ln \left| \frac{t-1}{t+1} \right| \right]_{\infty}^2$$

$$\Rightarrow I = - \frac{1}{2} \left[\ln \left| \frac{2-1}{2+1} \right| - \ln \left| \frac{\infty-1}{\infty+1} \right| \right]$$

$$\Rightarrow I = - \frac{1}{2} \left[\ln \frac{1}{3} - 0 \right]$$

$$\Rightarrow I = - \frac{1}{2} \ln 3^{-1}$$

$$\Rightarrow I = \ln 3^{\frac{1}{2}} = \ln \sqrt{3}$$

$$\therefore \int_0^1 \frac{1-x^2}{x^4+x^2+1} dx = \ln \sqrt{3}$$

34. Question

Evaluate the following Integrals:

$$\int_0^1 \frac{24x^3}{(1+x^2)^4} dx$$

Answer

$$\text{Let } I = \int_0^1 \frac{24x^3}{(1+x^2)^4} dx$$

$$\text{Put } 1+x^2 = t$$

$$\Rightarrow 2xdx = dt \text{ (Differentiating both sides)}$$

$$\text{When } x = 0, t = 1 + 0^2 = 1$$

$$\text{When } x = 1, t = 1 + 1^2 = 2$$

So, the new limits are 1 and 2.

In numerator, we can write $24x^3 dx = 12x^2 \times 2xdx$

$$\text{But, } x^2 = t-1 \text{ and } 2xdx = dt$$



$$\Rightarrow 24x^3 dx = 12(t - 1) dt$$

Substituting this in the original integral,

$$I = \int_1^2 \frac{12(t - 1)}{t^4} dt$$

$$\Rightarrow I = 12 \int_1^2 \left(\frac{t}{t^4} - \frac{1}{t^4} \right) dt$$

$$\Rightarrow I = 12 \left(\int_1^2 \frac{1}{t^3} dt - \int_1^2 \frac{1}{t^4} dt \right)$$

$$\Rightarrow I = 12 \left(\int_1^2 t^{-3} dt - \int_1^2 t^{-4} dt \right)$$

$$\text{Recall } \int x^n dx = \frac{1}{n+1} x^{n+1} + C$$

$$\Rightarrow I = 12 \left[\frac{t^{-3+1}}{(-3+1)} - \frac{t^{-4+1}}{(-4+1)} \right]_1^2$$

$$\Rightarrow I = 12 \left[\frac{-1}{2t^2} + \frac{1}{3t^3} \right]_1^2$$

$$\Rightarrow I = 12 \left[\left(\frac{-1}{2(2)^2} + \frac{1}{3(2)^3} \right) - \left(\frac{-1}{2(1)^2} + \frac{1}{3(1)^3} \right) \right]$$

$$\Rightarrow I = 12 \left[\left(\frac{-1}{8} + \frac{1}{24} \right) - \left(\frac{-1}{2} + \frac{1}{3} \right) \right]$$

$$\Rightarrow I = 12 \left[\frac{-1}{12} - \left(\frac{-1}{6} \right) \right]$$

$$\Rightarrow I = 12 \left[\frac{1}{6} - \frac{1}{12} \right] = 1$$

$$\therefore \int_0^1 \frac{24x^3}{(1+x^2)^4} dx = 1$$

35. Question

Evaluate the following Integrals:

$$\int_4^{12} x(x-4)^{1/3} dx$$

Answer

$$\text{Let } I = \int_4^{12} x(x-4)^{1/3} dx$$

$$\text{Put } x - 4 = t^3$$

$$\Rightarrow dx = 3t^2 dt \text{ (Differentiating both sides)}$$

$$\text{When } x = 4, t^3 = 4 - 4 = 0 \Rightarrow t = 0$$

When $x = 12$, $t^3 = 12 - 4 = 8 \Rightarrow t = 2$

So, the new limits are 0 and 2.

We can write $x = t^3 + 4$

Substituting this in the original integral,

$$I = \int_0^2 (t^3 + 4)t(3t^2) dt$$

$$\Rightarrow I = 3 \int_0^2 (t^6 + 4t^3) dt$$

$$\Rightarrow I = 3 \left(\int_0^2 t^6 dt + 4 \int_0^2 t^3 dt \right)$$

Recall,

$$\int x^n dx = \frac{1}{n+1} x^{n+1} + c$$

$$\Rightarrow I = 3 \left(\left[\frac{t^7}{7} \right]_0^2 + 4 \left[\frac{t^4}{4} \right]_0^2 \right)$$

$$\Rightarrow I = 3 \left(\left[\frac{t^7}{7} \right]_0^2 + 4 \left[\frac{t^4}{4} \right]_0^2 \right)$$

$$\Rightarrow I = 3 \left[\left(\frac{2^7}{7} - \frac{0^7}{7} \right) + 4 \left(\frac{2^4}{4} - \frac{0^4}{4} \right) \right]$$

$$\Rightarrow I = 3 \left(\frac{128}{7} + 16 \right)$$

$$\Rightarrow I = 3 \left(\frac{240}{7} \right) = \frac{720}{7}$$

$$\therefore \int_4^{12} x(x-4)^{\frac{1}{3}} dx = \frac{720}{7}$$

36. Question

Evaluate the following Integrals:

$$\int_0^{\pi/2} x^2 \sin x dx$$

Answer

Let $I = \int_0^{\pi/2} x^2 \sin x dx$

We will use integration by parts.

Recall,

$$\int f(x)g(x)dx = f(x) \left[\int g(x)dx \right] - \int \left[f'(x) \int g(x)dx \right] dx + c$$

Here, take $f(x) = x^2$ and $g(x) = \sin x$

$$\Rightarrow \int g(x)dx = \int \sin x dx = -\cos x$$

Now,

$$f'(x) = \frac{df(x)}{dx} = \frac{d}{dx}(x^2)$$

$$\Rightarrow f'(x) = 2x$$

Substituting these values, we evaluate the integral.

$$\Rightarrow I = [x^2(-\cos x)]_0^{\frac{\pi}{2}} - \int_0^{\frac{\pi}{2}} (2x)(-\cos x)dx$$

$$\Rightarrow I = [-x^2 \cos x]_0^{\frac{\pi}{2}} + 2 \int_0^{\frac{\pi}{2}} x \cos x dx$$

$$\text{Let } I_1 = \int_0^{\frac{\pi}{2}} x \cos x dx$$

We use integration by parts again.

Here, take $f(x) = x$ and $g(x) = \cos x$

$$\Rightarrow \int g(x)dx = \int \cos x dx = \sin x$$

$$\text{Now, } f'(x) = \frac{df(x)}{dx} = \frac{d}{dx}(x)$$

$$\Rightarrow f'(x) = 1$$

Using these values in equation for I_1

$$\Rightarrow I_1 = [x \sin x]_0^{\frac{\pi}{2}} - \int_0^{\frac{\pi}{2}} (1)(\sin x) dx$$

$$\Rightarrow I_1 = [x \sin x]_0^{\frac{\pi}{2}} - \int_0^{\frac{\pi}{2}} \sin x dx$$

$$\Rightarrow I_1 = [x \sin x]_0^{\frac{\pi}{2}} + [\cos x]_0^{\frac{\pi}{2}}$$

Substituting I_1 in I , we get

$$I = [-x^2 \cos x]_0^{\frac{\pi}{2}} + 2 \left([x \sin x]_0^{\frac{\pi}{2}} + [\cos x]_0^{\frac{\pi}{2}} \right)$$

$$\Rightarrow I = [-x^2 \cos x + 2x \sin x + 2 \cos x]_0^{\frac{\pi}{2}}$$

$$\Rightarrow I = \left[-\left(\frac{\pi}{2}\right)^2 \cos \frac{\pi}{2} + 2\left(\frac{\pi}{2}\right) \sin \frac{\pi}{2} + 2 \cos \frac{\pi}{2} \right] - (0 + 0 + 2 \cos 0)$$

$$\Rightarrow I = (0 + \pi + 0) - (2)$$

$$\Rightarrow I = \pi - 2$$

$$\therefore \int_0^{\frac{\pi}{2}} x^2 \sin x \, dx = \pi - 2$$

37. Question

Evaluate the following Integrals:

$$\int_0^1 \sqrt{\frac{1-x}{1+x}} \, dx$$

Answer

$$\text{Let } I = \int_0^1 \sqrt{\frac{1-x}{1+x}} \, dx$$

As we have the trigonometric identity $\frac{1-\cos 2\theta}{1+\cos 2\theta} = \tan^2 \theta$, to evaluate this integral we use $x = \cos 2\theta$

$\Rightarrow dx = -2\sin(2\theta)d\theta$ (Differentiating both sides)

$$\text{When } x = 0, \cos 2\theta = 0 \Rightarrow 2\theta = \frac{\pi}{2} \Rightarrow \theta = \frac{\pi}{4}$$

$$\text{When } x = 1, \cos 2\theta = 1 \Rightarrow 2\theta = 0 \Rightarrow \theta = 0$$

So, the new limits are $\frac{\pi}{4}$ and 0.

Substituting this in the original integral,

$$I = \int_{\frac{\pi}{4}}^0 \sqrt{\frac{1 - \cos 2\theta}{1 + \cos 2\theta}} (-2 \sin 2\theta \, d\theta)$$

$$\Rightarrow I = -2 \int_{\frac{\pi}{4}}^0 \sqrt{\tan^2 \theta} \sin 2\theta \, d\theta$$

$$\Rightarrow I = -2 \int_{\frac{\pi}{4}}^0 \tan \theta \sin 2\theta \, d\theta$$

We have $\tan \theta = \frac{\sin \theta}{\cos \theta}$ and $\sin 2\theta = 2 \sin \theta \cos \theta$

$$\Rightarrow I = -2 \int_{\frac{\pi}{4}}^0 \frac{\sin \theta}{\cos \theta} \times 2 \sin \theta \cos \theta \, d\theta$$

$$\Rightarrow I = -4 \int_{\frac{\pi}{4}}^0 \sin^2 \theta \, d\theta$$

But,

$$\sin^2 \theta = \frac{1 - \cos 2\theta}{2}$$

$$\Rightarrow I = -4 \int_{\frac{\pi}{4}}^0 \left[\frac{1 - \cos 2\theta}{2} \right] \, d\theta$$

$$\Rightarrow I = -2 \int_{\frac{\pi}{4}}^0 [1 - \cos 2\theta] d\theta$$

$$\Rightarrow I = -2 \left(\int_{\frac{\pi}{4}}^0 d\theta - \int_{\pi/4}^0 \cos 2\theta d\theta \right)$$

$$\Rightarrow I = -2 \left([\theta]_{\frac{\pi}{4}}^0 - \left[\frac{\sin 2\theta}{2} \right]_{\frac{\pi}{4}}^0 \right)$$

$$\Rightarrow I = -2 \left[\left(0 - \frac{\pi}{4} \right) - \left[\frac{\sin 0}{2} - \frac{\sin(2 \times \frac{\pi}{4})}{2} \right] \right]$$

$$\Rightarrow I = -2 \left(-\frac{\pi}{4} + \frac{1}{2} \sin \frac{\pi}{2} \right)$$

$$\Rightarrow I = -2 \left(-\frac{\pi}{4} + \frac{1}{2} \right) = \frac{\pi}{2} - 1$$

$$\therefore \int_0^1 \sqrt{\frac{1-x}{1+x}} dx = \frac{\pi}{2} - 1$$

38. Question

Evaluate the following Integrals:

$$\int_0^1 \frac{1-x^2}{(1+x^2)^2} dx$$

Answer

$$\text{Let } I = \int_0^1 \frac{1-x^2}{(1+x^2)^2} dx$$

As we have the trigonometric identity $1 + \tan^2 \theta = \sec^2 \theta$, to evaluate this integral we use $x = \tan \theta$

$\Rightarrow dx = \sec^2 \theta d\theta$ (Differentiating both sides)

When $x = 0, \tan \theta = 0 \Rightarrow \theta = 0$

$$\text{When } x = 1, \tan \theta = 1 \Rightarrow \theta = \frac{\pi}{4}$$

So, the new limits are 0 and $\frac{\pi}{4}$.

Substituting this in the original integral,

$$I = \int_0^{\frac{\pi}{4}} \frac{(1 - \tan^2 \theta)}{(\sec^2 \theta)^2} [\sec^2 \theta] d\theta$$

$$\Rightarrow I = \int_0^{\frac{\pi}{4}} \frac{(1 - \tan^2 \theta)}{\sec^2 \theta} d\theta$$

$$\Rightarrow I = \int_0^{\frac{\pi}{4}} \frac{\left(1 - \frac{\sin^2 \theta}{\cos^2 \theta}\right)}{\left(\frac{1}{\cos^2 \theta}\right)} d\theta$$

$$\Rightarrow I = \int_0^{\frac{\pi}{4}} (\cos^2 \theta - \sin^2 \theta) d\theta$$

We have $\cos^2 \theta - \sin^2 \theta = \cos 2\theta$

$$\Rightarrow I = \int_0^{\frac{\pi}{4}} \cos 2\theta d\theta$$

$$\Rightarrow I = \left[\frac{\sin 2\theta}{2} \right]_0^{\frac{\pi}{4}}$$

$$\Rightarrow I = \frac{\sin(2 \times \frac{\pi}{4})}{2} - \frac{\sin(2 \times 0)}{2}$$

$$\Rightarrow I = \frac{1}{2} \sin \frac{\pi}{2} - 0 = \frac{1}{2}$$

$$\therefore \int_0^1 \frac{1 - x^2}{(1 + x^2)^2} dx = \frac{1}{2}$$

39. Question

Evaluate the following Integrals:

$$\int_{-1}^1 5x^4 \sqrt{x^5 + 1} dx$$

Answer

$$\text{Let } I = \int_{-1}^1 5x^4 \sqrt{x^5 + 1} dx$$

$$\text{Put } x^5 + 1 = t$$

$$\Rightarrow 5x^4 dx = dt \text{ (Differentiating both sides)}$$

$$\text{When } x = -1, t = (-1)^5 + 1 = 0$$

$$\text{When } x = 1, t = 1^5 + 1 = 2$$

So, the new limits are 0 and 2.

Substituting this in the original integral,

$$I = \int_0^2 \sqrt{t} dt$$

$$\Rightarrow I = \int_0^2 t^{\frac{1}{2}} dt$$

$$\text{Recall } \int x^n dx = \frac{1}{n+1} x^{n+1} + C$$

$$\Rightarrow I = \left[\frac{t^{\frac{1}{2}+1}}{\frac{1}{2}+1} \right]_0^2$$

$$\Rightarrow I = \frac{2}{3} \left[t^{\frac{3}{2}} \right]_0^2$$

$$\Rightarrow I = \frac{2}{3} \left(2^{\frac{3}{2}} - 0^{\frac{3}{2}} \right)$$

$$\Rightarrow I = \frac{2}{3} \times \sqrt{8} = \frac{4\sqrt{2}}{3}$$

$$\therefore \int_{-1}^1 5x^4 \sqrt{x^5 + 1} dx = \frac{4\sqrt{2}}{3}$$

40. Question

Evaluate the following Integrals:

$$\int_0^{\pi/2} \frac{\cos^2 x}{1 + 3 \sin^2 x} dx$$

Answer

$$\text{Let } I = \int_0^{\frac{\pi}{2}} \frac{\cos^2 x}{1 + 3 \sin^2 x} dx$$

Dividing numerator and denominator with $\cos^2 x$, we have $I = \int_0^{\frac{\pi}{2}} \frac{1}{\sec^2 x + 3 \tan^2 x} dx$

$$\Rightarrow I = \int_0^{\frac{\pi}{2}} \frac{1}{1 + 4\tan^2 x} dx \quad [\because \sec^2 x = 1 + \tan^2 x]$$

Put $\tan x = t$

$\Rightarrow \sec^2 x dx = dt$ (Differentiating both sides)

$$\Rightarrow dx = \frac{dt}{1 + \tan^2 x} = \frac{dt}{1 + t^2}$$

When $x = 0, t = \tan 0 = 0$

When $x = \frac{\pi}{2}, t = \tan \frac{\pi}{2} = \infty$

So, the new limits are 0 and ∞ .

Substituting this in the original integral,

$$I = \int_0^{\infty} \frac{1}{1 + 4t^2} \left(\frac{dt}{1 + t^2} \right)$$

$$\Rightarrow I = \int_0^{\infty} \frac{1}{(1 + t^2)(1 + 4t^2)} dt$$

Multiplying numerator and denominator with 3, we have

$$I = \int_0^{\infty} \frac{3}{3(1+t^2)(1+4t^2)} dt$$

Now, we can write

$$\begin{aligned}\frac{3}{3(1+t^2)(1+4t^2)} &= \frac{[(4+4t^2) - (1+4t^2)]}{3(1+t^2)(1+4t^2)} \\ \Rightarrow \frac{3}{3(1+t^2)(1+4t^2)} &= \frac{4(1+t^2) - (1+4t^2)}{3(1+t^2)(1+4t^2)} \\ \Rightarrow \frac{3}{3(1+t^2)(1+4t^2)} &= \frac{4}{3} \left(\frac{1}{1+4t^2} \right) - \frac{1}{3} \left(\frac{1}{1+t^2} \right)\end{aligned}$$

Substituting this in the original integral,

$$\begin{aligned}I &= \int_0^{\infty} \left[\frac{4}{3} \left(\frac{1}{1+4t^2} \right) - \frac{1}{3} \left(\frac{1}{1+t^2} \right) \right] dt \\ \Rightarrow I &= \frac{4}{3} \int_0^{\infty} \left(\frac{1}{1+4t^2} \right) dt - \frac{1}{3} \int_0^{\infty} \left(\frac{1}{1+t^2} \right) dt \\ \Rightarrow I &= \frac{1}{3} \int_0^{\infty} \left(\frac{1}{\frac{1}{4} + t^2} \right) dt - \frac{1}{3} \int_0^{\infty} \left(\frac{1}{1+t^2} \right) dt \\ \Rightarrow I &= \frac{1}{3} \int_0^{\infty} \left(\frac{1}{\left(\frac{1}{2}\right)^2 + t^2} \right) dt - \frac{1}{3} \int_0^{\infty} \left(\frac{1}{1^2 + t^2} \right) dt\end{aligned}$$

Recall $\int \frac{1}{a^2+x^2} dx = \frac{1}{a} \tan^{-1} \left(\frac{x}{a} \right) + C$

$$\begin{aligned}\Rightarrow I &= \frac{1}{3} \left[\frac{1}{\left(\frac{1}{2}\right)} \tan^{-1} \left(\frac{x}{\left(\frac{1}{2}\right)} \right) \right]_0^{\infty} - \frac{1}{3} \left[\tan^{-1} \left(\frac{x}{1} \right) \right]_0^{\infty} \\ \Rightarrow I &= \frac{2}{3} [\tan^{-1} 2x]_0^{\infty} - \frac{1}{3} [\tan^{-1} x]_0^{\infty} \\ \Rightarrow I &= \frac{2}{3} [\tan^{-1}(\infty) - \tan^{-1}(0)] - \frac{1}{3} [\tan^{-1}(\infty) - \tan^{-1}(0)] \\ \Rightarrow I &= \frac{2}{3} \left(\frac{\pi}{2} \right) - \frac{1}{3} \left(\frac{\pi}{2} \right) = \frac{\pi}{6} \\ \therefore \int_0^{\frac{\pi}{2}} \frac{\cos^2 x}{1+3\sin^2 x} dx &= \frac{\pi}{6}\end{aligned}$$

41. Question

Evaluate the following Integrals:

$$\int_0^{\pi/4} \sin^3 2t \cos 2t dt$$

Answer

$$\text{Let } I = \int_0^{\frac{\pi}{4}} \sin^3 2t \cos 2t dt$$

Put $\sin 2t = x$

$\Rightarrow 2\cos(2t)dt = dx$ (Differentiating both sides)

$$\Rightarrow \cos(2t)dt = \frac{dx}{2}$$

When $t = 0, x = \sin 0 = 0$

$$\text{When } t = \frac{\pi}{4}, x = \sin 2 \times \frac{\pi}{4} = \sin \frac{\pi}{2} = 1$$

So, the new limits are 0 and 1.

Substituting this in the original integral,

$$I = \int_0^1 x^3 \left(\frac{dx}{2}\right)$$

$$\Rightarrow I = \frac{1}{2} \int_0^1 x^3 dx$$

$$\text{Recall } \int x^n dx = \frac{1}{n+1} x^{n+1} + C$$

$$\Rightarrow I = \frac{1}{2} \left[\frac{x^4}{4} \right]_0^1$$

$$\Rightarrow I = \frac{1}{8} [x^4]_0^1$$

$$\Rightarrow I = \frac{1}{8} (1^4 - 0^4) = \frac{1}{8}$$

$$\therefore \int_0^{\frac{\pi}{4}} \sin^3 2t \cos 2t dt = \frac{1}{8}$$

42. Question

Evaluate the following Integrals:

$$\int_0^{\frac{\pi}{4}} 5(5 - 4\cos \theta)^{1/4} \sin \theta d\theta$$

Answer

$$\text{Let } I = \int_0^{\frac{\pi}{4}} 5(5 - 4\cos \theta)^{\frac{1}{4}} \sin \theta d\theta$$

Put $5 - 4 \cos \theta = x$

$\Rightarrow 4\sin(\theta)d\theta = dx$ (Differentiating both sides)

$$\Rightarrow \sin(\theta)d\theta = \frac{dx}{4}$$

When $\theta = 0, x = 5 - 4 \cos 0 = 5 - 4 = 1$

When $\theta = \pi, x = 5 - 4 \cos \pi = 5 - (-4) = 9$

So, the new limits are 1 and 9.

Substituting this in the original integral,

$$I = \int_1^9 5x^{\frac{1}{4}} \left(\frac{dx}{4}\right)$$

$$\Rightarrow I = \frac{5}{4} \int_1^9 x^{\frac{1}{4}} dx$$

Recall $\int x^n dx = \frac{1}{n+1} x^{n+1} + C$

$$\Rightarrow I = \frac{5}{4} \left[\frac{x^{\frac{1}{4}+1}}{\frac{1}{4}+1} \right]_1^9$$

$$\Rightarrow I = \left[x^{\frac{5}{4}} \right]_1^9$$

$$\Rightarrow I = 9^{\frac{5}{4}} - 1$$

$$\Rightarrow I = (3^2)^{\frac{5}{4}} - 1$$

$$\Rightarrow I = \sqrt{3^5} - 1 = 9\sqrt{3} - 1$$

$$\therefore \int_0^{\pi} 5(5 - 4 \cos \theta)^{\frac{1}{4}} \sin \theta d\theta = 9\sqrt{3} - 1$$

43. Question

Evaluate the following Integrals:

$$\int_0^{\pi/6} \cos^{-3} 2\theta \sin 2\theta d\theta$$

Answer

$$\text{Let } I = \int_0^{\pi/6} \cos^{-3} 2\theta \sin 2\theta d\theta$$

Put $\cos 2\theta = x$

$\Rightarrow -2\sin(2\theta)d\theta = dx$ (Differentiating both sides)

$$\Rightarrow \sin(2\theta)d\theta = -\frac{dx}{2}$$

When $\theta = 0, x = \cos 0 = 1$

$$\text{When } \theta = \frac{\pi}{6}, x = \cos 2 \times \frac{\pi}{6} = \cos \frac{\pi}{3} = \frac{1}{2}$$

So, the new limits are 1 and $\frac{1}{2}$.

Substituting this in the original integral,

$$I = \int_1^{\frac{1}{2}} x^{-3} \left(-\frac{dx}{2} \right)$$

$$\Rightarrow I = -\frac{1}{2} \int_1^{\frac{1}{2}} x^{-3} dx$$

$$\text{Recall } \int x^n dx = \frac{1}{n+1} x^{n+1} + C$$

$$\Rightarrow I = -\frac{1}{2} \left[\frac{x^{-3+1}}{-3+1} \right]_1^{\frac{1}{2}}$$

$$\Rightarrow I = \frac{1}{4} [x^{-2}]_1^{\frac{1}{2}}$$

$$\Rightarrow I = \frac{1}{4} \left[\frac{1}{x^2} \right]_1^{\frac{1}{2}}$$

$$\Rightarrow I = \frac{1}{4} \left[\frac{1}{(\frac{1}{2})^2} - \frac{1}{1^2} \right]$$

$$\Rightarrow I = \frac{1}{4} (4 - 1) = \frac{3}{4}$$

$$\therefore \int_0^{\frac{\pi}{6}} \cos^{-3} 2\theta \sin 2\theta d\theta = \frac{3}{4}$$

44. Question

Evaluate the following Integrals:

$$\int_0^{(\pi)^{2/3}} \sqrt{x} \cos^2 x^{3/2} dx$$

Answer

$$\text{Let } I = \int_0^{\pi^{2/3}} \sqrt{x} \cos^2 x^{3/2} dx$$

$$\text{Put } x^{3/2} = t$$

$$\Rightarrow \frac{3}{2} x^{1/2} dx = dt$$

(Differentiating both sides)

$$\Rightarrow \sqrt{x} dx = \frac{2}{3} dt$$

$$\text{When } x = 0, t = 0^{\frac{3}{2}} = 0$$

$$\text{When } x = \pi^{\frac{2}{3}}, t = (\pi^{\frac{2}{3}})^{\frac{3}{2}} = \pi$$

So, the new limits are 0 and π .

Substituting this in the original integral,

$$I = \int_0^\pi \cos^2 t \left(\frac{2}{3} dt\right)$$

$$\Rightarrow I = \frac{2}{3} \int_0^\pi \cos^2 t dt$$

But,

$$\cos^2 t = \frac{1 + \cos 2t}{2}$$

$$\Rightarrow I = \frac{2}{3} \int_0^\pi \left[\frac{1 + \cos 2t}{2} \right] dt$$

$$\Rightarrow I = \frac{1}{3} \int_0^\pi [1 + \cos 2t] dt$$

$$\Rightarrow I = \frac{1}{3} \left(\int_0^\pi dt + \int_0^\pi \cos 2t dt \right)$$

$$\Rightarrow I = \frac{1}{3} \left([t]_0^\pi + \left[\frac{\sin 2t}{2} \right]_0^\pi \right)$$

$$\Rightarrow I = \frac{1}{3} \left[(\pi - 0) + \left[\frac{\sin 2\pi}{2} - \frac{\sin 0}{2} \right] \right]$$

$$\Rightarrow I = \frac{1}{3} (\pi + 0) = \frac{\pi}{3}$$

$$\therefore \int_0^{\frac{\pi}{2}} \sqrt{x} \cos^2 x^{\frac{3}{2}} dx = \frac{\pi}{3}$$

45. Question

Evaluate the following Integrals:

$$\int_1^2 \frac{1}{x(1+\log x)^2} dx$$

Answer

$$\text{Let } I = \int_1^2 \frac{1}{x(1+\log x)^2} dx$$

$$\text{Put } 1 + \log x = t$$

Differentiating both sides, we get,

$$\Rightarrow \frac{1}{x} dx = dt$$

Now, changing the limits,

$$\text{When } x = 1, t = 1 + \log 1 = 1$$



When $x = 2$, $t = 1 + \log 2$

So, the new limits are 1 and $1 + \log 2$.

Substituting this in the original integral,

$$I = \int_1^{1+\log 2} \frac{1}{t^2} dt$$

$$\Rightarrow I = \int_1^{1+\log 2} t^{-2} dt$$

Recall $\int x^n dx = \frac{1}{n+1} x^{n+1} + C$

$$\Rightarrow I = \left[\frac{t^{-2+1}}{-2+1} \right]_1^{1+\log 2}$$

$$\Rightarrow I = -[t^{-1}]_1^{1+\log 2}$$

$$\Rightarrow I = -\left[\frac{1}{t} \right]_1^{1+\log 2}$$

$$\Rightarrow I = -\left[\frac{1}{1+\log 2} - \frac{1}{1} \right]$$

$$\Rightarrow I = 1 - \frac{1}{1+\log 2} = \frac{\log 2}{1+\log 2}$$

$$\therefore \int_1^2 \frac{1}{x(1+\log x)^2} dx = \frac{\log 2}{1+\log 2}$$

46. Question

Evaluate the following Integrals:

$$\int_0^{\pi/2} \cos^5 x dx$$

Answer

$$\text{Let } I = \int_0^{\pi/2} \cos^5 x dx$$

Note that we can write $\cos^5 x = \cos^4 x \times \cos x$

$$\Rightarrow \cos^5 x = (\cos^2 x)^2 \times \cos x$$

We also have $\sin^2 x + \cos^2 x = 1$

$$\Rightarrow \cos^5 x = (1 - \sin^2 x)^2 \cos x$$

$$\text{So, } I = \int_0^{\pi/2} (1 - \sin^2 x)^2 \cos x dx$$

Put $\sin x = t$

$$\Rightarrow \cos x dx = dt \text{ (Differentiating both sides)}$$

When $x = 0$, $t = \sin 0 = 0$

$$\text{When } x = \frac{\pi}{2}, t = \sin \frac{\pi}{2} = 1$$

So, the new limits are 0 and 1.

Substituting this in the original integral,

$$I = \int_0^1 (1 - t^2)^2 dt$$

$$\Rightarrow I = \int_0^1 (1 - 2t^2 + t^4) dt$$

$$\Rightarrow I = \int_0^1 dt - 2 \int_0^1 t^2 dt + \int_0^1 t^4 dt$$

$$\text{Recall } \int x^n dx = \frac{1}{n+1} x^{n+1} + C$$

$$\Rightarrow I = [t]_0^1 - 2 \left[\frac{t^3}{3+1} \right]_0^1 + \left[\frac{t^5}{5+1} \right]_0^1$$

$$\Rightarrow I = [t]_0^1 - \frac{2}{3} [t^3]_0^1 + \frac{1}{5} [t^5]_0^1$$

$$\Rightarrow I = (1 - 0) - \frac{2}{3}(1^3 - 0) + \frac{1}{5}(1^5 - 0)$$

$$\Rightarrow I = 1 - \frac{2}{3} + \frac{1}{5} = \frac{8}{15}$$

$$\therefore \int_0^{\frac{\pi}{2}} \cos^5 x dx = \frac{8}{15}$$

47. Question

Evaluate the following Integrals:

$$\int_4^9 \frac{\sqrt{x}}{(30-x^{3/2})^2} dx$$

Answer

$$\text{Let } I = \int_4^9 \frac{\sqrt{x}}{(30-x^{3/2})^2} dx$$

$$\text{Put } 30 - x^{3/2} = t$$

$$\Rightarrow -\frac{3}{2} x^{1/2} dx = dt$$

(Differentiating both sides)

$$\Rightarrow \sqrt{x} dx = -\frac{2}{3} dt$$

$$\text{When } x = 4, t = 30 - (4)^{\frac{3}{2}} = 30 - 8 = 22$$

$$\text{When } x = 9, t = 30 - (9)^{\frac{3}{2}} = 30 - 27 = 3$$

So, the new limits are 22 and 3.

Substituting this in the original integral,

$$I = \int_{22}^3 \frac{1}{t^2} \left(-\frac{2}{3} dt \right)$$

$$\Rightarrow I = -\frac{2}{3} \int_{22}^3 t^{-2} dt$$

$$\text{Recall } \int x^n dx = \frac{1}{n+1} x^{n+1} + C$$

$$\Rightarrow I = -\frac{2}{3} \left[\frac{t^{-2+1}}{-2+1} \right]_{22}$$

$$\Rightarrow I = \frac{2}{3} [t^{-1}]_{22}^3$$

$$\Rightarrow I = \frac{2}{3} \left[\frac{1}{t} \right]_{22}^3$$

$$\Rightarrow I = \frac{2}{3} \left[\frac{1}{3} - \frac{1}{22} \right]$$

$$\Rightarrow I = \frac{2}{3} \times \frac{19}{66} = \frac{19}{99}$$

$$\therefore \int_4^9 \frac{\sqrt{x}}{(30 - x^2)^2} dx = \frac{19}{99}$$

48. Question

Evaluate the following Integrals:

$$\int_0^\pi \sin^3 x (1 + 2 \cos x) (1 + \cos x)^2 dx$$

Answer

$$\text{Let } I = \int_0^\pi \sin^3 x (1 + 2 \cos x) (1 + \cos x)^2 dx$$

Note that we can write $\sin^3 x = \sin^2 x \times \sin x$

We also have $\sin^2 x + \cos^2 x = 1$

$$\Rightarrow \sin^3 x = (1 - \cos^2 x) \sin x$$

$$\text{So, } I = \int_0^\pi (1 - \cos^2 x) (1 + 2 \cos x) (1 + \cos x)^2 \sin x dx$$

Put $\cos x = t$

$\Rightarrow -\sin(x)dx = dt$ (Differentiating both sides)

$$\Rightarrow \sin(x)dx = -dt$$

When $x = 0, t = \cos 0 = 1$

When $x = \pi, t = \cos \pi = -1$

So, the new limits are 1 and -1.

Substituting this in the original integral,

$$I = \int_1^{-1} (1-t^2)(1+2t)(1+t)^2(-dt)$$

$$\Rightarrow I = - \int_1^{-1} (1-t^2)(1+2t)(1+2t+t^2)dt$$

$$\Rightarrow I = - \int_1^{-1} (1+2t-t^2-2t^3)(1+2t+t^2)dt$$

$$\Rightarrow I = - \int_1^{-1} \left[\frac{1(1+2t+t^2) + 2t(1+2t+t^2) - t^2(1+2t+t^2)}{2t^3(1+2t+t^2)} \right] dt$$

$$\Rightarrow I = - \int_1^{-1} (1+2t+t^2+2t+4t^2+2t^3-t^2-2t^3-t^4-2t^3-4t^4-2t^5)dt$$

$$\Rightarrow I = - \int_1^{-1} (1+4t+4t^2-2t^3-5t^4-2t^5)dt$$

$$\Rightarrow I = - \left(\int_1^{-1} dt + 4 \int_1^{-1} tdt + 4 \int_1^{-1} t^2 dt - 2 \int_1^{-1} t^3 dt - 5 \int_1^{-1} t^4 dt - 2 \int_1^{-1} t^5 dt \right)$$

Recall $\int x^n dx = \frac{1}{n+1}x^{n+1} + C$

$$\Rightarrow I = - \left([t]_1^{-1} + 4 \left[\frac{t^{1+1}}{1+1} \right]_1^{-1} + 4 \left[\frac{t^{2+1}}{2+1} \right]_1^{-1} - 2 \left[\frac{t^{3+1}}{3+1} \right]_1^{-1} - 5 \left[\frac{t^{4+1}}{4+1} \right]_1^{-1} - 2 \left[\frac{t^{5+1}}{5+1} \right]_1^{-1} \right)$$

$$\Rightarrow I = - \left([t]_1^{-1} + 2[t^2]_1^{-1} + \frac{4}{3}[t^3]_1^{-1} - \frac{1}{2}[t^4]_1^{-1} - [t^5]_1^{-1} - \frac{1}{3}[t^6]_1^{-1} \right)$$

$$\Rightarrow I = - \left[(-1-1) + 2((-1)^2 - 1^2) + \frac{4}{3}((-1)^3 - 1^3) - \frac{1}{2}((-1)^4 - 1^4) - ((-1)^5 - 1^5) - \frac{1}{3}((-1)^6 - 1^6) \right]$$

$$\Rightarrow I = - \left[-2 + 2(0) + \frac{4}{3}(-2) - \frac{1}{2}(0) - (-2) - \frac{1}{3}(0) \right]$$

$$\Rightarrow I = 2 + \frac{8}{3} + (-2) = \frac{8}{3}$$

$$\therefore \int_0^{\pi} \sin^3 x (1+2\cos x)(1+\cos x)^2 dx = \frac{8}{3}$$

49. Question

Evaluate the following Integrals:

$$\int_0^{\pi/2} 2 \sin x \cos x \tan^{-1}(\sin x) dx$$

Answer

Let $I = \int_0^{\frac{\pi}{2}} 2 \sin x \cos x \tan^{-1}(\sin x) dx$

Put $\sin x = t$

$\Rightarrow \cos x dx = dt$ (Differentiating both sides)

When $x = 0, t = \sin 0 = 0$

When $x = \frac{\pi}{2}, t = \sin \frac{\pi}{2} = 1$

So, the new limits are 0 and 1.

Substituting this in the original integral,

$$I = \int_0^1 2t \tan^{-1} t dt$$

$$\Rightarrow I = 2 \int_0^1 t \tan^{-1} t dt$$

We will use integration by parts.

Recall $\int f(x)g(x)dx = f(x)[\int g(x)dx] - \int [f'(x)\int g(x)dx]dx + c$

Here, take $f(t) = \tan^{-1} t$ and $g(t) = t$

$$\Rightarrow \int g(t) dt = \int t dt = \frac{t^2}{2}$$

Now,

$$f'(t) = \frac{df(t)}{dt} = \frac{d}{dt}(\tan^{-1} t)$$

$$\Rightarrow f'(t) = \frac{1}{1+t^2}$$

Substituting these values, we evaluate the integral.

$$\Rightarrow I = 2 \left(\left[\tan^{-1} t \left(\frac{t^2}{2} \right) \right]_0^1 - \int_0^1 \left(\frac{1}{1+t^2} \right) \left(\frac{t^2}{2} \right) dt \right)$$

$$\Rightarrow I = 2 \left[\frac{t^2}{2} \tan^{-1} t \right]_0^1 - \int_0^1 \left(\frac{t^2}{1+t^2} \right) dt$$

We can write,

$$\frac{t^2}{1+t^2} = 1 - \frac{1}{1+t^2}$$

$$\Rightarrow I = 2 \left[\frac{t^2}{2} \tan^{-1} t \right]_0^1 - \int_0^1 \left[1 - \frac{1}{1+t^2} \right] dt$$

$$\Rightarrow I = 2 \left[\frac{1^2}{2} \tan^{-1}(1) - \frac{0^2}{2} \tan^{-1}(0) \right] - \left(\int_0^1 dt - \int_0^1 \frac{1}{1+t^2} dt \right)$$

Recall $\int \frac{1}{1+x^2} dx = \tan^{-1} x + C$

$$\Rightarrow I = 2 \times \frac{1}{2} \times \frac{\pi}{4} - ([t]_0^1 - [\tan^{-1} t]_0^1)$$

$$\Rightarrow I = \frac{\pi}{4} - ([1 - 0] - [\tan^{-1}(1) - \tan^{-1}(0)])$$

$$\Rightarrow I = \frac{\pi}{4} - \left(1 - \left[\frac{\pi}{4} - 0 \right] \right)$$

$$\Rightarrow I = \frac{\pi}{4} - 1 + \frac{\pi}{4} = \frac{\pi}{2} - 1$$

$$\therefore \int_0^{\frac{\pi}{2}} 2 \sin x \cos x \tan^{-1}(\sin x) dx = \frac{\pi}{2} - 1$$

50. Question

Evaluate the following Integrals:

$$\int_0^{\pi/2} \sin 2x \tan^{-1}(\sin x) dx$$

Answer

$$\text{Let } I = \int_0^{\frac{\pi}{2}} \sin 2x \tan^{-1}(\sin x) dx$$

We have $\sin 2x = 2 \sin x \cos x$

$$\Rightarrow I = \int_0^{\frac{\pi}{2}} 2 \sin x \cos x \tan^{-1}(\sin x) dx$$

Put $\sin x = t$

$\Rightarrow \cos x dx = dt$ (Differentiating both sides)

When $x = 0, t = \sin 0 = 0$

When $x = \frac{\pi}{2}, t = \sin \frac{\pi}{2} = 1$

So, the new limits are 0 and 1.

Substituting this in the original integral,

$$I = \int_0^1 2t \tan^{-1} t dt$$

$$\Rightarrow I = 2 \int_0^1 t \tan^{-1} t dt$$

We will use integration by parts.

Recall $\int f(x)g(x)dx = f(x)[\int g(x)dx] - \int [f'(x)\int g(x)dx]dx + C$

Here, take $f(t) = \tan^{-1}t$ and $g(t) = t$

$$\Rightarrow \int g(t) dt = \int t dt = \frac{t^2}{2}$$

$$\text{Now, } f'(t) = \frac{df(t)}{dt} = \frac{d}{dt}(\tan^{-1}t)$$

$$\Rightarrow f'(t) = \frac{1}{1+t^2}$$

Substituting these values, we evaluate the integral.

$$\Rightarrow I = 2 \left(\left[\tan^{-1} t \left(\frac{t^2}{2} \right) \right]_0^1 - \int_0^1 \left(\frac{1}{1+t^2} \right) \left(\frac{t^2}{2} \right) dt \right)$$

$$\Rightarrow I = 2 \left[\frac{t^2}{2} \tan^{-1} t \right]_0^1 - \int_0^1 \left(\frac{t^2}{1+t^2} \right) dt$$

$$\text{We can write } \frac{t^2}{1+t^2} = 1 - \frac{1}{1+t^2}$$

$$\Rightarrow I = 2 \left[\frac{t^2}{2} \tan^{-1} t \right]_0^1 - \int_0^1 \left[1 - \frac{1}{1+t^2} \right] dt$$

$$\Rightarrow I = 2 \left[\frac{1^2}{2} \tan^{-1}(1) - 0 \right] - \left(\int_0^1 dt - \int_0^1 \frac{1}{1+t^2} dt \right)$$

$$\text{Recall } \int \frac{1}{1+x^2} dx = \tan^{-1} x + c$$

$$\Rightarrow I = \frac{\pi}{4} - ([t]_0^1 - [\tan^{-1} t]_0^1)$$

$$\Rightarrow I = \frac{\pi}{4} - ([1-0] - [\tan^{-1}(1) - \tan^{-1}(0)])$$

$$\Rightarrow I = \frac{\pi}{4} - \left(1 - \left[\frac{\pi}{4} - 0 \right] \right)$$

$$\Rightarrow I = \frac{\pi}{4} - 1 + \frac{\pi}{4} = \frac{\pi}{2} - 1$$

$$\therefore \int_0^{\frac{\pi}{2}} \sin 2x \tan^{-1}(\sin x) dx = \frac{\pi}{2} - 1$$

51. Question

Evaluate the following Integrals:

$$\int_0^1 (\cos^{-1} x^2) dx$$

Answer

$$\text{Let } I = \int_0^1 (\cos^{-1} x)^2 dx$$

$$\text{Put } \cos^{-1} x = t$$

$$\Rightarrow x = \cos t$$

$\Rightarrow dx = -\sin t dt$ (Differentiating both sides)

$$\text{When } x = 0, t = \cos^{-1}(0) = \frac{\pi}{2}$$

$$\text{When } x = 1, t = \cos^{-1}(1) = 0$$

So, the new limits are $\frac{\pi}{2}$ and 0.

Substituting this in the original integral,

$$I = \int_{\frac{\pi}{2}}^0 t^2 (-\sin t dt)$$

$$\Rightarrow I = - \int_{\frac{\pi}{2}}^0 t^2 \sin t dt$$

We will use integration by parts.

$$\text{Recall } \int f(x)g(x)dx = f(x)[\int g(x)dx] - \int [f'(x)\int g(x)dx]dx + c$$

Here, take $f(t) = t^2$ and $g(t) = \sin t$

$$\Rightarrow \int g(t)dt = \int \sin t dt = -\cos t$$

$$\text{Now, } f'(t) = \frac{df(t)}{dt} = \frac{d}{dt}(t^2)$$

$$\Rightarrow f'(t) = 2t$$

Substituting these values, we evaluate the integral.

$$\Rightarrow I = - \left([t^2(-\cos t)]_{\frac{\pi}{2}}^0 - \int_{\frac{\pi}{2}}^0 (2t)(-\cos t) dt \right)$$

$$\Rightarrow I = [t^2 \cos t]_{\frac{\pi}{2}}^0 - 2 \int_{\frac{\pi}{2}}^0 t \cos t dt$$

$$\text{Let } I_1 = \int_{\frac{\pi}{2}}^0 t \cos t dt$$

We use integration by parts again.

Here, take $f(t) = t$ and $g(t) = \cos t$

$$\Rightarrow \int g(t)dt = \int \cos t dt = \sin t$$

$$\text{Now, } f'(t) = \frac{df(t)}{dt} = \frac{d}{dt}(t)$$

$$\Rightarrow f'(t) = 1$$

Using these values in equation for I_1

$$\Rightarrow I_1 = [t \sin t]_{\frac{\pi}{2}}^0 - \int_{\frac{\pi}{2}}^0 (1)(\sin t) dt$$

$$\Rightarrow I_1 = [t \sin t]_{\frac{\pi}{2}}^0 + [\cos t]_{\frac{\pi}{2}}^0$$

Substituting I_1 in I , we get

$$I = [t^2 \cos t]_{\frac{\pi}{2}}^0 - 2 \left([t \sin t]_{\frac{\pi}{2}}^0 + [\cos t]_{\frac{\pi}{2}}^0 \right)$$

$$\Rightarrow I = [t^2 \cos t - 2t \sin t - 2 \cos t]_{\frac{\pi}{2}}^0$$

$$\Rightarrow I = (0 - 0 - 2 \cos 0) - \left[\left(\frac{\pi}{2}\right)^2 \cos\left(\frac{\pi}{2}\right) - 2\left(\frac{\pi}{2}\right) \sin\left(\frac{\pi}{2}\right) - 2 \cos\left(\frac{\pi}{2}\right) \right]$$

$$\Rightarrow I = -2 - (-\pi) = \pi - 2$$

$$\therefore \int_0^1 (\cos^{-1} x)^2 dx = \pi - 2$$

52. Question

Evaluate the following Integrals:

$$\int_0^a \sin^{-1} \sqrt{\frac{x}{a+x}} dx$$

Answer

$$\text{Let } I = \int_0^a \sin^{-1} \sqrt{\frac{x}{a+x}} dx$$

$$\text{Put } x = \tan^2 \theta$$

$$\Rightarrow x = 2a \tan \theta \sec^2 \theta d\theta \text{ (Differentiating both sides)}$$

$$\text{When } x = 0, \tan^2 \theta = 0 \Rightarrow \tan \theta = 0 \Rightarrow \theta = 0$$

$$\text{When } x = a, \tan^2 \theta = a \Rightarrow \tan \theta = 1 \Rightarrow \theta = \frac{\pi}{4}$$

$$\text{So, the new limits are 0 and } \frac{\pi}{4}.$$

Also,

$$\sqrt{\frac{x}{a+x}} = \sqrt{\frac{a \tan^2 \theta}{a + \tan^2 \theta}}$$

$$\Rightarrow \sqrt{\frac{x}{a+x}} = \sqrt{\frac{\tan^2 \theta}{1 + \tan^2 \theta}}$$

We have the trigonometric identity $1 + \tan^2 \theta = \sec^2 \theta$

$$\Rightarrow \sqrt{\frac{\tan^2 \theta}{1 + \tan^2 \theta}} = \sqrt{\frac{\tan^2 \theta}{\sec^2 \theta}} = \sqrt{\left(\frac{\sin^2 \theta}{\cos^2 \theta}\right)} = \sqrt{\left(\frac{1}{\cos^2 \theta}\right)}$$

$$\Rightarrow \sqrt{\frac{\tan^2 \theta}{1 + \tan^2 \theta}} = \sqrt{\sin^2 \theta} = \sin \theta$$

Substituting this in the original integral,

$$I = \int_0^{\frac{\pi}{4}} \sin^{-1}(\sin \theta) (2a \tan \theta \sec^2 \theta d\theta)$$

$$\Rightarrow I = 2a \int_0^{\frac{\pi}{4}} \theta \tan \theta \sec^2 \theta d\theta$$

Now, put $\tan \theta = t$

$$\Rightarrow \sec^2 \theta d\theta = dt \text{ (Differentiating both sides)}$$

When $\theta = 0, t = \tan 0 = 0$

$$\text{When } \theta = \frac{\pi}{4}, t = \tan \frac{\pi}{4} = 1$$

So, the new limits are 0 and 1.

Substituting this in the original integral,

$$I = 2a \int_0^1 (\tan^{-1} t) (t) dt$$

$$\Rightarrow I = 2a \int_0^1 t \tan^{-1} t dt$$

We will use integration by parts.

$$\text{Recall } \int f(x)g(x)dx = f(x)[\int g(x)dx] - \int [f'(x)\int g(x)dx]dx + c$$

Here, take $f(t) = \tan^{-1} t$ and $g(t) = t$

$$\Rightarrow \int g(t) dt = \int t dt = \frac{t^2}{2}$$

Now,

$$f'(t) = \frac{df(t)}{dt} = \frac{d}{dt}(\tan^{-1} t)$$

$$\Rightarrow f'(t) = \frac{1}{1+t^2}$$

Substituting these values, we evaluate the integral.

$$\Rightarrow I = 2a \left(\left[\tan^{-1} t \left(\frac{t^2}{2} \right) \right]_0^1 - \int_0^1 \left(\frac{1}{1+t^2} \right) \left(\frac{t^2}{2} \right) dt \right)$$

$$\Rightarrow I = 2a \left[\frac{t^2}{2} \tan^{-1} t \right]_0^1 - a \int_0^1 \left(\frac{t^2}{1+t^2} \right) dt$$

$$\text{We can write } \frac{t^2}{1+t^2} = 1 - \frac{1}{1+t^2}$$

$$\Rightarrow I = 2a \left[\frac{t^2}{2} \tan^{-1} t \right]_0^1 - a \int_0^1 \left[1 - \frac{1}{1+t^2} \right] dt$$

$$\Rightarrow I = 2a \left[\frac{1^2}{2} \tan^{-1}(1) - 0 \right] - a \left(\int_0^1 dt - \int_0^1 \frac{1}{1+t^2} dt \right)$$

Recall $\int \frac{1}{1+x^2} dx = \tan^{-1} x + C$

$$\Rightarrow I = 2a \times \frac{1}{2} \times \frac{\pi}{4} - a([t]_0^1 - [\tan^{-1} t]_0^1)$$

$$\Rightarrow I = \frac{\pi}{4} a - a([1 - 0] - [\tan^{-1}(1) - \tan^{-1}(0)])$$

$$\Rightarrow I = \frac{\pi}{4} a - a \left(1 - \left[\frac{\pi}{4} - 0 \right] \right)$$

$$\Rightarrow I = \frac{\pi}{4} a - a + \frac{\pi}{4} a = \left(\frac{\pi}{2} - 1 \right) a$$

$$\therefore \int_0^a \sin^{-1} \sqrt{\frac{x}{a+x}} dx = \left(\frac{\pi}{2} - 1 \right) a$$

53. Question

Evaluate the following Integrals:

$$\int_{\pi/3}^{\pi/2} \frac{\sqrt{1+\cos x}}{(1-\cos x)^{3/2}} dx$$

Answer

$$\text{Let } I = \int_{\pi/3}^{\pi/2} \frac{\sqrt{1+\cos x}}{(1-\cos x)^{3/2}} dx$$

In the denominator, we can write

$$(1-\cos x)^{\frac{3}{2}} = (1-\cos x)\sqrt{1-\cos x}$$

$$\Rightarrow I = \int_{\pi/3}^{\pi/2} \frac{\sqrt{1+\cos x}}{(1-\cos x)\sqrt{1-\cos x}} dx$$

$$\Rightarrow I = \int_{\pi/3}^{\pi/2} \frac{1}{(1-\cos x)} \sqrt{\frac{1+\cos x}{1-\cos x}} dx$$

Recall the trigonometric identity,

$$\frac{1-\cos(2\theta)}{1+\cos(2\theta)} = \tan^2 \theta$$

Here, we have,

$$\sqrt{\frac{1+\cos x}{1-\cos x}} = \sqrt{\frac{1}{\tan^2(\frac{x}{2})}} = \frac{1}{\tan(\frac{x}{2})}$$

We also have,

$$1 - \cos x = 2 \sin^2\left(\frac{x}{2}\right)$$

$$\Rightarrow I = \int_{\frac{\pi}{3}}^{\frac{\pi}{2}} \frac{1}{2 \sin^2\left(\frac{x}{2}\right)} \left(\frac{1}{\tan\left(\frac{x}{2}\right)} \right) dx$$

$$\Rightarrow I = \frac{1}{2} \int_{\frac{\pi}{3}}^{\frac{\pi}{2}} \operatorname{cosec}^2\left(\frac{x}{2}\right) \cot\left(\frac{x}{2}\right) dx$$

$$\text{Put } \cot\left(\frac{x}{2}\right) = t$$

$$\Rightarrow -\frac{1}{2} \operatorname{cosec}^2\left(\frac{x}{2}\right) dx = dt$$

(Differentiating both sides)

$$\Rightarrow \frac{1}{2} \operatorname{cosec}^2\left(\frac{x}{2}\right) dx = -dt$$

$$\text{When } x = \frac{\pi}{3}, t = \cot\left(\frac{\pi}{6}\right) = \cot\frac{\pi}{6} = \sqrt{3}$$

$$\text{When } x = \frac{\pi}{2}, t = \cot\left(\frac{\pi}{4}\right) = \cot\frac{\pi}{4} = 1$$

So, the new limits are $\sqrt{3}$ and 1.

Substituting this in the original integral,

$$I = \int_{\sqrt{3}}^1 t(-dt)$$

$$\Rightarrow I = - \int_{\sqrt{3}}^1 t dt$$

$$\text{Recall } \int x^n dx = \frac{1}{n+1} x^{n+1} + C$$

$$\Rightarrow I = - \left[\frac{t^{1+1}}{1+1} \right]_{\sqrt{3}}^1$$

$$\Rightarrow I = -\frac{1}{2} [t^2]_{\sqrt{3}}^1$$

$$\Rightarrow I = -\frac{1}{2} [1^2 - (\sqrt{3})^2]$$

$$\Rightarrow I = -\frac{1}{2} [1 - 3] = 1$$

$$\therefore \int_{\pi/3}^{\pi/2} \frac{\sqrt{1 + \cos x}}{(1 - \cos x)^{3/2}} dx = 1$$

54. Question

Evaluate the following Integrals:

$$\int_0^a x \sqrt{\frac{a^2 - x^2}{a^2 + x^2}} dx$$

Answer

$$\text{Let } I = \int_0^a x \sqrt{\frac{a^2 - x^2}{a^2 + x^2}} dx$$

As we have the trigonometric identity

$$\frac{1 - \cos(2\theta)}{1 + \cos(2\theta)} = \tan^2 \theta$$

to evaluate this integral we use $x^2 = a^2 \cos 2\theta$

$$\Rightarrow 2x dx = -2a^2 \sin(2\theta) d\theta \text{ (Differentiating both sides)}$$

$$\Rightarrow x dx = -a^2 \sin(2\theta) d\theta$$

$$\text{When } x = 0, a^2 \cos 2\theta = 0 \Rightarrow \cos 2\theta = 0$$

$$\Rightarrow 2\theta = \frac{\pi}{2} \Rightarrow \theta = \frac{\pi}{4}$$

$$\text{When } x = a, a^2 \cos 2\theta = a^2 \Rightarrow \cos 2\theta = 1$$

$$\Rightarrow 2\theta = 0 \Rightarrow \theta = 0$$

So, the new limits are $\frac{\pi}{4}$ and 0.

Also,

$$\sqrt{\frac{a^2 - x^2}{a^2 + x^2}} = \sqrt{\frac{a^2 - a^2 \cos 2\theta}{a^2 + a^2 \cos 2\theta}}$$

$$\Rightarrow \sqrt{\frac{a^2 - x^2}{a^2 + x^2}} = \sqrt{\frac{1 - \cos 2\theta}{1 + \cos 2\theta}}$$

$$\Rightarrow \sqrt{\frac{a^2 - x^2}{a^2 + x^2}} = \sqrt{\tan^2 \theta} = \tan \theta$$

Substituting this in the original integral,

$$I = \int_{\frac{\pi}{4}}^0 \tan \theta (-a^2 \sin 2\theta d\theta)$$

$$\Rightarrow I = -a^2 \int_{\frac{\pi}{4}}^0 \tan \theta \sin 2\theta d\theta$$

$$\Rightarrow I = -a^2 \int_{\frac{\pi}{4}}^0 \frac{\sin \theta}{\cos \theta} \times 2 \sin \theta \cos \theta d\theta$$

$$\Rightarrow I = -a^2 \int_{\frac{\pi}{4}}^0 2 \sin^2 \theta d\theta$$

But, we have $2 \sin^2 \theta = 1 - \cos 2\theta$

$$\Rightarrow I = -a^2 \int_{\frac{\pi}{4}}^0 (1 - \cos 2\theta) d\theta$$

$$\Rightarrow I = -a^2 \left[\int_{\frac{\pi}{4}}^0 d\theta - \int_{\frac{\pi}{4}}^0 \cos 2\theta d\theta \right]$$

$$\Rightarrow I = -a^2 \left([0]_{\frac{\pi}{4}}^0 - \left[\frac{\sin 2\theta}{2} \right]_{\frac{\pi}{4}}^0 \right)$$

$$\Rightarrow I = -a^2 \left[\left(0 - \frac{\pi}{4} \right) - \frac{1}{2} \left(\sin 0 - \sin \left(2 \times \frac{\pi}{4} \right) \right) \right]$$

$$\Rightarrow I = -a^2 \left[-\frac{\pi}{4} - \frac{1}{2} \left(-\sin \left(\frac{\pi}{2} \right) \right) \right]$$

$$\Rightarrow I = -a^2 \left[-\frac{\pi}{4} + \frac{1}{2} \right] = a^2 \left(\frac{\pi}{4} - \frac{1}{2} \right)$$

$$\therefore \int_0^a x \sqrt{\frac{a^2 - x^2}{a^2 + x^2}} dx = a^2 \left(\frac{\pi}{4} - \frac{1}{2} \right)$$

55. Question

Evaluate the following Integrals:

$$\int_{-a}^a \sqrt{\frac{a-x}{a+x}} dx$$

Answer

$$\text{Let } I = \int_{-a}^a \sqrt{\frac{a-x}{a+x}} dx$$

As we have the trigonometric identity

$$\frac{1 - \cos(2\theta)}{1 + \cos(2\theta)} = \tan^2 \theta$$

to evaluate this integral we use $x = a \cos 2\theta$

$\Rightarrow dx = -2a \sin(2\theta) d\theta$ (Differentiating both sides)

When $x = -a$, $a \cos 2\theta = -a \Rightarrow \cos 2\theta = -1$

$$\Rightarrow 2\theta = \pi \Rightarrow \theta = \frac{\pi}{2}$$

When $x = a$, $a \cos 2\theta = a \Rightarrow \cos 2\theta = 1$

$$\Rightarrow 2\theta = 0 \Rightarrow \theta = 0$$

So, the new limits are $\frac{\pi}{2}$ and 0.

Also,

$$\sqrt{\frac{a-x}{a+x}} = \sqrt{\frac{a - a\cos 2\theta}{a + a\cos 2\theta}}$$

$$\Rightarrow \sqrt{\frac{a-x}{a+x}} = \sqrt{\frac{1 - \cos 2\theta}{1 + \cos 2\theta}}$$

$$\Rightarrow \sqrt{\frac{a-x}{a+x}} = \sqrt{\tan^2 \theta} = \tan \theta$$

Substituting this in the original integral,

$$I = \int_{\frac{\pi}{2}}^0 \tan \theta (-2a \sin 2\theta d\theta)$$

$$\Rightarrow I = -2a \int_{\frac{\pi}{2}}^0 \tan \theta \sin 2\theta d\theta$$

$$\Rightarrow I = -2a \int_{\frac{\pi}{2}}^0 \frac{\sin \theta}{\cos \theta} \times 2 \sin \theta \cos \theta d\theta$$

$$\Rightarrow I = -2a \int_{\frac{\pi}{2}}^0 2 \sin^2 \theta d\theta$$

But, we have $2 \sin^2 \theta = 1 - \cos 2\theta$

$$\Rightarrow I = -2a \int_{\frac{\pi}{2}}^0 (1 - \cos 2\theta) d\theta$$

$$\Rightarrow I = -2a \left[\int_{\frac{\pi}{2}}^0 d\theta - \int_{\frac{\pi}{2}}^0 \cos 2\theta d\theta \right]$$

$$\Rightarrow I = -2a \left([\theta]_{\frac{\pi}{2}}^0 - \left[\frac{\sin 2\theta}{2} \right]_{\frac{\pi}{2}}^0 \right)$$

$$\Rightarrow I = -2a \left[\left(0 - \frac{\pi}{2} \right) - \frac{1}{2} (\sin 0 - \sin(2 \times \frac{\pi}{2})) \right]$$

$$\Rightarrow I = -2a \left[-\frac{\pi}{2} - \frac{1}{2}(0) \right]$$

$$\Rightarrow I = -2a \left[-\frac{\pi}{2} \right] = \pi a$$

$$\therefore \int_{-a}^a \sqrt{\frac{a-x}{a+x}} dx = \pi a$$

56. Question

Evaluate the following Integrals:

$$\int_0^{\pi/2} \frac{\sin x \cos x}{\cos^2 x + 3 \cos x + 2} dx$$

Answer

$$\text{Let } I = \int_0^{\pi/2} \frac{\sin x \cos x}{\cos^2 x + 3 \cos x + 2} dx$$

In the denominator, we can write

$$\cos^2 x + 3 \cos x + 2 = (\cos x + 1)(\cos x + 2)$$

$$\Rightarrow I = \int_0^{\pi/2} \frac{\sin x \cos x}{(\cos x + 1)(\cos x + 2)} dx$$

Put $\cos x = t$

$$\Rightarrow -\sin(x)dx = dt \text{ (Differentiating both sides)}$$

$$\Rightarrow \sin(x)dx = -dt$$

When $x = 0, t = \cos 0 = 1$

$$\text{When } x = \frac{\pi}{2}, t = \cos \frac{\pi}{2} = 0$$

So, the new limits are 1 and 0.

Substituting this in the original integral,

$$I = \int_1^0 \frac{t}{(t+1)(t+2)} (-dt)$$

$$\Rightarrow I = - \int_1^0 \frac{t}{(t+1)(t+2)} dt$$

We can write,

$$\frac{t}{(t+1)(t+2)} = \frac{2t+2 - (t+2)}{(t+1)(t+2)}$$

$$\Rightarrow \frac{t}{(t+1)(t+2)} = \frac{2(t+1) - (t+2)}{(t+1)(t+2)} = \frac{2}{t+2} - \frac{1}{t+1}$$

Using this, we have

$$I = - \int_1^0 \left(\frac{2}{t+2} - \frac{1}{t+1} \right) dt$$

$$\Rightarrow I = - \left(2 \int_1^0 \frac{1}{t+2} dt - \int_1^0 \frac{1}{t+1} dt \right)$$

$$\text{Recall } \int \frac{1}{x+a} dx = \ln|x+a| + c$$

$$\Rightarrow I = -(2[\ln|x+2|]_1^0 - [\ln|x+1|]_1^0)$$

$$\Rightarrow I = -[2(\ln|0+2| - \ln|1+2|) - (\ln|0+1| - \ln|1+1|)]$$

$$\Rightarrow I = -[2(\ln 2 - \ln 3) - (\ln 1 - \ln 2)]$$

$$\Rightarrow I = -(2 \ln 2 - 2 \ln 3 - 0 + \ln 2)$$

$$\Rightarrow I = -(3 \ln 2 - 2 \ln 3)$$

$$\Rightarrow I = 2 \ln 3 - 3 \ln 2$$

$$\Rightarrow I = \ln 9 - \ln 8 = \ln \frac{9}{8}$$

$$\therefore \int_0^{\frac{\pi}{2}} \frac{\sin x \cos x}{\cos^2 x + 3 \cos x + 2} dx = \ln \frac{9}{8}$$

57. Question

Evaluate the following Integrals:

$$\int_0^{\pi/2} \frac{\tan x}{1 + m^2 \tan^2 x} dx$$

Answer

$$\text{Let } I = \int_0^{\frac{\pi}{2}} \frac{\tan x}{1 + m^2 \tan^2 x} dx$$

$$\Rightarrow I = \int_0^{\frac{\pi}{2}} \frac{\frac{\sin x}{\cos x}}{1 + m^2 \left(\frac{\sin^2 x}{\cos^2 x}\right)} dx$$

$$\Rightarrow I = \int_0^{\frac{\pi}{2}} \frac{\sin x \cos x}{\cos^2 x + m^2 \sin^2 x} dx$$

We have $\sin^2 x + \cos^2 x = 1$

$$\Rightarrow I = \int_0^{\frac{\pi}{2}} \frac{\sin x \cos x}{1 + (m^2 - 1) \sin^2 x} dx$$

Put $\sin^2 x = t$

$\Rightarrow 2 \sin x \cos x dx = dt$ (Differentiating both sides)

$$\Rightarrow \sin x \cos x dx = \frac{1}{2} dt$$

When $x = 0, t = \sin^2 0 = 0$

$$\text{When } x = \frac{\pi}{2}, t = \sin^2 \frac{\pi}{2} = 1$$

So, the new limits are 0 and 1.

Substituting this in the original integral,

$$I = \int_0^1 \frac{1}{1 + (m^2 - 1)t} \left(\frac{1}{2} dt\right)$$

$$\Rightarrow I = \frac{1}{2} \int_0^1 \frac{1}{1 + (m^2 - 1)t} dt$$

Recall $\int \frac{1}{ax+b} dx = \frac{1}{a} \ln|ax+b| + c$

$$\Rightarrow I = \frac{1}{2} \left(\frac{1}{m^2 - 1} [\ln|1 + (m^2 - 1)t|]_0^1 \right)$$

$$\Rightarrow I = \frac{1}{2(m^2 - 1)} (\ln|1 + (m^2 - 1) \times 1| - \ln|1 + (m^2 - 1) \times 0|)$$

$$\Rightarrow I = \frac{1}{2(m^2 - 1)} (\ln|m^2| - \ln|1|)$$

$$\Rightarrow I = \frac{1}{2(m^2 - 1)} (2 \ln|m|) = \frac{\ln|m|}{m^2 - 1}$$

$$\therefore \int_0^{\frac{\pi}{2}} \frac{\tan x}{1 + m^2 \tan^2 x} dx = \frac{\ln|m|}{m^2 - 1}$$

58. Question

Evaluate the following Integrals:

$$\int_0^{1/2} \frac{1}{(1+x^2)\sqrt{1-x^2}} dx$$

Answer

$$\text{Let } I = \int_0^{\frac{1}{2}} \frac{1}{(1+x^2)\sqrt{1-x^2}} dx$$

Put $x = \sin \theta$

$\Rightarrow dx = \cos \theta d\theta$ (Differentiating both sides)

Also, $\sqrt{1-x^2} = \sqrt{1-\sin^2 \theta} = \cos \theta$

When $x = 0, \sin \theta = 0 \Rightarrow \theta = 0$

When $x = \frac{1}{2}, \sin \theta = \frac{1}{2} \Rightarrow \theta = \frac{\pi}{6}$

So, the new limits are 0 and $\frac{\pi}{6}$.

Substituting this in the original integral,

$$I = \int_0^{\frac{\pi}{6}} \frac{1}{(1 + \sin^2 \theta) \cos \theta} (\cos \theta d\theta)$$

$$\Rightarrow I = \int_0^{\frac{\pi}{6}} \frac{1}{1 + \sin^2 \theta} d\theta$$

Dividing numerator and denominator with $\cos^2 \theta$, we have

$$I = \int_0^{\frac{\pi}{6}} \frac{\sec^2 \theta}{\sec^2 \theta + \tan^2 \theta} d\theta$$

$$\Rightarrow I = \int_0^{\frac{\pi}{6}} \frac{\sec^2 \theta}{1 + 2\tan^2 \theta} d\theta$$

[$\because \sec^2 \theta = 1 + \tan^2 \theta$]

Put $\tan \theta = t$

$\Rightarrow \sec^2 \theta d\theta = dt$ (Differentiating both sides)

When $\theta = 0, t = \tan 0 = 0$

$$\text{When } \theta = \frac{\pi}{6}, t = \tan \frac{\pi}{6} = \frac{1}{\sqrt{3}}$$

So, the new limits are 0 and $\frac{1}{\sqrt{3}}$.

Substituting this in the original integral,

$$I = \int_0^{\frac{1}{\sqrt{3}}} \frac{1}{1 + 2t^2} dt$$

$$\Rightarrow I = \frac{1}{2} \int_0^{\frac{1}{\sqrt{3}}} \frac{1}{\frac{1}{2} + t^2} dt$$

Recall $\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \left(\frac{x}{a} \right) + C$

$$\Rightarrow I = \frac{1}{2} \left(\frac{1}{\left(\frac{1}{\sqrt{2}}\right)} \left[\tan^{-1} \left(\frac{t}{\frac{1}{\sqrt{2}}} \right) \right]_0^{\frac{1}{\sqrt{3}}} \right)$$

$$\Rightarrow I = \frac{1}{\sqrt{2}} \left[\tan^{-1} (\sqrt{2}t) \right]_0^{\frac{1}{\sqrt{3}}}$$

$$\Rightarrow I = \frac{1}{\sqrt{2}} \left[\tan^{-1} \left(\frac{\sqrt{2}}{\sqrt{3}} \right) - \tan^{-1}(0) \right]$$

$$\Rightarrow I = \frac{1}{\sqrt{2}} \left(\tan^{-1} \left(\sqrt{\frac{2}{3}} \right) - 0 \right) = \frac{1}{\sqrt{2}} \tan^{-1} \left(\sqrt{\frac{2}{3}} \right)$$

$$\therefore \int_0^{\frac{1}{2}} \frac{1}{(1+x^2)\sqrt{1-x^2}} dx = \frac{1}{\sqrt{2}} \tan^{-1} \left(\sqrt{\frac{2}{3}} \right)$$

59. Question

Evaluate the following Integrals:

$$\int_{1/3}^1 \frac{(x-x^3)^{1/3}}{x^4} dx$$

Answer

$$\text{Let } I = \int_{\frac{1}{3}}^1 \frac{(x-x^2)^{\frac{1}{3}}}{x^4} dx$$

$$\Rightarrow I = \int_{\frac{1}{3}}^1 \frac{x^3 \left(\frac{1}{x^2} - 1\right)^{\frac{1}{3}}}{x^4} dx$$

(taking x^3 common)

$$\Rightarrow I = \int_{\frac{1}{3}}^1 \frac{x \left(\frac{1}{x^2} - 1\right)^{\frac{1}{3}}}{x^4} dx$$

$$\Rightarrow I = \int_{\frac{1}{3}}^1 \frac{\left(\frac{1}{x^2} - 1\right)^{\frac{1}{3}}}{x^3} dx$$

$$\text{Put } \frac{1}{x^2} - 1 = t$$

$\Rightarrow -2x^{-2-1}dx = dt$ (Differentiating both sides)

$$\Rightarrow -\frac{2}{x^3} dx = dt$$

$$\Rightarrow \frac{1}{x^3} dx = -\frac{1}{2} dt$$

$$\text{When } x = \frac{1}{3}, t = \frac{1}{\left(\frac{1}{3}\right)^2} - 1 = 32 - 1 = 8$$

$$\text{When } x = 1, t = \frac{1}{1^2} - 1 = 0$$

So, the new limits are 8 and 0.

Substituting this in the original integral,

$$I = \int_8^0 t^{\frac{1}{3}} \left(-\frac{dt}{2}\right)$$

$$\Rightarrow I = -\frac{1}{2} \int_8^0 t^{\frac{1}{3}} dt$$

$$\text{Recall } \int x^n dx = \frac{1}{n+1} x^{n+1} + C$$

$$\Rightarrow I = -\frac{1}{2} \left[\frac{t^{\frac{1}{3}+1}}{\frac{1}{3}+1} \right]_8^0$$

$$\Rightarrow I = -\frac{3}{8} \left[t^{\frac{4}{3}} \right]_8^0$$

$$\Rightarrow I = -\frac{3}{8} \left[0 - (8)^{\frac{4}{3}} \right]$$

$$\Rightarrow I = -\frac{3}{8}[-16] = 6$$

$$\therefore \int_{\frac{1}{3}}^1 \frac{(x-x^3)^{\frac{1}{3}}}{x^4} dx = 6$$

60. Question

Evaluate the following Integrals:

$$\int_0^{\pi/4} \frac{\sin^2 x \cos^2 x}{(\sin^3 x + \cos^3 x)^2} dx$$

Answer

$$\text{Let } I = \int_0^{\pi/4} \frac{\sin^2 x \cos^2 x}{(\sin^2 x + \cos^2 x)^2} dx$$

$$\Rightarrow I = \int_0^{\pi/4} \frac{\sin^2 x \cos^2 x}{\left[(\cos^3 x) \left(\frac{\sin^3 x}{\cos^3 x} + 1\right)\right]^2} dx$$

(taking $\cos^3 x$ common)

$$\Rightarrow I = \int_0^{\pi/4} \frac{\sin^2 x \cos^2 x}{\cos^6 x (\tan^3 x + 1)^2} dx$$

$$\Rightarrow I = \int_0^{\pi/4} \frac{\frac{\sin^2 x}{\cos^2 x} \times \frac{1}{\cos^2 x}}{(\tan^3 x + 1)^2} dx$$

$$\Rightarrow I = \int_0^{\pi/4} \frac{\tan^2 x \sec^2 x}{(\tan^3 x + 1)^2} dx$$

Put $\tan x = t$

$\Rightarrow \sec^2 x dx = dt$ (Differentiating both sides)

When $x = 0, t = \tan 0 = 0$

When $x = \frac{\pi}{4}, t = \tan \frac{\pi}{4} = 1$

So, the new limits are 0 and 1.

Substituting this in the original integral,

$$I = \int_0^1 \frac{t^2}{(1 + t^3)^2} dt$$

Put $t^3 = u$

$\Rightarrow 3t^2 dt = du$ (Differentiating both sides)

$$\Rightarrow t^2 dt = \frac{1}{3} du$$

When $t = 0$, $u = 0^3 = 0$

When $t = 1$, $u = 1^3 = 1$

So, the new limits are 0 and 1.

Substituting this in the original integral,

$$I = \int_0^1 \frac{1}{(1+u)^2} \left(\frac{1}{3} du\right)$$

$$\Rightarrow I = \frac{1}{3} \int_0^1 (u+1)^{-2} du$$

Recall $\int (x+a)^n dx = \frac{1}{n+1} (x+a)^{n+1} + c$

$$\Rightarrow I = \frac{1}{3} \left[\frac{(u+1)^{-2+1}}{-2+1} \right]_0^1$$

$$\Rightarrow I = -\frac{1}{3} \left[\frac{1}{u+1} \right]_0^1$$

$$\Rightarrow I = -\frac{1}{3} \left(\frac{1}{1+1} - \frac{1}{0+1} \right)$$

$$\Rightarrow I = -\frac{1}{3} \left(\frac{1}{2} - 1 \right) = \frac{1}{6}$$

$$\therefore \int_0^{\frac{\pi}{4}} \frac{\sin^2 x \cos^2 x}{(\sin^3 x + \cos^3 x)^2} dx = \frac{1}{6}$$

61. Question

Evaluate the following Integrals:

$$\int_0^{\pi/2} \sqrt{\cos x - \cos^3 x} (\sec^2 x - 1) \cos^2 x dx$$

Answer

$$\text{Let } I = \int_0^{\frac{\pi}{2}} \sqrt{\cos x - \cos^3 x} (\sec^2 x - 1) \cos^2 x dx$$

$$\Rightarrow I = \int_0^{\frac{\pi}{2}} \sqrt{\cos x (1 - \cos^2 x)} (\sec^2 x - 1) \cos^2 x dx$$

We have $\sin^2 x + \cos^2 x = 1$ and $\sec^2 x - \tan^2 x = 1$

$$\Rightarrow I = \int_0^{\frac{\pi}{2}} \sqrt{\cos x \sin^2 x} (\tan^2 x) \cos^2 x dx$$

$$\Rightarrow I = \int_0^{\frac{\pi}{2}} \sqrt{\cos x} \sin x \left(\frac{\sin^2 x}{\cos^2 x} \right) \cos^2 x dx$$

$$\Rightarrow I = \int_0^{\frac{\pi}{2}} \sqrt{\cos x} \sin^3 x \, dx$$

We can write $\sin^3 x = \sin^2 x \times \sin x = (1 - \cos^2 x) \sin x$

$$\Rightarrow I = \int_0^{\frac{\pi}{2}} \sqrt{\cos x} (1 - \cos^2 x) \sin x \, dx$$

Put $\cos x = t$

$\Rightarrow -\sin(x)dx = dt$ (Differentiating both sides)

$\Rightarrow \sin(x)dx = -dt$

When $x = 0, t = \cos 0 = 1$

When $x = \frac{\pi}{2}, t = \cos \frac{\pi}{2} = 0$

So, the new limits are 1 and 0.

Substituting this in the original integral,

$$I = \int_1^0 \sqrt{t}(1-t^2)(-dt)$$

$$\Rightarrow I = - \int_1^0 t^{\frac{1}{2}}(1-t^2)dt$$

$$\Rightarrow I = - \int_1^0 \left(t^{\frac{1}{2}} - t^{\frac{5}{2}} \right) dt$$

$$\Rightarrow I = - \left(\int_1^0 t^{\frac{1}{2}} dt - \int_1^0 t^{\frac{5}{2}} dt \right)$$

Recall $\int x^n dx = \frac{1}{n+1}x^{n+1} + C$

$$\Rightarrow I = - \left(\left[\frac{t^{\frac{1}{2}+1}}{\frac{1}{2}+1} \right]_1^0 - \left[\frac{t^{\frac{5}{2}+1}}{\frac{5}{2}+1} \right]_1^0 \right)$$

$$\Rightarrow I = - \left(\frac{2}{3} \left[t^{\frac{3}{2}} \right]_1^0 - \frac{2}{7} \left[t^{\frac{7}{2}} \right]_1^0 \right)$$

$$\Rightarrow I = - \left[\frac{2}{3} (0-1) - \frac{2}{7} (0-1) \right]$$

$$\Rightarrow I = - \left(-\frac{2}{3} + \frac{2}{7} \right) = \frac{8}{21}$$

$$\therefore \int_0^{\frac{\pi}{2}} \sqrt{\cos x - \cos^3 x} (\sec^2 x - 1) \cos^2 x \, dx = \frac{8}{21}$$

62. Question

Evaluate the following Integrals:

$$\int_0^{\pi/2} \frac{\cos x}{\left(\cos \frac{x}{2} + \sin \frac{x}{2}\right)^n} dx$$

Answer

$$\text{Let } I = \int_0^{\pi/2} \frac{\cos x}{\left(\cos \frac{x}{2} + \sin \frac{x}{2}\right)^n} dx$$

We can write,

$$\begin{aligned} \cos x &= \cos\left(2 \times \frac{x}{2}\right) = \cos^2 \frac{x}{2} - \sin^2 \frac{x}{2} \\ \Rightarrow \cos x &= \left(\cos \frac{x}{2} + \sin \frac{x}{2}\right)\left(\cos \frac{x}{2} - \sin \frac{x}{2}\right) \end{aligned}$$

Putting this value in the integral

$$I = \int_0^{\pi/2} \frac{\left(\cos \frac{x}{2} + \sin \frac{x}{2}\right)\left(\cos \frac{x}{2} - \sin \frac{x}{2}\right)}{\left(\cos \frac{x}{2} + \sin \frac{x}{2}\right)^n} dx$$

$$\Rightarrow I = \int_0^{\pi/2} \frac{\cos \frac{x}{2} - \sin \frac{x}{2}}{\left(\cos \frac{x}{2} + \sin \frac{x}{2}\right)^{n-1}} dx$$

$$\text{Put } \cos \frac{x}{2} + \sin \frac{x}{2} = t$$

$$\Rightarrow \left(-\frac{1}{2} \sin \frac{x}{2} + \frac{1}{2} \cos \frac{x}{2}\right) dx = dt$$

(Differentiating both sides)

$$\Rightarrow \left(\cos \frac{x}{2} - \sin \frac{x}{2}\right) dx = 2dt$$

When $x = 0$, $t = \cos 0 + \sin 0 = 1$

$$\text{When } x = \frac{\pi}{2}, t = \cos \frac{\left(\frac{\pi}{2}\right)}{2} + \sin \frac{\left(\frac{\pi}{2}\right)}{2} = \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}} = \sqrt{2}$$

So, the new limits are 1 and $\sqrt{2}$.

Substituting this in the original integral,

$$I = \int_1^{\sqrt{2}} \frac{1}{t^{n-1}} (2dt)$$

$$\Rightarrow I = 2 \int_1^{\sqrt{2}} t^{-(n-1)} dt$$

$$\text{Recall } \int x^n dx = \frac{1}{n+1} x^{n+1} + c$$

$$\Rightarrow I = 2 \left[\frac{t^{-(n-1)+1}}{-(n-1) + 1} \right]_1^{\sqrt{2}}$$

$$\Rightarrow I = \frac{2}{2-n} [t^{2-n}]_1^{\sqrt{2}}$$

$$\Rightarrow I = \frac{2}{2-n} \left[(\sqrt{2})^{2-n} \right]_{-1^{2-n}}$$

$$\Rightarrow I = \frac{2}{2-n} \left[\left(2^{\frac{1}{2}} \right)^{2-n} \right]_{-1} = \frac{2}{2-n} \left(2^{1-\frac{n}{2}} - 1 \right)$$

$$\therefore \int_0^{\frac{\pi}{2}} \frac{\cos x}{(\cos \frac{x}{2} + \sin \frac{x}{2})^n} dx = \frac{2}{2-n} \left(2^{1-\frac{n}{2}} - 1 \right)$$

Exercise 20.3

1 A. Question

Evaluate the following Integrals:

$$\int_1^4 f(x) dx, \text{ where } f(x) = \begin{cases} 4x+3, & \text{if } 1 \leq x \leq 2 \\ 3x+5, & \text{if } 2 \leq x \leq 4 \end{cases}$$

Answer

$$\begin{aligned} \text{We have, } & \int_1^4 f(x) dx \\ &= \int_1^2 (4x+3) dx + \int_2^4 (3x+5) dx \\ &= \left[\frac{4x^2}{2} + 3x \right]_1^2 + \left[\frac{3x^2}{2} + 5x \right]_2^4 \\ &= \left[\left(\frac{16}{2} + 6 \right) - \left(\frac{4}{2} + 3 \right) \right] + \left[\left(\frac{48}{2} + 20 \right) - \left(\frac{12}{2} + 10 \right) \right] \\ &= [14 - 5] + [44 - 16] \\ &= 9 + 28 \\ &= 37 \end{aligned}$$

1 B. Question

Evaluate the following Integrals:

$$\int_0^9 f(x) dx, \text{ where } f(x) = \begin{cases} \sin x, & 0 \leq x \leq \pi/2 \\ 1, & \pi/2 \leq x \leq 3 \\ e^{x-3}, & 3 \leq x \leq 9 \end{cases}$$

Answer

$$\begin{aligned} \text{We have, } & \int_0^9 f(x) dx \\ &= \int_0^{\frac{\pi}{2}} \sin x dx + \int_{\frac{\pi}{2}}^3 1 dx + \int_3^9 e^{x-3} dx \\ &= [-\cos x]_0^{\frac{\pi}{2}} + [x]_{\frac{\pi}{2}}^3 + [e^{x-3}]_3^9 \end{aligned}$$

$$= \left[-\cos \frac{\pi}{2} + \cos 0 \right] + \left[3 - \frac{\pi}{2} \right] + [e^{9-3} - e^{3-3}]$$

$$= [0 + 1] + \left[3 - \frac{\pi}{2} \right] + [e^6 - e^0]$$

$$= 1 + 3 - \frac{\pi}{2} + e^6 - 1$$

$$\text{Hence, } 3 - \frac{\pi}{2} + e^6$$

1 C. Question

Evaluate the following Integrals:

$$\int_1^4 f(x) dx, \text{ where } f(x) = \begin{cases} 7x + 3, & \text{if } 1 \leq x \leq 3 \\ 8x, & \text{if } 3 \leq x \leq 4 \end{cases}$$

Answer

$$\text{We have, } \int_1^4 f(x) dx$$

$$= \int_1^3 (7x + 3) + \int_3^4 8x dx$$

$$= \left[\frac{7x^2}{2} + 3x \right]_1^3 + \left[\frac{8x^2}{2} \right]_3^4$$

$$= \left[\frac{7 \times 9}{2} + 3 \times 3 \right] - \left[\frac{7 \times 1}{2} + 3 \times 1 \right] + \left[\left(\frac{8 \times 16}{2} \right) - \left(\frac{8 \times 9}{2} \right) \right]$$

$$= \left[\frac{63}{2} + 9 - \frac{7}{2} + 3 \right] + [64 - 36]$$

$$= 34 + 28$$

$$\text{Hence, } 62$$

2. Question

Evaluate the following Integrals:

$$\int_{-4}^4 |x + 2| dx$$

Answer

We have,

$$= \int_{-4}^4 |x + 2| dx$$

$$= \int_{-4}^{-2} -(x + 2) dx + \int_{-2}^4 (x + 2) dx$$

$$= - \left[\frac{x^2}{2} + 2x \right]_{-4}^{-2} + \left[\frac{x^2}{2} + 2x \right]_{-2}^4$$

$$= - \left[\frac{4}{2} - 4 \right] - \left[\frac{16}{2} - 8 \right] + \left[\left(\frac{16}{2} + 8 \right) - \left(\frac{4}{2} - 4 \right) \right]$$

$$= -[2 - 4] - [8 - 8] + [(8 + 8) - (2 - 4)]$$

$$= -[-2] + 16 + 2$$

$$= 2 + 2 + 16$$

$$\text{Hence, } 20$$

3. Question

Evaluate the following Integrals:

$$\int_{-3}^3 |x + 1| dx$$

Answer

We have, $\int_{-3}^3 |x + 1| dx$

$$\begin{aligned}&= \int_{-3}^3 |x + 1| dx \\&= \int_{-3}^{-1} -(x + 1) dx + \int_{-1}^3 (x + 1) dx \\&= -\left[\frac{x^2}{2} + x\right]_{-3}^{-1} + \left[\frac{x^2}{2} + x\right]_{-1}^3 \\&= -\left[\frac{1}{2} - 1\right] + \left[\frac{9}{2} - 3\right] + \left[\left(\frac{9}{2} + 3\right) - \left(\frac{1}{2} - 1\right)\right] \\&= -\left[-\frac{1}{2}\right] + \left[\frac{3}{2}\right] + \left[\left(\frac{15}{2}\right) - \left(-\frac{1}{2}\right)\right] \\&= \frac{1}{2} + \frac{3}{2} + \frac{16}{2} \\&= \frac{20}{2} = 10\end{aligned}$$

Hence, $\int_{-3}^3 |x + 1| dx = 10$

4. Question

Evaluate the following Integrals:

$$\int_{-1}^1 |2x + 1| dx$$

Answer

We have, $\int_{-1}^1 |2x + 1| dx$

$$\begin{aligned}&= \int_{-1}^1 |2x + 1| dx \\&= \int_{-1}^{-\frac{1}{2}} -(2x + 1) dx + \int_{-\frac{1}{2}}^1 (2x + 1) dx \\&= -\left[\frac{2x^2}{2} + x\right]_{-1}^{-\frac{1}{2}} + \left[\frac{2x^2}{2} + x\right]_{-\frac{1}{2}}^1 \\&= -\left[\frac{2}{8} - \frac{1}{2}\right] - \left[\frac{2}{2} - 1\right] + \left[\left(\frac{2}{2} + 1\right) - \left(\frac{2}{8} - \frac{1}{2}\right)\right] \\&= -\left[\frac{1}{4} - \frac{1}{2}\right] - [1 - 1] + \left[(1 + 1) - \left(\frac{1}{4} - \frac{1}{2}\right)\right] \\&= -\left[-\frac{1}{4}\right] + \left[2 + \left(\frac{1}{4}\right)\right] \\&= \frac{1}{4} + 2 + \frac{1}{4} \\&= \frac{2}{4} + 2\end{aligned}$$

Hence, $\int_{-1}^1 |2x + 1| dx = \frac{5}{2}$

5. Question

Evaluate the following Integrals:

$$\int_{-2}^2 |2x + 3| dx$$

Answer

We have, $\int_{-2}^2 |2x + 3| dx$

$$= \int_{-2}^2 |2x + 3| dx$$

$$= \int_{-2}^{\frac{3}{2}} -(2x + 3) dx + \int_{\frac{3}{2}}^2 (2x + 3) dx$$

$$= -\left[\frac{2x^2}{2} + 3x\right]_{-2}^{\frac{3}{2}} + \left[\frac{2x^2}{2} + 3x\right]_{\frac{3}{2}}^2$$

$$= -\left[\frac{2 \times 9}{2 \times 4} - \frac{9}{2}\right] - \left[\frac{2 \times 4}{2} - 6\right] + \left[\left(\frac{2 \times 4}{2} + 6\right) - \left(\frac{2 \times 9}{2 \times 4} - \frac{9}{2}\right)\right]$$

$$= -\left[\frac{18}{8} - \frac{9}{2}\right] - \left[\frac{8}{2} - 6\right] + \left[\left(\frac{8}{2} + 6\right) - \left(\frac{18}{8} - \frac{9}{2}\right)\right]$$

$$= -\left[\frac{9}{4} - \frac{9}{2}\right] - [-2] + \left[(10) - \left(\frac{9}{4} - \frac{9}{2}\right)\right]$$

$$= -\left[-\frac{9}{4} + 2\right] + \left[(10) + \frac{9}{4}\right]$$

$$= \frac{9}{4} - 2 + 10 + \frac{9}{4}$$

$$= \frac{9}{2} + 8$$

Hence, $\int_{-2}^2 |2x + 3| dx = \frac{25}{2}$

6. Question

Evaluate the following Integrals:

$$\int_0^2 |x^2 - 3x + 2| dx$$

Answer

We have, $\int_0^2 |x^2 - 3x + 2| dx$

$$= \int_0^2 |x^2 - 3x + 2| dx$$

$$= \int_0^1 (x^2 - 3x + 2) dx + \int_1^2 -(x^2 - 3x + 2) dx$$

$$= \left[\frac{x^3}{3} - \frac{3x^2}{2} + 2x\right]_0^1 - \left[\frac{x^3}{3} - \frac{3x^2}{2} + 2x\right]_1^2$$

$$= \left[\frac{1}{3} - \frac{3}{2} + 2 - 0\right] - \left[\left(\frac{8}{3} - \frac{12}{2} + 4 - \frac{1}{3} + \frac{3}{2} + 2\right)\right]$$

$$= \left[\frac{1}{6}\right] - \left[-\frac{5}{6}\right]$$

$$= \frac{1}{6} + \frac{5}{6}$$

$$= \frac{6}{6} = 1$$

Hence, $\int_0^2 |x^2 - 3x + 2| dx = 1$

7. Question

Evaluate the following Integrals:

$$\int_0^3 |3x - 1| dx$$

Answer

We have, $\int_0^3 |3x - 1| dx$

$$= \int_0^3 |3x - 1| dx$$

$$= \int_0^{\frac{1}{3}} -(3x - 1) dx + \int_{\frac{1}{3}}^3 (3x - 1) dx$$

$$= -\left[\frac{3x^2}{2} - x\right]_0^{\frac{1}{3}} + \left[\frac{3x^2}{2} - x\right]_{\frac{1}{3}}^3$$

$$= -\left[\frac{3}{9 \times 2} - \frac{1}{3} - 0\right] + \left[\left(\frac{3 \times 9}{2} - 3\right) - \left(\frac{3}{9 \times 2}\right)\right]$$

$$= -\left[\frac{1}{6} - \frac{1}{3}\right] + \left[\left(\frac{27}{2} - 3\right) - \left(\frac{1}{6} - \frac{1}{3}\right)\right]$$

$$= -\left[-\frac{1}{6}\right] + \left[\left(\frac{21}{2}\right) - \left(-\frac{1}{6}\right)\right]$$

$$= \left[\frac{1}{6}\right] + \left[\left(\frac{21}{2}\right) + \left(\frac{1}{6}\right)\right]$$

$$= \frac{2}{6} + \frac{21}{2}$$

$$= \frac{1}{3} + \frac{21}{2} = \frac{2 + 63}{6}$$

$$= \frac{65}{6}$$

Hence, $\int_0^3 |3x - 1| dx = \frac{65}{6}$

8. Question

Evaluate the following Integrals:

$$\int_{-6}^6 |x + 2| dx$$

Answer

We have, $\int_{-6}^6 |x + 2| dx$

$$= \int_{-6}^2 -(x + 2) + \int_2^6 (x + 2) dx$$

$$= -\left[\frac{x^2}{2} + 2x\right]_{-6}^{-2} + \left[\frac{x^2}{2} + 2x\right]_2^6$$

$$\begin{aligned}
&= - \left[\left(\frac{4}{2} + 2 \times 2 \right) - \left(\frac{36}{2} - 12 \right) \right] + \left[\left(\frac{36}{2} + 12 \right) - \left(\frac{4}{2} - 4 \right) \right] \\
&= -[(2 - 4) - (18 - 12)] + [(18 + 12) - (2 - 4)] \\
&= -(-8) + (30 + 2) \\
&= 8 + 32 \\
&= 40
\end{aligned}$$

Hence, $\int_{-6}^6 |x + 2| dx = 40$

9. Question

Evaluate the following Integrals:

$$\int_{-2}^2 |x + 1| dx$$

Answer

$$\begin{aligned}
\text{We have, } &\int_{-2}^2 |x + 1| dx \\
&= \int_{-2}^{-1} -(x + 1) dx + \int_{-1}^2 (x + 1) dx \\
&= - \left[\frac{x^2}{2} + x \right]_{-2}^{-1} + \left[\frac{x^2}{2} + x \right]_{-1}^2 \\
&= - \left[\left(\frac{1}{2} - 1 \right) - \left(\frac{4}{2} - 2 \right) \right] + \left[\left(\frac{4}{2} + 2 \right) - \left(\frac{1}{2} - 1 \right) \right] \\
&= - \left[\left(-\frac{1}{2} \right) - 0 \right] + \left[4 + \frac{1}{2} \right] \\
&= \frac{1}{2} + \frac{9}{2} \\
&= \frac{10}{2}
\end{aligned}$$

Hence, $\int_{-2}^2 |x + 1| dx = 5$

10. Question

Evaluate the following Integrals:

$$\int_1^2 |x - 3| dx$$

Answer

$$\begin{aligned}
\text{We have, } &\int_1^2 |x - 3| dx \\
&= \int_1^2 -(x - 3) dx \\
&= [x - 3 < 0 \text{ for } 1 > x > 2] \\
&= - \left[\frac{x^2}{2} - 3x \right]_1^2 \\
&= - \left[\left(\frac{4}{2} - 6 \right) - \left(\frac{1}{2} - 3 \right) \right] \\
&= - \left[(-4) - \left(-\frac{5}{2} \right) \right]
\end{aligned}$$

$$= - \left[-4 + \frac{5}{2} \right]$$

$$= - \left[-\frac{3}{2} \right]$$

$$= \frac{3}{2}$$

$$\text{Hence, } \int_1^2 |x - 3| dx = \frac{3}{2}$$

11. Question

Evaluate the following Integrals:

$$\int_0^{\pi/2} |\cos 2x| dx$$

Answer

$$\text{We have, } \int_0^{\pi/2} |\cos 2x| dx$$

$$= \int_0^{\pi/2} |\cos 2x| dx$$

$$= \int_0^{\pi/4} -(\cos 2x) dx + \int_{\pi/4}^{\pi/2} (\cos 2x) dx$$

$$= \left[\frac{\sin 2x}{2} \right]_0^{\pi/4} + \left[\frac{-\sin 2x}{2} \right]_{\pi/4}^{\pi/2}$$

$$= \frac{1}{2} \left[\sin \frac{\pi}{2} - \sin 0 \right] + \frac{1}{2} [\sin \pi + \sin \frac{\pi}{2}]$$

$$= \frac{1}{2}(1) + \frac{1}{2}(1)$$

$$= \frac{1}{2} + \frac{1}{2}$$

$$= \frac{2}{2} = 1$$

$$\text{Hence, } \int_0^{\pi/2} |\cos 2x| dx = 1$$

12. Question

Evaluate the following Integrals:

$$\int_0^{2\pi} |\sin x| dx$$

Answer

$$\text{We have, } \int_0^{\pi} |\sin x| dx$$

$$= \int_0^{\pi} \sin x dx = \int_0^{\pi} \sin x dx + \int_{\pi}^{2\pi} -\sin x dx$$

$$= [-\cos x]_0^{\pi} + [\cos x]_{\pi}^{2\pi}$$

$$= [-\cos \pi + \cos 0] + [\cos 2\pi - \cos \pi]$$

$$= [1 + 1] + [1 + 1]$$

$$= 2 + 2$$

= 4

Hence, $\int_0^{\frac{\pi}{2}} |\sin x| dx = 4$

13. Question

Evaluate the following Integrals:

$$\int_{-\pi/4}^{\pi/4} |\sin x| dx$$

Answer

We have, $\int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} |\sin x| dx$

$$= \int_{-\frac{\pi}{4}}^0 -\sin x dx + \int_0^{\frac{\pi}{4}} \sin x dx$$

$$= [\cos x]_{-\frac{\pi}{4}}^0 + [-\cos x]_0^{\frac{\pi}{4}}$$

$$= [\cos 0 + \cos(-\frac{\pi}{4})] + [-\cos \frac{\pi}{4} + \cos 0]$$

$$= \left[\left(1 - \frac{1}{\sqrt{2}}\right) - \left(\frac{1}{\sqrt{2}} - 1\right) \right]$$

$$= (2 - \sqrt{2})$$

Hence, $\int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} |\sin x| dx = (2 - \sqrt{2})$

14. Question

Evaluate the following Integrals:

$$\int_2^8 |x - 5| dx$$

Answer

We have,

$$\int_2^8 |x - 5| dx$$

We have,

$$|x - 5| = \begin{cases} x - 5 & \text{if } x \in (5, 8) \\ -(x - 5) & \text{if } x \in (2, 5) \end{cases}$$

Hence,

$$= \int_2^5 -(x - 5) dx + \int_5^8 (x - 5) dx$$

$$= -\left[\frac{x^2}{2} - 5x\right]_2^5 + \left[\frac{x^2}{2} - 5x\right]_5^8$$

$$= -\left[\left(\frac{25}{2} - 25\right) - \left(\frac{4}{2} - 10\right) + \left(\frac{64}{2} - 40\right) - \left(\frac{25}{2} - 25\right)\right]$$

$$= -\left[-\frac{25}{2} + 8\right] + \left[(-8) + \left(\frac{25}{2}\right)\right]$$

$$= \frac{25}{2} + \frac{25}{2} - 8 - 8$$

$$= 25 - 16 = 9$$

Hence, $\int_2^8 |x - 5| dx = 9$

15. Question

Evaluate the following Integrals:

$$\int_{-\pi/2}^{\pi/2} (\sin|x| + \cos|x|) dx$$

Answer

We have,

$$I = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} (\sin|x| + \cos|x|) dx$$

Let $f(x) = \sin|x| + \cos|x|$

Then, $f(x) = f(-x)$

Since, $(f(x))$ is an even function.

$$\text{So, } I = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} (\sin|x| + \cos|x|) dx$$

$$= 2 \int_0^{\frac{\pi}{2}} (\sin x + \cos x) dx$$

$$= 2[-\cos x + \sin x]_0^{\frac{\pi}{2}}$$

$$= 2[-\cos \frac{\pi}{2} + \sin \frac{\pi}{2} + \cos 0 - \sin 0]$$

$$= 2[0 + 1 + 1 - 0]$$

$$= 2(2)$$

Hence, $\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} (\sin|x| + \cos|x|) dx = 4$

16. Question

Evaluate the following Integrals:

$$\int_0^4 |x - 1| dx$$

Answer

We have, $\int_0^4 |x - 1| dx$

It can be seen that $(x-1) \leq 0$ when $0 \leq x \leq 1$ and $(x-1) \geq 0$ when $1 \leq x \leq 4$

$$= I = \int_1^4 |x - 1| dx$$

$$= \int_0^1 -(x - 1) dx + \int_1^4 (x - 1) dx$$

$$\begin{aligned}
&= -\left[\frac{x^2}{2} - x\right]_0^1 + \left[\frac{x^2}{2} - x\right]_1^4 \\
&= -\left[\left(\frac{1}{2} - 1 - 0\right) - \left(\frac{16}{2} - 4 - \frac{1}{2} + 1\right)\right] \\
&= \left[1 - \frac{1}{2} + 8 - 4 - \frac{1}{2} + 1\right] \\
&= \left[\frac{1}{2} + 4 - \frac{1}{2} + 1\right] \\
&= 5
\end{aligned}$$

Hence, $\int_1^4 |x - 1| dx = 5$

17. Question

Evaluate the following Integrals:

$$\int_1^4 \{|x - 1| + |x - 2| + |x - 4|\} dx$$

Answer

$$\begin{aligned}
\text{Let } I &= \int_1^4 \{|x - 1| + |x - 2| + |x - 4|\} dx \\
&= \int_1^2 \{(x - 1 - x + 2 - x + 4)\} dx + \int_2^4 \{(x - 1 + x - 2 - x + 4)\} dx \\
&= \int_1^2 (5 - x) dx + \int_2^4 (x + 1) dx \\
&= \left[5x - \frac{x^2}{2}\right]_1^2 + \left[\frac{x^2}{2} + x\right]_2^4 \\
&= \left[10 - \frac{4}{2} - 5 + \frac{1}{2}\right] + \left[\frac{16}{2} + 4 - \frac{4}{2} - 2\right] \\
&= \left[5 - \frac{3}{2}\right] + [8 + 4 - 4] \\
&= \left[\frac{7}{2}\right] + 8
\end{aligned}$$

Hence, $\int_1^4 \{|x - 1| + |x - 2| + |x - 4|\} dx = \frac{23}{2}$

18. Question

Evaluate the following Integrals:

$$\int_{-5}^0 f(x) dx, \text{ where } f(x) = |x| + |x + 2| + |x + 5|$$

Answer

We have,

$$\begin{aligned}
I &= \int_{-5}^0 \{|x| + |x + 2| + |x + 5|\} dx \\
&= \int_{-5}^0 \{|x| dx + \int_{-5}^0 |x + 2| dx + \int_{-5}^0 |x + 5| dx\} \\
&= I = \int_{-5}^0 -x dx + \int_{-5}^{-2} -(x + 2) dx + \int_{-2}^0 (x + 2) dx + \int_{-5}^0 |x + 5| dx \\
&= \left[-\frac{x^2}{2}\right]_{-5}^0 + \left[-\frac{x^2}{2} - 2x\right]_{-5}^{-2} + \left[\frac{x^2}{2} + 2x\right]_{-2}^0 + \left[\frac{x^2}{2} + 5x\right]_{-5}^0
\end{aligned}$$

$$\begin{aligned}
&= \left[0 + \frac{25}{2} \right] - \left[\frac{4}{2} + 4 - \frac{25}{2} + 10 \right] + \left[0 - \frac{4}{2} + 4 \right] + \left[0 - \frac{25}{2} + 25 \right] \\
&= \frac{25}{2} - \left[8 - \frac{25}{2} \right] + (2) + \left[25 - \frac{25}{2} \right] \\
&= \frac{25}{2} - 8 + \frac{25}{2} + 2 + 25 - \frac{25}{2} \\
&= 19 + \frac{25}{2}
\end{aligned}$$

Hence, $\int_{-5}^0 (|x| + |x+2| + |x+5|) dx = \frac{62}{2}$

19. Question

Evaluate the following Integrals:

$$\int_0^4 (|x| + |x-2| + |x-4|) dx$$

Answer

We have

$$\begin{aligned}
I &= \int_0^4 (|x| + |x-2| + |x-4|) dx \\
&= \int_0^2 (|x| + |x-2| + |x-4|) dx + \int_2^4 (|x| + |x-2| + |x-4|) dx \\
&= \int_0^2 (x + 2 - x + 4 - x) dx + \int_2^4 (x + x - 2 + 4 - x) dx \\
&= \int_0^2 (6 - x) dx + \int_2^4 (2 + x) dx \\
&= \left[6x - \frac{x^2}{2} \right]_0^2 + \left[2x + \frac{x^2}{2} \right]_2^4 \\
&= [12 - 2 - 0 - 0] + [8 + 8 - 4 - 2] \\
&= [10 + 10] \\
&= 20
\end{aligned}$$

Hence, $\int_0^4 (|x| + |x-2| + |x-4|) dx = 20$

20. Question

Evaluate the following Integrals:

$$\int_{-1}^2 (|x+1| + |x| + |x-1|) dx$$

Answer

We have, $\int_{-1}^2 (|x+1| + |x| + |x-1|) dx$

Now, we can write as

$$\begin{aligned}
&\int_{-1}^2 |x+1| dx \int_{-1}^2 |x| dx + \int_{-1}^2 |x-1| dx \\
&\int_{-1}^2 (x+1) dx - \int_{-1}^0 x dx + \int_0^2 x dx - \int_{-1}^1 (x-1) dx + \int_1^2 (x-1) dx
\end{aligned}$$

$$\left\{ \frac{x^2}{2} + x \right\}_{-1}^2 - \left\{ \frac{x^2}{2} \right\}_{-1}^0 + \left\{ \frac{x^2}{2} \right\}_0^2 - \left\{ \frac{x^2}{2} - x \right\}_{-1}^1 + \left\{ \frac{x^2}{2} - x \right\}_1^2$$

While putting the limits

$$\left\{ \left(\frac{4}{2} + 2 - \frac{1}{2} - 1 \right) - \left(0 - \frac{1}{2} \right) + \left(\frac{4}{2} - 0 \right) - \left(\frac{1}{2} - 1 + \frac{1}{2} + 1 \right) + \left(\frac{4}{2} - 2 - \frac{1}{2} + 1 \right) \right\}$$

$$\left\{ \left(4 - \left(-\frac{1}{2} \right) \right) - \left(-\frac{1}{2} \right) + (2) - \left(-\frac{1}{2} - \frac{3}{2} \right) + \left(2 - 2 + \frac{1}{2} \right) \right\}$$

$$\left\{ \left(4 + \frac{1}{2} \right) + \frac{1}{2} + (2) - \left(-\frac{1}{2} - \frac{3}{2} \right) + \left(0 + \frac{1}{2} \right) \right\}$$

$$\left\{ \left(4 + \frac{1}{2} \right) + \frac{1}{2} + (2) - \left(-\frac{1}{2} - \frac{3}{2} \right) + \left(0 + \frac{1}{2} \right) \right\}$$

$$\left\{ \left(\frac{9}{2} \right) + \frac{1}{2} + (2) + (2) + \left(\frac{1}{2} \right) \right\}$$

$$\left\{ \frac{11}{2} + (4) \right\}$$

Hence, $\frac{19}{2}$

21. Question

Evaluate the following Integrals:

$$\int_{-2}^2 x e^{|x|} dx$$

Answer

$$\begin{aligned} & \int_{-2}^2 x e^{|x|} dx \\ &= \int_{-2}^0 x e^{-x} dx + \int_0^2 x e^x dx \end{aligned}$$

Let's Say $I = I_1 + I_2$

$$\text{And, } I_1 = \int_{-2}^0 x e^{-x} dx$$

Using Integration By parts

$$\int f'g = fg - \int fg'$$

$$f' = e^{-x}, g = x$$

$$f = -e^{-x}, g' = 1$$

$$\int_{-2}^0 x e^{-x} dx = \{-xe^{-x}\}_{-2}^0 - \int_{-2}^0 e^{-x} dx$$

$$\int_{-2}^0 x e^{-x} dx = \{-xe^{-x} - e^{-x}\}_{-2}^0$$

$$\int_{-2}^0 x e^{-x} dx = \{(-1) - (2e^2 - e^0)\}$$

$$I_1 = \int_{-2}^0 x e^{-x} dx = \{-1 - e^2\}$$

$$\text{For } I_2 = \int_0^2 x e^x dx$$

Using Integration By parts

$$\int f'g = fg - \int fg'$$

$$f' = e^x, g = x$$

$$f = e^x, g' = 1$$

$$\int_0^2 x e^x dx = \{xe^x\}_0^2 - \int_0^2 e^x dx$$

$$\int_0^2 x e^x dx = \{xe^x - e^x\}_0^2$$

$$\int_0^2 x e^x dx = \{(2e^2 - e^2 + 1)\}$$

$$I_2 = \int_0^2 x e^x dx = e^2 + 1$$

$$\text{Now, } I = I_1 + I_2$$

$$\int_{-2}^0 x e^{|x|} dx = -1 - e^2 + e^2 + 1 = 0$$

Hence, 0

22. Question

Evaluate the following Integrals:

$$\int_{-\frac{\pi}{4}}^{\frac{\pi}{2}} (\sin x |\sin x|) dx$$

Answer

$$\begin{aligned} & - \int_{-\frac{\pi}{4}}^0 \sin^2 x dx + \int_0^{\frac{\pi}{2}} \sin^2 x dx \\ &= \sin^2 x = \frac{1 - \cos 2x}{2} \\ &= - \int_{-\frac{\pi}{4}}^0 \frac{1 - \cos 2x}{2} dx + \int_0^{\frac{\pi}{2}} \frac{1 - \cos 2x}{2} dx \\ &= -\frac{1}{2} \left\{ x - \frac{\sin 2x}{2} \right\}_{-\frac{\pi}{4}}^0 + \frac{1}{2} \left\{ x + \frac{\sin 2x}{2} \right\}_0^{\frac{\pi}{2}} \\ &= -\frac{1}{2} \left\{ -0 \left(-\frac{\pi}{4} + \frac{1}{2} \right) \right\} + \frac{1}{2} \left\{ \frac{\pi}{2} - 0 \right\} \\ &= \left\{ -\frac{\pi}{8} + \frac{1}{4} \right\} + \left\{ \frac{\pi}{4} \right\} \end{aligned}$$

$$= \frac{\pi}{8} + \frac{1}{4}$$

$$= \frac{\pi+2}{8}$$

23. Question

Evaluate the following Integrals:

$$\int_0^{\pi} (\cos x |\cos x|) dx$$

Answer

$$\int_0^{\pi/2} \cos^2 x dx - \int_{\pi/2}^{\pi} \cos^2 x dx$$

$$= \cos^2 x = \frac{1 + \cos 2x}{2}$$

$$= \int_0^{\frac{\pi}{2}} \frac{1 + \cos 2x}{2} dx - \int_{\frac{\pi}{2}}^{\pi} \frac{1 + \cos 2x}{2} dx$$

$$= \frac{1}{2} \left\{ \frac{1 + \cos 2x}{2} \right\}_{0}^{\frac{\pi}{2}} - \frac{1}{2} \left\{ x + \frac{\sin 2x}{2} \right\}_{\frac{\pi}{2}}^{\pi}$$

$$= \frac{\pi}{4} - \frac{\pi}{4}$$

$$= 0$$

24. Question

Evaluate the following Integrals:

$$\int_{-\pi/4}^{\pi/2} (2\sin|x| + \cos|x|) dx$$

Answer

$$\int_{-\pi/4}^{\pi/2} (2\sin|x| + \cos|x|) dx$$

$$= \int_{-\frac{\pi}{4}}^0 (-2\sin x + \cos x) dx + \int_0^{\frac{\pi}{2}} (2\sin x + \cos x) dx$$

$$= [2\cos x + \sin x]_{-\frac{\pi}{4}}^0 + [-2\cos x + \sin x]_0^{\frac{\pi}{2}}$$

$$= [2\cos(0) + \sin(0) - 2\cos(-\frac{\pi}{4}) - \sin(-\frac{\pi}{4})] + [-2\cos(\frac{\pi}{2}) + \sin(\frac{\pi}{2}) + 2\cos(0) - \sin(0)]$$

$$= [2 + 0 - 0 + 1] + [0 + 1 + 2 - 0]$$

$$= 6$$

25. Question

Evaluate the following Integrals:

$$\int_{-\pi/2}^{\pi} \sin^{-1}(\sin x) dx$$

Answer

$$\begin{aligned}
\int_{-\pi/2}^{\pi} \sin^{-1}(\sin x) dx &= \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} x dx + \int_{\frac{\pi}{2}}^{\pi} (\pi - x) dx \\
&= \left\{ \frac{x^2}{2} \right\}_{-\frac{\pi}{2}}^{\frac{\pi}{2}} + \left\{ \pi x - \frac{x^2}{2} \right\}_{\frac{\pi}{2}}^{\pi} \\
&= \left\{ \left(\pi^2 - \frac{\pi^2}{2} \right) - \left(\frac{\pi^2}{2} - \frac{\pi^2}{8} \right) \right\} \\
&= \left\{ \left(\frac{2\pi^2 - \pi^2}{2} \right) - \left(\frac{4\pi^2 - \pi^2}{8} \right) \right\} \\
&= \left\{ \left(\frac{\pi^2}{2} - \frac{3\pi^2}{8} \right) \right\} \\
&= \left(\frac{4\pi^2 - 3\pi^2}{8} \right) \\
&= \frac{\pi^2}{8}
\end{aligned}$$

26. Question

Evaluate the following Integrals:

$$\int_{-\pi/2}^{\pi/2} \frac{-\pi/2}{\sqrt{\cos x \sin^2 x}} dx$$

Answer

$$\text{Let } f(x) = \frac{-\frac{\pi}{2}}{\sqrt{\cos x \sin^2 x}}$$

$$f(-x) = f(x)$$

And thus $f(x)$ is an even function.

So,

$$\int_{-\pi/2}^{\pi/2} \frac{-\frac{\pi}{2}}{\sqrt{\cos x \sin^2 x}} dx = 2 \int_0^{\pi/2} \frac{-\frac{\pi}{2}}{\sqrt{\cos x \sin^2 x}} dx$$

$$\int_{-\pi/2}^{\pi/2} \frac{-\frac{\pi}{2}}{\sqrt{\cos x \sin^2 x}} dx = 2 \int_0^{\pi/2} \frac{-\frac{\pi}{2}}{\sin x \sqrt{\cos x}} dx$$

$$\text{Let } \cos x = t$$

Differentiating both sides we get,

$$-\sin x dx = dt$$

$$-\sqrt{1-t^2} dx = dt$$

Limits will also change,

$$\text{At } x = 0, t = 1 \text{ and at } x = \pi/2, t = 0$$

Now the Expression becomes,

$$\int_{-\pi/2}^{\pi/2} \frac{-\frac{\pi}{2}}{\sqrt{\cos x \sin^2 x}} dx = \int_1^0 \frac{\pi}{\sqrt{t}\sqrt{1-t^2}} dt$$

$$\int_{\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{-\frac{\pi}{2}}{\sqrt{\cos x \sin^2 x}} dx = \int_1^0 \frac{\pi}{\sqrt{t-t^3}} dt$$

$$\int_{\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{-\frac{\pi}{2}}{\sqrt{\cos x \sin^2 x}} dx = 1 + \frac{1}{3} = \frac{4}{3}$$

27. Question

Evaluate the following Integrals:

$$\int_0^2 2x[x] dx$$

Answer

$[x] = 0$ for 0

and $[x] = 1$ for 1

Hence

$$= \int_0^1 0 + \int_1^2 2x dx$$

$$= 0 + \left[\frac{2x^2}{2} \right]_1^2$$

$$= [x^2]_1^2$$

$$= (2^2 - 1^2)$$

$$= 4 - 1$$

$$= 3.$$

28. Question

Evaluate the following Integrals:

$$\int_0^{2\pi} \cos^{-1}(\cos x) dx$$

Answer

$$\int_0^{2\pi} \cos^{-1}(\cos x) dx$$

$$= - \int_0^\pi \cos^{-1}(\cos x) dx + \int_0^{2\pi} \cos^{-1}(\cos x) dx$$

$$= - \int_0^\pi x dx + \int_0^{2\pi} x dx$$

$$= - \left[\frac{x^2}{2} \right]_0^\pi + \left[\frac{x^2}{2} \right]_\pi^{2\pi}$$

$$= - \frac{\pi^2}{2} + \frac{4\pi^2}{2} - \frac{\pi^2}{2}$$

$$= \pi^2$$

Exercise 20.4

1. Question

Evaluate of each of the following integral:

$$\int_0^{2\pi} \frac{e^{\sin x}}{e^{\sin x} + e^{-\sin x}} dx$$

Answer

let us assume $I = \int_0^{2\pi} \frac{e^{\sin x}}{e^{\sin x} + e^{-\sin x}} dx \dots\dots \text{equation 1}$

By property, we know that $\int_a^b f(x) dx = \int_a^b f(a+b-x) dx$

$$I = \int_0^{2\pi} \frac{e^{\sin(2\pi-x)}}{e^{\sin(2\pi-x)} + e^{-\sin(2\pi-x)}} dx \dots\dots \text{equation 2}$$

Adding equation 1 and 2

$$2I = \int_0^{2\pi} \frac{e^{\sin x}}{e^{\sin x} + e^{-\sin x}} dx + \int_0^{2\pi} \frac{e^{\sin(2\pi-x)}}{e^{\sin(2\pi-x)} + e^{-\sin(2\pi-x)}} dx$$

We know $\sin(2\pi - x) = -\sin x$

Thus

$$2I = \int_0^{2\pi} \frac{e^{\sin x}}{e^{\sin x} + e^{-\sin x}} dx + \int_0^{2\pi} \frac{e^{-\sin x}}{e^{-\sin x} + e^{\sin x}} dx$$

We know

$$\int_m^n [f(x) + g(x)] dx = \int_m^n f(x) dx + \int_m^n g(x) dx$$

thus

$$2I = \int_0^{2\pi} \frac{e^{\sin x} + e^{-\sin x}}{e^{\sin x} + e^{-\sin x}} dx$$

$$2I = \int_0^{2\pi} 1 dx$$

$$2I = [x]_0^{2\pi}$$

We know

$$[f(x)]_a^b = f(b) - f(a)$$

$$2I = [2\pi - 0]$$

$$I = \pi$$

2. Question

Evaluate of each of the following integral:

$$\int_0^{2\pi} \log(\sec x + \tan x) dx$$

Answer

Let us assume $I = \int_0^{2\pi} \log(\sec x + \tan x) dx \dots\dots \text{equation 1}$

By property, we know that $\int_a^b f(x) dx = \int_a^b f(a+b-x) dx$

$$I = \int_0^{2\pi} \log(\sec(2\pi - x) + \tan(2\pi - x)) dx$$

We know that $\sec(2\pi - x) = \sec(x)$

$\tan(2\pi - x) = -\tan(x)$

thus $I = \int_0^{2\pi} \log(\sec(x) - \tan(x)) dx$ equation 2

Adding equations 1 and equation 2, we get,

$$2I = \int_0^{2\pi} \log(\sec x + \tan x) dx + \int_0^{2\pi} \log(\sec x - \tan x) dx$$

We know $\log(a) + \log(b) = \log(ab)$

$$\int_m^n [f(x) + g(x)] dx = \int_m^n f(x) dx + \int_m^n g(x) dx$$

Thus

$$2I = \int_0^{2\pi} \log(\sec x + \tan x)(\sec x - \tan x) dx$$

$$2I = \int_0^{2\pi} \log(\sec^2 x - \tan^2 x) dx$$

We know Trigonometric identity $\sec^2 x - \tan^2 x = 1$

$$2I = \int_0^{2\pi} \log(1) dx$$

We know

$[f(x)]_a^b = f(b) - f(a)$ where b is the upper limit and a is lower and $f(x)$ is integral function

$$2I = [0]_0^{2\pi}$$

Thus $I = 0$.

3. Question

Evaluate each of the following integral:

$$\int_{\pi/6}^{\pi/3} \frac{\sqrt{\tan x}}{\sqrt{\tan x} + \sqrt{\cot x}} dx$$

Answer

Let us assume,

$$I = \int_{\pi/6}^{\pi/3} \frac{\sqrt{\tan x}}{\sqrt{\tan x} + \sqrt{\cot x}} dx \text{ equation 1}$$

By property, we know that $\int_a^b f(x) dx = \int_a^b f(a + b - x) dx$

$$I = \int_{\pi/6}^{\pi/3} \frac{\sqrt{\tan\left(\frac{\pi}{3} + \frac{\pi}{6} - x\right)}}{\sqrt{\tan\left(\frac{\pi}{3} + \frac{\pi}{6} - x\right)} + \sqrt{\cot\left(\frac{\pi}{3} + \frac{\pi}{6} - x\right)}} dx$$

$$I = \int_{\pi/6}^{\pi/3} \frac{\sqrt{\tan(\frac{\pi}{2} - x)}}{\sqrt{\tan(\frac{\pi}{2} - x)} + \sqrt{\cot(\frac{\pi}{2} - x)}} dx$$

Trigonometric property

$$\tan\left(\frac{\pi}{2} - x\right) = \cot x$$

$$\cot\left(\frac{\pi}{2} - x\right) = \tan x$$

$$I = \int_{\pi/6}^{\pi/3} \frac{\sqrt{\cot x}}{\sqrt{\cot x} + \sqrt{\tan x}} dx \dots\dots \text{equation 2}$$

Adding equation 1 and equation 2

$$2I = \int_{\pi/6}^{\pi/3} \frac{\sqrt{\tan x} + \sqrt{\cot x}}{\sqrt{\cot x} + \sqrt{\tan x}} dx$$

$$2I = \int_{\pi/6}^{\pi/3} 1 dx$$

$$2I = [x]_{\pi/6}^{\pi/3}$$

We know $[f(x)]_a^b = f(b) - f(a)$ where b and a are upper and lower limits respectively and $f(x)$ is a function

$$2I = \frac{\pi}{3} - \frac{\pi}{6}$$

$$2I = \frac{\pi}{3} - \frac{\pi}{6}$$

$$2I = \frac{\pi}{6}$$

$$I = \frac{\pi}{12}$$

4. Question

Evaluate each of the following integral:

$$\int_{\pi/6}^{\pi/3} \frac{\sqrt{\sin x}}{\sqrt{\sin x} + \sqrt{\cos x}} dx$$

Answer

Let us assume ,

$$I = \int_{\pi/6}^{\pi/3} \frac{\sqrt{\sin x}}{\sqrt{\sin x} + \sqrt{\cos x}} dx \dots\dots \text{equation 1}$$

By property, we know that $\int_a^b f(x) dx = \int_a^b f(a + b - x) dx$

Thus

$$I = \int_{\pi/6}^{\pi/3} \frac{\sqrt{\sin\left(\frac{\pi}{2} - x\right)}}{\sqrt{\sin\left(\frac{\pi}{2} - x\right)} + \sqrt{\cos\left(\frac{\pi}{2} - x\right)}} dx$$

Trigonometric property

$$\sin\left(\frac{\pi}{2} - x\right) = \cos x$$

$$\cos\left(\frac{\pi}{2} - x\right) = \sin x$$

$$I = \int_{\pi/6}^{\pi/3} \frac{\sqrt{\cos x}}{\sqrt{\cos x} + \sqrt{\sin x}} dx \dots\dots \text{equation 2}$$

Adding equations 1 and equation 2, we get,

$$2I = \int_{\pi/6}^{\pi/3} \frac{\sqrt{\sin x}}{\sqrt{\sin x} + \sqrt{\cos x}} dx + \int_{\pi/6}^{\pi/3} \frac{\sqrt{\cos x}}{\sqrt{\cos x} + \sqrt{\sin x}} dx$$

We know

$$\int_m^n [f(x) + g(x)] dx = \int_m^n f(x) dx + \int_m^n g(x) dx$$

$$2I = \int_{\pi/6}^{\pi/3} \frac{\sqrt{\sin x} + \sqrt{\cos x}}{\sqrt{\sin x} + \sqrt{\cos x}} dx$$

$$2I = \int_{\pi/6}^{\pi/3} 1 dx$$

$$2I = [x]_{\pi/6}^{\pi/3}$$

Since $[f(x)]_a^b = f(b) - f(a)$ where b and a are upper and lower limits respectively

$$2I = \frac{\pi}{3} - \frac{\pi}{6}$$

$$2I = \frac{\pi}{6}$$

$$I = \frac{\pi}{12}$$

5. Question

Evaluate each of the following integral:

$$\int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \frac{\tan^2 x}{1 + e^x} dx$$

Answer

$$\text{Let us assume } I = \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \frac{\tan^2 x}{1 + e^x} dx \dots\dots \text{equation 1}$$

By property, we know that $\int_a^b f(x) dx = \int_a^b f(a + b - x) dx$

Thus

$$I = \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \frac{\tan^2(\frac{\pi}{4} + \frac{-\pi}{4} - x)}{1 + e^{(\frac{\pi}{4} + \frac{-\pi}{4} - x)}} dx$$

$$I = \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \frac{\tan^2(-x)}{1 + e^{-x}} dx$$

$$I = \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \frac{e^x \tan^2(x)}{1 + e^x} dx$$

We know $\tan(-x) = -\tan x$

$$I = \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \frac{e^x \tan^2(x)}{e^{(x)} + 1} dx \dots\dots \text{equation 2}$$

Adding equations 1 and equation 2, we get,

$$2I = \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \frac{\tan^2(x)}{1 + e^{(x)}} dx + \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \frac{e^x \tan^2(x)}{e^{(x)} + 1} dx$$

We know

$$\int_m^n [f(x) + g(x)] dx = \int_m^n f(x) dx + \int_m^n g(x) dx$$

$$2I = \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \frac{\tan^2(x) + e^x \tan^2(x)}{1 + e^{(x)}} dx$$

$$2I = \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \frac{(1 + e^{(x)}) \tan^2(x)}{1 + e^{(x)}} dx$$

$$2I = \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \frac{\tan^2(x)}{1} dx$$

Trigonometric identity $\sec^2 \theta$

$$2I = \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} (\sec^2 x - 1) dx$$

We know $\int \sec^2 \theta d\theta = \tan \theta$

Thus

We know

$$\int_m^n [f(x) + g(x)] dx = \int_m^n f(x) dx + \int_m^n g(x) dx$$

Thus

$$2I = [\tan x - x]_{-\pi/4}^{\pi/4}$$

We know $[f(x)]_a^b = f(b) - f(a)$ b and a being the upper and lower limits respectively.

$$2I = \left[\tan\left(\frac{\pi}{4}\right) - \left(\frac{\pi}{4}\right) \right] - \left[\tan\left(-\frac{\pi}{4}\right) - \left(-\frac{\pi}{4}\right) \right]$$

Since $\tan\frac{\pi}{4} = 1$ and $\tan(-\theta) = -\tan \theta$

Thus

$$2I = \left[1 - \frac{\pi}{4} \right] - \left[-1 + \frac{\pi}{4} \right]$$

$$I = \frac{1}{2} \left[2 - \frac{\pi}{2} \right]$$

6. Question

Evaluate of each of the following integral:

$$\int_{-a}^a \frac{1}{1+a^x} dx$$

Answer

Let us assume $I = \int_{-a}^a \frac{1}{1+a^x} dx$

By integration property, we know,

$$\int_{-m}^m f(x) dx = \int_0^m f(x) dx + \int_0^m f(-x) dx$$

Thus

$$I = \int_0^a \frac{1}{1+a^x} dx + \int_0^a \frac{1}{1+a^{-x}} dx$$

$$I = \int_0^a \frac{1}{1+a^x} dx + \int_0^a \frac{a^x}{1+a^x} dx$$

We know

$$\int_m^n [f(x) + g(x)] dx = \int_m^n f(x) dx + \int_m^n g(x) dx$$

$$I = \int_0^a \frac{1+a^x}{1+a^x} dx$$

$$I = \int_0^a 1 dx$$

$$I = [x]_0^a$$

We know $[f(x)]_a^b = f(b) - f(a)$ and a being the upper and lower limits respectively

$$I = [a - 0]$$

$$I = a.$$

7. Question

Evaluate of each of the following integral:

$$\int_{-\frac{\pi}{3}}^{\frac{\pi}{3}} \frac{1}{1+e^{\tan x}} dx$$

Answer

Let us assume,

$$I = \int_{-\frac{\pi}{3}}^{\frac{\pi}{3}} \frac{1}{1+e^{\tan x}} dx$$

By integration property,

$$\int_{-m}^m f(x)dx = \int_0^m f(x)dx + \int_0^m f(-x)dx$$

Thus

$$I = \int_0^{\frac{\pi}{3}} \frac{1}{1 + e^{\tan x}} dx + \int_0^{\frac{\pi}{3}} \frac{1}{1 + e^{-\tan x}} dx$$

We know

$$\int_m^n [f(x) + g(x)]dx = \int_m^n f(x)dx + \int_m^n g(x)dx$$

$$I = \int_0^{\frac{\pi}{3}} \frac{1 + e^{\tan x}}{1 + e^{\tan x}} dx$$

$$I = \int_0^{\frac{\pi}{3}} 1 dx$$

$$I = [x]_0^{\frac{\pi}{3}}$$

we know since $[f(x)]_a^b = f(b) - f(a)$ b and a being the upper and lower limits

$$I = [\frac{\pi}{3} - 0]$$

$$I = \frac{\pi}{3}$$

8. Question

Evaluate of each of the following integral:

$$\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{\cos^2 x}{1 + e^x} dx.$$

Answer

$$\text{Let us assume } I = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{\cos^2 x}{1 + e^x} dx \dots\dots \text{equation 1}$$

By property, we know that $\int_a^b f(x)dx = \int_a^b f(a + b - x)dx$

Thus

$$I = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{\cos^2(\frac{\pi}{4} + \frac{-\pi}{4} - x)}{1 + e^{(\frac{\pi}{4} + \frac{-\pi}{4} - x)}} dx$$

$$I = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{\cos^2(-x)}{1 + e^{-x}} dx$$

$$I = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{\cos^2(-x)}{1 + \frac{1}{e^x}} dx$$

$$I = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{e^x \cos^2(x)}{e^{(x)} + 1} dx \dots\dots \text{equation 2}$$

Adding the equations 1 and 2, we get,

$$2I = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{\cos^2(x)}{1+e^{(x)}} dx + \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{e^x \cos^2(x)}{e^{(x)}+1} dx$$

We know

$$\int_m^n [f(x) + g(x)] dx = \int_m^n f(x) dx + \int_m^n g(x) dx$$

$$2I = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{\cos^2(x) + e^x \cos^2(x)}{1+e^{(x)}} dx$$

$$2I = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{(1+e^{(x)})\cos^2(x)}{1+e^{(x)}} dx$$

$$2I = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{\cos^2(x)}{1} dx$$

Trigonometric formula

$$2\cos^2\theta - 1 = \cos 2\theta$$

$$2I = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{\cos 2x + 1}{2} dx$$

We know $\int \cos \theta d\theta = \sin \theta$

Thus

We know

$$\int_m^n [f(x) + g(x)] dx = \int_m^n f(x) dx + \int_m^n g(x) dx$$

$$2I = \frac{1}{2} \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \cos 2x dx + 1/2 \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} dx$$

Thus

$$2I = \frac{\left[\frac{\sin 2x}{2} \right]_{-\frac{\pi}{2}}^{\frac{\pi}{2}} + [x]_{-\frac{\pi}{2}}^{\frac{\pi}{2}}}{2}$$

we know $[f(x)]_a^b = f(b) - f(a)$ b and a being the upper and lower limit

$$4I = \left[\frac{\sin\left(\frac{2\pi}{2}\right)}{2} - \frac{\sin\left(\frac{-2\pi}{2}\right)}{2} \right] + \left[\left(\frac{\pi}{2}\right) - \left(-\frac{\pi}{2}\right) \right]$$

Since $\sin \pi = 0$ and $\sin(-\theta) = -\sin \theta$

Thus

$$4I = [0 - 0] + \left[\frac{\pi}{1} \right]$$

$$I = \left[\frac{\pi}{4} \right]$$

9. Question

Evaluate of each of the following integral:

$$\int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \frac{x^{11} - 3x^9 + 5x^7 - x^5 + 1}{\cos^2 x} dx$$

Answer

Let us assume $I = \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \frac{x^{11} - 3x^9 + 5x^7 - x^5 + 1}{\cos^2 x} dx$

By property, we know,

$$\int_{-m}^m f(x) dx = \int_0^m f(x) dx + \int_0^m f(-x) dx$$

$$I = \int_0^{\frac{\pi}{4}} \frac{x^{11} - 3x^9 + 5x^7 - x^5 + 1}{\cos^2 x} dx + \int_0^{\frac{\pi}{4}} \frac{-x^{11} + 3x^9 - 5x^7 + x^5 + 1}{\cos^2 x} dx$$

We know

$$\int_m^n [f(x) + g(x)] dx = \int_m^n f(x) dx + \int_m^n g(x) dx$$

$$I = \int_0^{\frac{\pi}{4}} \frac{x^{11} - 3x^9 + 5x^7 - x^5 + 1 - x^{11} + 3x^9 - 5x^7 + x^5 + 1}{\cos^2 x} dx$$

$$I = \int_0^{\frac{\pi}{4}} \frac{2}{\cos^2 x} dx$$

$$I = \int_0^{\frac{\pi}{4}} 2 \sec^2 x dx$$

$$I = [2 \tan x]_0^{\frac{\pi}{4}}$$

We know $[f(x)]_a^b = f(b) - f(a)$ b and a being the upper and lower limits

$$I = [2(1 - 0)]$$

Since $\tan \frac{\pi}{4} = 1$

$I = 2.$

10. Question

Evaluate of each of the following integral:

$$\int_a^b \frac{x^{1/n}}{x^{1/n} + (a + b - x)^{1/n}} dx, n \in N, n \geq 2$$

Answer

Let us assume $I = \int_a^b \frac{x^{1/n}}{x^{1/n} + (a + b - x)^{1/n}} dx \dots \text{equation 1}$

By property, we know that,

$$\int_a^b f(x)dx = \int_a^b f(a+b-x)dx$$

$$I = \int_a^b \frac{(a+b-x)^{1/n}}{(a+b-x)^{1/n} + (x)^{1/n}} dx \dots\dots \text{equation 2}$$

Adding equation 1 and equation 2

$$2I = \int_a^b \frac{x^{1/n}}{x^{1/n} + (a+b-x)^{1/n}} dx + \int_a^b \frac{(a+b-x)^{1/n}}{(a+b-x)^{1/n} + (x)^{1/n}} dx$$

We know

$$\int_m^n [f(x) + g(x)]dx = \int_m^n f(x)dx + \int_m^n g(x)dx$$

$$2I = \int_a^b \frac{x^{1/n} + (a+b-x)^{1/n}}{x^{1/n} + (a+b-x)^{1/n}} dx$$

$$I = \int_a^b 1 dx$$

$$2I = [x]_a^b$$

We know $[f(x)]_a^b = f(b) - f(a)$ b and a being the upper and lower limit

$$I = \frac{b-a}{2}$$

11. Question

Evaluate of each of the following integral:

$$\int_0^{\frac{\pi}{2}} 2\log\cos x - \log\sin 2x dx$$

Answer

$$\text{Let us assume } I = \int_0^{\frac{\pi}{2}} 2\log\cos x - \log\sin 2x dx$$

We know $n\log m = \log m^n$ and $\sin 2\theta = 2\sin\theta\cos\theta$

Thus

$$I = \int_0^{\frac{\pi}{2}} \log\cos^2 x - \log 2\sin x \cos x dx$$

We know

$$\log m - \log n = \log\left(\frac{m}{n}\right)$$

Thus

$$I = \int_0^{\frac{\pi}{2}} \log\left(\frac{\cos^2 x}{2\sin x \cos x}\right) dx$$

$$\text{Since } \tan x = \frac{\sin x}{\cos x}, \cot x = \frac{1}{\tan x} = \frac{\cos x}{\sin x}$$

$$I = \int_0^{\frac{\pi}{2}} \log\left(\frac{\cos x}{2\sin x}\right) dx$$

$$I = \int_0^{\frac{\pi}{2}} \log\left(\frac{\cot x}{2}\right) dx \dots\dots \text{equation 1}$$

By property, we know that,

$$\int_a^b f(x)dx = \int_a^b f(a+b-x)dx$$

Thus

$$I = \int_0^{\frac{\pi}{2}} \log\left(\frac{\cot\left(\frac{\pi}{2} + 0 - x\right)}{2}\right) dx$$

$$\text{Since } \cot\left(\frac{\pi}{2} - \theta\right) = \tan\theta$$

$$I = \int_0^{\frac{\pi}{2}} \log\left(\frac{\tan x}{2}\right) dx \dots\dots \text{equation 2}$$

Adding equations 1 and 2

$$2I = \int_0^{\frac{\pi}{2}} \log\left(\frac{\tan x}{2}\right) dx + \int_0^{\frac{\pi}{2}} \log\left(\frac{\cot x}{2}\right) dx$$

We know

$$\int_m^n [f(x) + g(x)]dx = \int_m^n f(x)dx + \int_m^n g(x)dx$$

$$2I = \int_0^{\frac{\pi}{2}} \log\left(\frac{\tan x}{2}\right) + \log\left(\frac{\cot x}{2}\right) dx$$

Since we know that $\log m + \log n = \log mn$

$$2I = \int_0^{\frac{\pi}{2}} \log \frac{\tan x}{2} * \frac{\cot x}{2} dx$$

Since $\tan x = 1/\cot x$

$$2I = \int_0^{\frac{\pi}{2}} \log \frac{1}{2} * \frac{1}{2} dx$$

$$2I = \left[\log\left(\frac{1}{4}\right)x \right]_0^{\frac{\pi}{2}}$$

We know $[f(x)]_a^b = f(b) - f(a)$

$$2I = \log\left(\frac{1}{4}\right)\left[\frac{\pi}{2} - 0\right]$$

$$I = \frac{\pi}{4} \log\left(\frac{1}{4}\right)$$

12. Question

Evaluate of each of the following integral:

$$\int_0^a \frac{\sqrt{x}}{\sqrt{x} + \sqrt{a-x}} dx$$

Answer

Let us assume $I = \int_0^a \frac{\sqrt{x}}{\sqrt{x} + \sqrt{a-x}} dx$equation 1

By property, we know that,

$$\int_a^b f(x)dx = \int_a^b f(a+b-x)dx$$

$$I = \int_0^a \frac{\sqrt{a-x}}{\sqrt{a-x} + \sqrt{x}} dx \text{equation 2}$$

$$\text{Adding equation 1 and 2 } 2I = \int_0^a \frac{\sqrt{x}}{\sqrt{x} + \sqrt{a-x}} dx + \int_0^a \frac{\sqrt{a-x}}{\sqrt{a-x} + \sqrt{x}} dx$$

We know

$$\int_m^n [f(x) + g(x)]dx = \int_m^n f(x)dx + \int_m^n g(x)dx$$

$$2I = \int_0^a \frac{\sqrt{x} + \sqrt{a-x}}{\sqrt{x} + \sqrt{a-x}} dx$$

$$2I = \int_0^a 1dx$$

We know $[f(x)]_a^b = f(b) - f(a)$

$$2I = [x]_0^a$$

$$2I = [a - 0]$$

$$2I = a$$

$$I = \frac{a}{2}$$

13. Question

Evaluate of each of the following integral:

$$\int_0^5 \frac{\sqrt[4]{x+4}}{\sqrt[4]{x+4} + \sqrt[4]{9-x}} dx$$

Answer

Let us assume $I = \int_0^5 \frac{\sqrt[4]{x+4}}{\sqrt[4]{x+4} + \sqrt[4]{9-x}} dx$ equation 1

By property we know that $\int_a^b f(x)dx = \int_a^b f(a+b-x)dx$

$$I = \int_0^5 \frac{\sqrt[4]{9-x}}{\sqrt[4]{9-x} + \sqrt[4]{x+4}} dx \text{equation 2}$$

Adding equation 1 and 2

$$2I = \int_0^5 \frac{\sqrt[4]{x+4}}{\sqrt[4]{x+4} + \sqrt[4]{9-x}} dx + \int_0^5 \frac{\sqrt[4]{9-x}}{\sqrt[4]{9-x} + \sqrt[4]{x+4}} dx$$

We know

$$\int_m^n [f(x) + g(x)]dx = \int_m^n f(x)dx + \int_m^n g(x)dx$$

$$2I = \int_0^5 \frac{\sqrt[4]{9-x} + \sqrt[4]{x+4}}{\sqrt[4]{9-x} + \sqrt[4]{x+4}} dx$$

$$2I = \int_0^5 1 dx$$

$$2I = [x]_0^5$$

We know $[f(x)]_a^b = f(b) - f(a)$

$$2I = [5 - 0]$$

$$2I = 5$$

$$I = \frac{5}{2}$$

14. Question

Evaluate each of the following integral:

$$\int_0^7 \frac{\sqrt[3]{x}}{\sqrt[3]{x} + \sqrt[3]{7-x}} dx$$

Answer

Let us assume $I = \int_0^7 \frac{\sqrt[3]{x}}{\sqrt[3]{x} + \sqrt[3]{7-x}} dx \dots\dots \text{equation 1}$

By property, we know that $\int_a^b f(x) dx = \int_a^b f(a+b-x) dx$

$$I = \int_0^7 \frac{\sqrt[3]{7-x}}{\sqrt[3]{7-x} + \sqrt[3]{x}} dx \dots\dots \text{equation 2}$$

Adding equation 1 and 2

$$2I = \int_0^7 \frac{\sqrt[3]{x}}{\sqrt[3]{x} + \sqrt[3]{7-x}} dx + \int_0^7 \frac{\sqrt[3]{7-x}}{\sqrt[3]{7-x} + \sqrt[3]{x}} dx$$

We know

$$\int_m^n [f(x) + g(x)] dx = \int_m^n f(x) dx + \int_m^n g(x) dx$$

$$2I = \int_0^7 \frac{\sqrt[3]{7-x} + \sqrt[3]{x}}{\sqrt[3]{7-x} + \sqrt[3]{x}} dx$$

$$2I = \int_0^7 1 dx$$

$$2I = [x]_0^7$$

We know $[f(x)]_a^b = f(b) - f(a)$

$$2I = [7 - 0]$$

$$2I = 7$$

$$I = \frac{7}{2}$$

15. Question

Evaluate each of the following integral:

$$\int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{1}{1 + \sqrt{\tan x}} dx$$

Answer

Let us assume $I = \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{1}{1 + \sqrt{\tan x}} dx$

We know

$$\tan x = \frac{\sin x}{\cos x}$$

$$I = \int_{\pi/6}^{\pi/3} \frac{\sqrt{\cos x}}{\sqrt{\sin x} + \sqrt{\cos x}} dx \dots\dots\dots \text{equation 1}$$

By property, we know that $\int_a^b f(x) dx = \int_a^b f(a + b - x) dx$

Thus

$$I = \int_{\pi/6}^{\pi/3} \frac{\sqrt{\cos(\frac{\pi}{2} - x)}}{\sqrt{\sin(\frac{\pi}{2} - x)} + \sqrt{\cos(\frac{\pi}{2} - x)}} dx$$

Trigonometric property

$$\sin\left(\frac{\pi}{2} - x\right) = \cos x$$

$$\cos\left(\frac{\pi}{2} - x\right) = \sin x$$

$$I = \int_{\pi/6}^{\pi/3} \frac{\sqrt{\sin x}}{\sqrt{\cos x} + \sqrt{\sin x}} dx \dots\dots\dots \text{equation 2}$$

Adding equations 1 and 2, we get,

$$2I = \int_{\pi/6}^{\pi/3} \frac{\sqrt{\cos x}}{\sqrt{\sin x} + \sqrt{\cos x}} dx + \int_{\pi/6}^{\pi/3} \frac{\sqrt{\sin x}}{\sqrt{\cos x} + \sqrt{\sin x}} dx$$

We know

$$\int_m^n [f(x) + g(x)] dx = \int_m^n f(x) dx + \int_m^n g(x) dx$$

$$2I = \int_{\pi/6}^{\pi/3} \frac{\sqrt{\sin x} + \sqrt{\cos x}}{\sqrt{\sin x} + \sqrt{\cos x}} dx$$

$$2I = \int_{\pi/6}^{\pi/3} 1 dx$$

$$2I = [x]_{\pi/6}^{\pi/3}$$

We know $[f(x)]_a^b = f(b) - f(a)$

$$2I = \left[\frac{\pi}{3} - \frac{\pi}{6} \right]$$

$$2I = \frac{\pi}{6}$$

$$I = \frac{\pi}{12}$$

16. Question

If $f(a + b - x) = f(x)$, then prove that $\int_a^b xf(x) dx = \frac{a+b}{2} \int_a^b f(x) dx$.

Answer

LHS

By property, we know that $\int_a^b f(x) dx = \int_a^b f(a + b - x) dx$

thus

$$\int_a^b xf(x) dx = \int_a^b (a + b - x)f(a + b - x) dx$$

Given $f(a + b - x) = f(x)$

$$\int_a^b xf(x) dx = \int_a^b (a + b - x)f(x) dx$$

We know

$$\int_m^n [f(x) + g(x)] dx = \int_m^n f(x) dx + \int_m^n g(x) dx$$

Thus

$$\int_a^b xf(x) dx = \int_a^b (a + b)f(x) dx - \int_a^b xf(x) dx$$

$$2 \int_a^b xf(x) dx = \int_a^b (a + b)f(x) dx$$

$$\int_a^b xf(x) dx = (a + b)/2 \int_a^b f(x) dx$$

Hence proved

Exercise 20.5

1. Question

Evaluate the following integral:

$$\int_0^{\frac{\pi}{2}} \frac{1}{1 + \tan x} dx$$

Answer

Let us assume $I = \int_0^{\frac{\pi}{2}} \frac{1}{1 + \tan x} dx$equation 1

We know that $\tan x = \frac{\sin x}{\cos x}$

Substituting the value in equation 1 we have,

$$I = \int_0^{\frac{\pi}{2}} \frac{1}{1 + \frac{\sin x}{\cos x}} dx$$

$$I = \int_0^{\frac{\pi}{2}} \frac{\cos x}{\cos x + \sin x} dx \dots \text{equation 2}$$

By property, we know that $\int_a^b f(x) dx = \int_a^b f(a + b - x) dx$

Thus in equation 2

$$I = \int_0^{\frac{\pi}{2}} \frac{\cos(\frac{\pi}{2} - x)}{\cos(\frac{\pi}{2} - x) + \sin(\frac{\pi}{2} - x)} dx$$

$$I = \int_0^{\frac{\pi}{2}} \frac{\sin x}{\sin x + \cos x} dx \dots \text{equation 3}$$

Adding equation 2 and 3

$$2I = \int_0^{\frac{\pi}{2}} \frac{\cos x}{\cos x + \sin x} dx + \int_0^{\frac{\pi}{2}} \frac{\sin x}{\cos x + \sin x} dx$$

We know

$$\int_m^n [f(x) + g(x)] dx = \int_m^n f(x) dx + \int_m^n g(x) dx$$

Thus

$$2I = \int_0^{\frac{\pi}{2}} \frac{\cos x + \sin x}{\cos x + \sin x} dx$$

$$2I = \int_0^{\frac{\pi}{2}} 1 dx$$

$$2I = [x]_0^{\frac{\pi}{2}}$$

$$2I = \left[\frac{\pi}{2} - 0 \right]$$

$$I = \frac{\pi}{4}$$

2. Question

Evaluate the following integral:

$$\int_0^{\frac{\pi}{2}} \frac{1}{1 + \cot x} dx$$

Answer

$$\text{Let us assume } I = \int_0^{\frac{\pi}{2}} \frac{1}{1 + \cot x} dx \dots \text{equation 1}$$

$$\text{We know that } \tan x = \frac{\cos x}{\sin x} \quad \cot x = \frac{1}{\tan x}$$

Substituting the value in equation 1 we have,

$$I = \int_0^{\frac{\pi}{2}} \frac{1}{1 + \frac{\cos x}{\sin x}} dx$$

$$I = \int_0^{\frac{\pi}{2}} \frac{\sin x}{\cos x + \sin x} dx \dots \text{equation 2}$$

By property, we know that $\int_a^b f(x) dx = \int_a^b f(a + b - x) dx$

Thus in equation 2

$$I = \int_0^{\frac{\pi}{2}} \frac{\sin(\frac{\pi}{2} - x)}{\sin(\frac{\pi}{2} - x) + \cos(\frac{\pi}{2} - x)} dx$$

$$I = \int_0^{\frac{\pi}{2}} \frac{\cos x}{\sin x + \cos x} dx \dots \text{equation 3}$$

Adding equation 2 and 3

$$2I = \int_0^{\frac{\pi}{2}} \frac{\sin x}{\cos x + \sin x} dx + \int_0^{\frac{\pi}{2}} \frac{\cos x}{\cos x + \sin x} dx$$

We know

$$\int_m^n [f(x) + g(x)] dx = \int_m^n f(x) dx + \int_m^n g(x) dx$$

Thus

$$2I = \int_0^{\frac{\pi}{2}} \frac{\cos x + \sin x}{\cos x + \sin x} dx$$

$$2I = \int_0^{\frac{\pi}{2}} 1 dx$$

$$2I = [x]_0^{\frac{\pi}{2}}$$

$$2I = \left[\frac{\pi}{2} - 0 \right]$$

$$I = \frac{\pi}{4}$$

3. Question

Evaluate the following integral:

$$\int_0^{\frac{\pi}{2}} \frac{\sqrt{\cot x}}{\sqrt{\tan x} + \sqrt{\cot x}} dx$$

Answer

$$\text{Let us assume } I = \int_0^{\frac{\pi}{2}} \frac{\sqrt{\cot x}}{\sqrt{\tan x} + \sqrt{\cot x}} dx \dots \text{equation 1}$$

By property, we know that $\int_a^b f(x) dx = \int_a^b f(a + b - x) dx$

Thus in equation 2

$$I = \int_0^{\frac{\pi}{2}} \frac{\sqrt{\cot\left(\frac{\pi}{2} - x\right)}}{\sqrt{\tan\left(\frac{\pi}{2} - x\right)} + \sqrt{\cot\left(\frac{\pi}{2} - x\right)}} dx$$

$$I = \int_0^{\frac{\pi}{2}} \frac{\sqrt{\tan x}}{\sqrt{\tan x} + \sqrt{\cot x}} dx \dots \text{equation 2}$$

Adding equation 1 and 2

$$2I = \int_0^{\frac{\pi}{2}} \frac{\sqrt{\cot x}}{\sqrt{\tan x} + \sqrt{\cot x}} dx + \int_0^{\frac{\pi}{2}} \frac{\sqrt{\tan x}}{\sqrt{\tan x} + \sqrt{\cot x}} dx$$

We know

$$\int_m^n [f(x) + g(x)] dx = \int_m^n f(x) dx + \int_m^n g(x) dx$$

Thus

$$2I = \int_0^{\frac{\pi}{2}} \frac{\sqrt{\cot x} + \sqrt{\tan x}}{\sqrt{\tan x} + \sqrt{\cot x}} dx$$

$$2I = \int_0^{\frac{\pi}{2}} 1 dx$$

$$2I = [x]_0^{\frac{\pi}{2}}$$

$$2I = \left[\frac{\pi}{2} - 0 \right]$$

$$I = \frac{\pi}{4}$$

4. Question

Evaluate the following integral:

$$\int_0^{\frac{\pi}{2}} \frac{\sin^{\frac{3}{2}} x}{\sin^{\frac{3}{2}} x + \cos^{\frac{3}{2}} x} dx$$

Answer

$$\text{Let us assume } I = \int_0^{\frac{\pi}{2}} \frac{\sin^{\frac{3}{2}} x}{\sin^{\frac{3}{2}} x + \cos^{\frac{3}{2}} x} \dots \text{equation 1}$$

$$\text{By property, we know that } \int_a^b f(x) dx = \int_a^b f(a + b - x) dx$$

Thus in equation 2

$$I = \int_0^{\frac{\pi}{2}} \frac{\sin^{\frac{3}{2}}(\frac{\pi}{2} - x)}{\sin^{\frac{3}{2}}(\frac{\pi}{2} - x) + \cos^{\frac{3}{2}}(\frac{\pi}{2} - x)} dx$$

$$I = \int_0^{\frac{\pi}{2}} \frac{\cos^{\frac{3}{2}} x}{\sin^{\frac{3}{2}} x + \cos^{\frac{3}{2}} x} \dots \text{equation 2}$$

Adding equation 1 and 2

$$2I = \int_0^{\frac{\pi}{2}} \frac{\sin^3 x}{\sin^3 x + \cos^3 x} dx + \int_0^{\frac{\pi}{2}} \frac{\cos^3 x}{\sin^3 x + \cos^3 x} dx$$

We know

$$\int_m^n [f(x) + g(x)] dx = \int_m^n f(x) dx + \int_m^n g(x) dx$$

Thus

$$2I = \int_0^{\frac{\pi}{2}} \frac{\cos^3 x + \sin^3 x}{\sin^3 x + \cos^3 x} dx$$

$$2I = \int_0^{\frac{\pi}{2}} 1 dx$$

$$2I = [x]_0^{\frac{\pi}{2}}$$

$$2I = \left[\frac{\pi}{2} - 0 \right]$$

$$I = \frac{\pi}{4}$$

5. Question

Evaluate the following integral:

$$\int_0^{\frac{\pi}{2}} \frac{\sin^n x}{\sin^n x + \cos^n x} dx$$

Answer

$$\text{Let us assume } I = \int_0^{\frac{\pi}{2}} \frac{\sin^n x}{\sin^n x + \cos^n x} \dots \text{equation 1}$$

$$\text{By property, we know that } \int_a^b f(x) dx = \int_a^b f(a + b - x) dx$$

Thus in equation 2

$$I = \int_0^{\frac{\pi}{2}} \frac{\sin^n(\frac{\pi}{2}-x)}{\sin^n(\frac{\pi}{2}-x) + \cos^n(\frac{\pi}{2}-x)} dx$$

$$I = \int_0^{\frac{\pi}{2}} \frac{\cos^n x}{\sin^n x + \cos^n x} \dots \text{equation 2}$$

Adding equation 1 and 2

$$2I = \int_0^{\frac{\pi}{2}} \frac{\sin^n x}{\sin^n x + \cos^n x} dx + \int_0^{\frac{\pi}{2}} \frac{\cos^n x}{\sin^n x + \cos^n x} dx$$

We know

$$\int_m^n [f(x) + g(x)] dx = \int_m^n f(x) dx + \int_m^n g(x) dx$$

Thus

$$2I = \int_0^{\frac{\pi}{2}} \frac{\cos^n x + \sin^n x}{\sin^n x + \cos^n x} dx$$

$$2I = \int_0^{\frac{\pi}{2}} 1 \, dx$$

$$2I = [x]_0^{\frac{\pi}{2}}$$

$$2I = \left[\frac{\pi}{2} - 0 \right]$$

$$I = \frac{\pi}{4}$$

6. Question

Evaluate the following integral:

$$\int_0^{\frac{\pi}{2}} \frac{1}{1 + \sqrt{\tan x}} \, dx$$

Answer

$$\text{Let us assume } I = \int_0^{\frac{\pi}{2}} \frac{1}{1 + \sqrt{\tan x}} \, dx \dots \text{equation 1}$$

$$\text{We know that } \tan x = \frac{\sin x}{\cos x}$$

Substituting the value in equation 1 we have,

$$I = \int_0^{\frac{\pi}{2}} \frac{1}{1 + \frac{\sqrt{\sin x}}{\sqrt{\cos x}}} \, dx$$

$$I = \int_0^{\frac{\pi}{2}} \frac{\sqrt{\cos x}}{\sqrt{\cos x} + \sqrt{\sin x}} \, dx \dots \text{equation 2}$$

$$\text{By property, we know that } \int_a^b f(x) \, dx = \int_a^b f(a + b - x) \, dx$$

Thus in equation 2

$$I = \int_0^{\frac{\pi}{2}} \frac{\sqrt{\cos(\frac{\pi}{2} - x)}}{\sqrt{\cos(\frac{\pi}{2} - x)} + \sqrt{\sin(\frac{\pi}{2} - x)}} \, dx$$

$$I = \int_0^{\frac{\pi}{2}} \frac{\sqrt{\sin x}}{\sqrt{\sin x} + \sqrt{\cos x}} \, dx \dots \text{equation 3}$$

Adding equation 2 and 3

$$2I = \int_0^{\frac{\pi}{2}} \frac{\sqrt{\cos x}}{\sqrt{\cos x} + \sqrt{\sin x}} \, dx + \int_0^{\frac{\pi}{2}} \frac{\sqrt{\sin x}}{\sqrt{\cos x} + \sqrt{\sin x}} \, dx$$

We know

$$\int_m^n [f(x) + g(x)] \, dx = \int_m^n f(x) \, dx + \int_m^n g(x) \, dx$$

Thus

$$2I = \int_0^{\frac{\pi}{2}} \frac{\sqrt{\cos x} + \sqrt{\sin x}}{\sqrt{\cos x} + \sqrt{\sin x}} \, dx$$

$$2I = \int_0^{\frac{\pi}{2}} 1 dx$$

$$2I = [x]_0^{\frac{\pi}{2}}$$

$$2I = \left[\frac{\pi}{2} - 0 \right]$$

$$I = \frac{\pi}{4}$$

7. Question

Evaluate the following integral:

$$\int_0^a \frac{1}{x + \sqrt{a^2 - x^2}} dx$$

Answer

$$\text{Let us assume } I = \int_0^a \frac{1}{x + \sqrt{a^2 - x^2}} dx \dots\dots\dots \text{equation 1}$$

$$\text{Let } x = a \cos \theta$$

thus

$$x = a \cos \theta$$

Differentiating both sides, we get,

$$dx = -a \sin \theta d\theta$$

Thus substituting old limits, we get a new upper limit and lower limit

$$\text{For } a = a \cos \theta$$

$$0 = \theta$$

$$\text{For } 0 = a \cos \theta$$

$$\frac{\pi}{2} = \theta$$

$$\text{We know that } \int_a^b -f(x) = \int_b^a f(x)$$

thus

Substituting the values in equation 1

$$I = \int_{\frac{\pi}{2}}^0 \frac{1}{a \cos \theta + \sqrt{a^2 - a^2 \cos^2 \theta}} (-a \sin \theta) d\theta$$

$$\text{We know that } \int_a^b -f(x) = \int_b^a f(x)$$

Trigonometric identity $1 - \cos^2 \theta = \sin^2 \theta$

$$I = \int_0^{\frac{\pi}{2}} \frac{1}{a \cos \theta + \sqrt{a^2 (1 - \cos^2 \theta)}} (a \sin \theta) d\theta$$

$$I = \int_0^{\frac{\pi}{2}} \frac{1}{a \cos \theta + \sqrt{a^2 \sin^2 \theta}} (a \sin \theta) d\theta$$

$$I = \int_0^{\frac{\pi}{2}} \frac{1}{\cos \theta + \sin \theta} (\sin \theta) d\theta$$

$$I = \int_0^{\frac{\pi}{2}} \frac{1}{\cos \theta + \sin \theta} (\cos \theta) d\theta \dots\dots \text{equation 2}$$

By property, we know that $\int_a^b f(x) dx = \int_a^b f(a + b - x) dx$

thus

$$I = \int_0^{\frac{\pi}{2}} \frac{1}{\cos(\frac{\pi}{2} - \theta) + \sin(\frac{\pi}{2} - \theta)} (\sin(\frac{\pi}{2} - \theta)) d\theta$$

$$I = \int_0^{\frac{\pi}{2}} \frac{1}{\sin \theta + \cos \theta} (\cos \theta) d\theta \dots\dots \text{equation 3}$$

Adding equation 3 and equation 2

Thus

$$2I = \int_0^{\frac{\pi}{2}} \frac{1}{\cos \theta + \sin \theta} (\sin \theta) d\theta + \int_0^{\frac{\pi}{2}} \frac{1}{\cos \theta + \sin \theta} (\cos \theta) d\theta$$

We know

$$\int_m^n [f(x) + g(x)] dx = \int_m^n f(x) dx + \int_m^n g(x) dx$$

$$2I = \int_0^{\frac{\pi}{2}} \frac{1}{\cos \theta + \sin \theta} (\sin \theta + \cos \theta) d\theta$$

$$2I = \int_0^{\frac{\pi}{2}} 1 d\theta$$

$$2I = [\theta]_0^{\frac{\pi}{2}}$$

We know $[f(x)]_a^b = f(b) - f(a)$ b and a being the upper and lower limits respectively.

$$2I = [\frac{\pi}{2} - 0]$$

$$I = \frac{\pi}{4}$$

8. Question

Evaluate the following integral:

$$\int_0^{\infty} \frac{\log x}{1+x^2} dx$$

Answer

$$\text{let us assume } I = \int_0^{\infty} \frac{\log x}{1+x^2} dx$$

$$\text{let } x = t \tan y$$

differentiating both sides

$$dx = \sec^2 y dy$$

for $x = \infty$

$$\frac{\pi}{2} = y$$

For $x = 0$

$$0 = y$$

thus

$$I = \int_0^{\frac{\pi}{2}} \frac{\log(\tan y)}{1 + \tan^2 y} \sec^2 y dy$$

$$I = \int_0^{\frac{\pi}{2}} \frac{\log(\tan y)}{\sec^2 y} \sec^2 y dy$$

(since $\sec^2 y - \tan^2 y = 1$)

$$I = \int_0^{\frac{\pi}{2}} \log(\tan y) dy \dots\dots \text{equation 1}$$

By property, we know that $\int_a^b f(x) dx = \int_a^b f(a + b - x) dx$

$$I = \int_0^{\frac{\pi}{2}} \log\left(\tan\left(\frac{\pi}{2} - y\right)\right) dy$$

$$I = \int_0^{\frac{\pi}{2}} \log(\cot y) dy \dots\dots \text{equation 2}$$

Adding equations 1 and 2, we get,

$$2I = \int_0^{\frac{\pi}{2}} \log(\tan y) dy + \int_0^{\frac{\pi}{2}} \log(\cot y) dy$$

We know

$$\int_m^n [f(x) + g(x)] dx = \int_m^n f(x) dx + \int_m^n g(x) dx$$

$$2I = \int_0^{\frac{\pi}{2}} [\log(\tan y) + \log(\cot y)] dy$$

$$2I = \int_0^{\frac{\pi}{2}} [\log(\tan y \times \cot y)] dy \text{ since } \log m + \log n = \log mn$$

$$2I = \int_0^{\frac{\pi}{2}} [\log 1] dy \text{ since } \tan y \times \cot y = 1$$

$$2I = \int_0^{\frac{\pi}{2}} 0 dy \text{ since } \log 1 = 0$$

Thus

$$2I = 0$$

$$I = 0$$

9. Question

Evaluate the following integral:

$$\int_0^1 \frac{\log(1+x)}{1+x^2} dx$$

Answer

Let us assume $I = \int_0^1 \frac{\log(1+x)}{1+x^2} dx$ equation 1

Let $x = \tan \theta$ thus

Differentiating both sides, we get,

$$dx = \sec^2 \theta d\theta$$

Thus substituting old limits, we get a new upper limit and lower limit

For $1 = \tan \theta$

$$\frac{\pi}{4} = \theta$$

For $0 = \tan \theta$

$$0 = \theta$$

substitute the values in equation 1

$$\text{we get } I = \int_0^{\frac{\pi}{4}} \frac{\log(1 + \tan \theta)}{1 + \tan^2 \theta} \sec^2 \theta \, d\theta \dots \text{equation 2}$$

trigonometric identity we know

$$1 + \tan^2 \theta = \sec^2 \theta$$

Thus substituting in equation 2 we have

$$I = \int_0^{\frac{\pi}{4}} \frac{\log(1 + \tan \theta)}{\sec^2 \theta} \sec^2 \theta d\theta$$

$$I = \int_{0}^{\pi} \log(1 + \tan \theta) d\theta \quad \dots \text{equation 3}$$

By property, we know that $\int_a^b f(x)dx = \int_a^b f(a + b - x)dx$

Thus

$$I = \int_0^{\frac{\pi}{4}} \log\left(1 + \tan\left(\frac{\pi}{4} - \theta\right)\right) d\theta \dots \text{equation 4}$$

Trigonometric formula:

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

$$\text{Thus } \tan\left(\frac{\pi}{4} - \theta\right) = \frac{\tan\frac{\pi}{4} - \tan\theta}{1 + \tan\frac{\pi}{4}\tan\theta}$$

We know by trigonometric property:

$$\tan \frac{\pi}{4} = 1$$

thus

$$\tan\left(\frac{\pi}{4} - \theta\right) = \frac{1 - \tan\theta}{1 + \tan\theta}$$

Substituting in equation 4

$$I = \int_0^{\frac{\pi}{4}} \log\left(1 + \frac{1 - \tan\theta}{1 + \tan\theta}\right) d\theta$$

$$I = \int_0^{\frac{\pi}{4}} \log\left(\frac{1 + \tan\theta + 1 - \tan\theta}{1 + \tan\theta}\right) d\theta$$

$$I = \int_0^{\frac{\pi}{4}} \log\left(\frac{2}{1 + \tan\theta}\right) d\theta$$

We know $\log\left(\frac{m}{n}\right) = \log m - \log n$

Thus

$$I = \int_0^{\frac{\pi}{4}} \log(2) - \log(1 + \tan\theta) d\theta \dots\dots \text{equation 6}$$

We know

$$\int_m^n [f(x) + g(x)] dx = \int_m^n f(x) dx + \int_m^n g(x) dx$$

Adding equation 3 and equation 6

$$2I = \int_0^{\frac{\pi}{4}} \log(1 + \tan\theta) d\theta + \int_0^{\frac{\pi}{4}} \log(2) - \log(1 + \tan\theta) d\theta$$

Thus

$$2I = \int_0^{\frac{\pi}{4}} \log(1 + \tan\theta) + \log(2) - \log(1 + \tan\theta) d\theta$$

$$2I = \int_0^{\frac{\pi}{4}} \log(2) d\theta$$

$$2I = \log(2) \int_0^{\frac{\pi}{4}} 1 d\theta$$

$$2I = \log 2 [\theta]_0^{\frac{\pi}{4}}$$

We know $[f(x)]_a^b = f(b) - f(a)$ b and a being the upper and lower limits respectively.

$$2I = \log 2 \left[\frac{\pi}{4} - 0\right]$$

$$I = \frac{\pi}{8} \log 2$$

10. Question

Evaluate the following integral:

$$\int_0^{\infty} \frac{x}{(1+x)(1+x^2)} dx$$

Answer

$$\text{Let us assume } I = \int_0^{\infty} \frac{x}{(1+x)(1+x^2)} dx$$

Adding -1 and +1

$$I = \int_0^{\infty} \frac{x+1-1}{(1+x)(1+x^2)} dx$$

$$I = \int_0^{\infty} \frac{x+1}{(1+x)(1+x^2)} - \frac{1}{(1+x)(1+x^2)} dx$$

$$I = \int_0^{\infty} \frac{1}{(1+x^2)} dx - \int_0^{\infty} \frac{1}{(1+x)(1+x^2)} dx$$

$$\text{Let } I_1 = \int_0^{\infty} \frac{1}{(1+x^2)} dx$$

$$I_2 = \int_0^{\infty} \frac{1}{(1+x)(1+x^2)} dx$$

Thus $I = I_1 - I_2$ equation 1

Solving for I_1

$$I_1 = \int_0^{\infty} \frac{1}{(1+x^2)} dx$$

$$I_1 = [\tan^{-1} x]_0^{\infty}$$

$$\text{since } \int \frac{1}{(1+x^2)} dx = \tan^{-1} x$$

$$I_1 = [\tan^{-1}(\infty) - \tan^{-1}(0)]$$

$$I_1 = \pi/2 \text{equation 2}$$

Solving for I_2

$$I_2 = \int_0^{\infty} \frac{1}{(1+x)(1+x^2)} dx$$

$$\text{Let } \frac{1}{(1+x)(1+x^2)} = \frac{a}{1+x} + \frac{bx+c}{1+x^2} \text{equation 3}$$

$$\frac{1}{(1+x)(1+x^2)} = \frac{a(1+x^2) + (bx+c)(1+x)}{(1+x)(1+x^2)}$$

$$\frac{1}{(1+x)(1+x^2)} = \frac{ax^2 + a + bx^2 + bx + cx + c}{(1+x)(1+x^2)}$$

$$a+b=0; a+c=1; b+c=0$$

solving we get

$$a=c=1/2$$

$$b=-1/2$$

substituting the values in equation 3

$$\frac{1}{(1+x)(1+x^2)} = \frac{\frac{1}{2}}{1+x} + \frac{-\frac{1}{2}x + \frac{1}{2}}{1+x^2}$$

$$\frac{1}{(1+x)(1+x^2)} = \frac{\frac{1}{2}}{1+x} + \frac{-\frac{1}{2}x}{1+x^2} + \frac{1/2}{1+x^2}$$

Thus substituting the values in I_2 , thus

$$I_2 = \int_0^{\infty} \frac{\frac{1}{2}}{1+x} + \frac{-\frac{1}{2}x}{1+x^2} + \frac{1/2}{1+x^2} dx$$

$$I_2 = \int_0^{\infty} \frac{1}{1+x} dx + \int_0^{\infty} \frac{-\frac{1}{2}x}{1+x^2} dx + \int_0^{\infty} \frac{1/2}{1+x^2} dx$$

Solving :

$$\int_0^{\infty} \frac{-\frac{1}{2}x}{1+x^2} dx$$

$$\text{Let } 1 + x^2 = y$$

$$2x dx = dy$$

For $x = \infty$

$$y = \infty$$

For $x = 0$

$$y = 0$$

substitution
 $1 \int^{\infty} dy$

$$-\frac{1}{2} \|\log y\|^\infty$$

1

$$I_2 = \int_{-\infty}^{\infty} \frac{1}{\frac{1}{2} + x^2} dx + \int_{-\infty}^{\infty} \frac{-\frac{1}{2}x}{\frac{1}{2} + x^2} dx + \int_{-\infty}^{\infty} \frac{1/2}{\frac{1}{2} + x^2} dx$$

$$I_2 = \frac{1}{2} [\log(1+x)]_0^\infty + -\frac{1}{2} [\log x]_0^\infty + \frac{1}{2} [\tan^{-1}x]_0^\infty$$

Substituting values equation 2 and equation 4 in equation 1

Thus

$$l_1 \equiv l_a - l_b$$

$$l = \pi/2 - \pi/4$$

1 - 11

11. Question

Evaluate the following integral:

$$\int_{\frac{\pi}{2}}^{\pi} \frac{x \tan x}{\sec x \cosec x} dx$$

Answer

Let us assume $I = \int_0^{\pi} \frac{x \tan x}{\sec x \cosec x} dx$equation 1

By property, we know that $\int_a^b f(x)dx = \int_a^b f(a + b - x)dx$

THUS

$$I = \int_0^{\pi} \frac{(\pi - x) \tan(\pi - x)}{\sec(\pi - x) \cosec(\pi - x)} dx$$

We know

$$\tan(\pi - x) = -\tan x$$

$$\sec(\pi - x) = -\sec x$$

$$\operatorname{cosec}(\pi - x) = \operatorname{cosecx}$$

Thus substituting values

Adding equation 1 and 2

$$2I = \int_0^{\pi} \frac{(x)(\tan x)}{\sec x \cosec x} dx + \int_0^{\pi} \frac{(\pi - x)(-\tan x)}{-\sec x \cosec x} dx$$

We know

$$\int_m^n [f(x) + g(x)]dx = \int_m^n f(x)dx + \int_m^n g(x)dx$$

Thus

$$2I = \int_0^{\pi} \frac{(\pi)(\tan x)}{\sec x \cosec x} dx$$

We know

$$\tan \theta = \frac{\sin \theta}{\cos \theta}$$

$$\sec \theta = \frac{1}{\cos \theta}$$

$$\operatorname{cosec} \theta = \frac{1}{\sin \theta}$$

Substituting the values we have

$$2I = \pi \int_0^{\pi} \sin^2 x \, dx$$

$$2I = \pi \int_0^{\frac{\pi}{2}} \frac{1 - \cos 2x}{2} dx \text{ by trigonometric formula}$$

$$2I = \frac{\pi}{2} \left[x - \frac{\sin 2x}{2} \right]_0^{\pi}$$

We know $[f(x)]_a^b = f(b) - f(a)$ b and a being the upper and lower limits respectively.

$$2I = \frac{\pi}{2} \left[\left[\pi - \frac{\sin 2\pi}{2} \right] - [0] \right]$$

$$I = \frac{\pi^2}{4}$$

12. Question

Evaluate the following integral:

$$\int_0^{\pi} x \sin x \cos^4 x \, dx$$

Answer

Let us assume $I = \int_0^{\pi} x \sin x \cos^4 x \, dx$equation 1

By property, we know that $\int_a^b f(x) \, dx = \int_a^b f(a + b - x) \, dx$

$$I = \int_0^{\pi} (\pi - x) \sin(\pi - x) \cos^4(\pi - x) \, dx$$

$$I = \int_0^{\pi} (\pi - x) \sin x \cos^4 x \, dx \text{equation 2}$$

Adding equation 1 and equation 2

$$2I = \int_0^{\pi} x \sin x \cos^4 x \, dx + \int_0^{\pi} (\pi - x) \sin x \cos^4 x \, dx$$

We know

$$\int_m^n [f(x) + g(x)] \, dx = \int_m^n f(x) \, dx + \int_m^n g(x) \, dx$$

$$2I = \int_0^{\pi} \pi \sin x \cos^4 x \, dx \text{equation 3}$$

Let $\cos x = y$

Differentiating both sides

$$-\sin x \, dx = dy$$

$$\sin x \, dx = -dy$$

for $x = 0$

$$\cos 0 = y$$

$$1 = y$$

For $x = \pi$

$$\cos \pi = y$$

$$-1 = y$$

Substituting equation 3 becomes

$$2I = \int_{-1}^{1} -\pi y^4 \, dy$$

$$2I = \int_{-1}^{1} \pi y^4 \, dy$$

$$2I = \pi \left[\frac{y^5}{5} \right]_{-1}^1$$

$$2I = \pi \left[\frac{(1)^5}{5} - \frac{(-1)^5}{5} \right]$$

$$2I = 2\pi/5$$

$$I = \pi/5$$



13. Question

Evaluate the following integral:

$$\int_0^{\pi} x \sin^3 x dx$$

Answer

Let us assume $I = \int_0^{\pi} x \sin^3 x dx$

$$I = \int_0^{\pi} x \sin^2 x \sin x dx$$

By property we know that $\int_a^b f(x)dx = \int_a^b f(a + b - x)dx$

Thus

$$I = \int_0^{\pi} (\pi - x)(1 - \cos^2(\pi - x))\sin(\pi - x) dx$$

$$I = \int_0^{\pi} (\pi - x)(1 - \cos^2 x) \sin x dx \dots \text{equation 2}$$

Adding equation 1 and equation 2

$$2I = \int_0^{\pi} x(1 - \cos^2 x) \sin x dx + \int_0^{\pi} (\pi - x)(1 - \cos^2 x) \sin x dx$$

We know

$$\int_m^n [f(x) + g(x)]dx = \int_m^n f(x)dx + \int_m^n g(x)dx$$

$$2I = \int_0^{\pi} \pi(1 - \cos^2 x) \sin x dx \dots \text{equation}$$

Let $\cos x = y$

Differentiating both sides

$$-\sin x dx = dy$$

$$\sin x dx = -dy$$

for $x = 0$

$$\cos\theta = \gamma$$

$$1 = \sqrt{}$$

For $x = \pi$

$$\cos \pi = -1$$

$$= 1 = \sqrt{}$$

Substituting equation 3 becomes

$$2I = \int_{-1}^1 -\pi(1-y^2) dy$$

$$2I = \int_{-1}^1 \pi(1-y^2) dy$$

$$2I = \pi \left[y - \frac{y^3}{3} \right]_{-1}^1$$

$$2I = \pi \left[\frac{3y - y^3}{3} \right]_{-1}^1$$

$$2I = \pi[\{3(1) - (1)^3\} - \{3(-1) - (-1)^3\}]/3$$

$$2I = \pi[2 - \{-3 + 1\}]/3$$

$$2l = \pi[2 + 2]/3$$

$$l = 2\pi/3$$

14. Question

Evaluate the following integral:

$$\int_0^{\pi} x \log \sin x \, dx$$

Answer

Let us assume $I = \int_0^{\pi} x \log \sin x \, dx$equation 1

By property, we know that $\int_a^b f(x)dx = \int_a^b f(a + b - x)dx$

Adding equation 1 and equation 2

$$2I = \int_0^{\pi} x \log \sin x \, dx + \int_0^{\pi} (\pi - x) \log \sin(\pi - x) \, dx$$

We know

$$\int_m^n [f(x) + g(x)]dx = \int_m^n f(x)dx + \int_m^n g(x)dx$$

$$2I = \int_0^{\pi} \pi \log \sin x \, dx \quad \dots \text{equation 3}$$

We know $\int_0^{2a} f(x) dx = \int_0^a f(x) dx + \int_0^a f(2a-x) dx$

$$= 2 \int_0^a f(x) dx \text{ if } f(2a - x) = f(x)$$

$$= 0 \text{ if } f(2a - x) = -f(x)$$

Thus equation 3 becomes

$$2I = 2\pi \int_0^{\frac{\pi}{2}} \log \sin x \, dx \quad \dots \text{equation 4 since } \log \sin(\pi - x) = \log \sin x$$

By property, we know that $\int_a^b f(x)dx = \int_a^b f(a + b - x)dx$

$$2I = 2\pi \int_0^{\frac{\pi}{2}} \log \sin\left(\frac{\pi}{2} - x\right) dx$$

$$2I = 2\pi \int_{0}^{\frac{\pi}{2}} \log \cos x \, dx \quad \dots \text{equation 5}$$

Adding equation 4 and equation 5

$$4I = 2\pi \int_0^{\frac{\pi}{2}} \log \sin x \, dx + 2\pi \int_0^{\frac{\pi}{2}} \log \cos x \, dx$$

We know

$$\int_m^n [f(x) + g(x)] dx = \int_m^n f(x) dx + \int_m^n g(x) dx$$

$$4I = 2\pi \int_0^{\frac{\pi}{2}} \log \sin x + \log \cos x \, dx$$

We know $\log m + \log n = \log mn$ thus

$$4I = 2\pi \int_0^{\frac{\pi}{2}} \log[(2\sin x \cos x)/2] \, dx$$

$$2I = \pi \int_0^{\frac{\pi}{2}} \log[(\sin 2x)/2] \, dx$$

$$2I = \pi \int_0^{\frac{\pi}{2}} \log \sin 2x - \log 2 \, dx \text{ since } \log(m/n) = \log m - \log n$$

$$2I = \pi \int_0^{\frac{\pi}{2}} \log \sin 2x \, dx - \pi \int_0^{\frac{\pi}{2}} \log 2 \, dx \dots \text{equation 6}$$

$$\text{Let } I_1 = \pi \int_0^{\frac{\pi}{2}} \log \sin 2x \, dx$$

$$\text{Let } 2x = y$$

$$2dx = dy$$

$$dx = dy/2$$

$$\text{For } x = 0$$

$$y = 0$$

$$\text{for } x = \frac{\pi}{2}$$

$$y = \pi$$

thus substituting value in I_1

$$I_1 = \frac{1}{2} \int_0^{\pi} \pi \log \sin y \, dy$$

From equation 3 we get

$$I_1 = \frac{1}{2}(2I)$$

$$I_1 = I$$

Thus substituting the value of I_1 in equation 6

$$2I = I - \pi \int_0^{\frac{\pi}{2}} \log 2 \, dx$$

$$I = -\pi \log 2 \int_0^{\frac{\pi}{2}} 1 \, dx$$

$$I = -\pi \log 2 \left[x \right]_0^{\frac{\pi}{2}}$$

$$I = -\pi \log_2 \left[\frac{\pi}{2} \right]$$

$$I = -\log_2 \left[\frac{\pi^2}{2} \right]$$

15. Question

Evaluate the following integral:

$$\int_0^{\pi} \frac{x \sin x}{1 + \sin x} dx$$

Answer

Let us assume $I = \int_0^{\pi} \frac{x \sin x}{1 + \sin x} dx$equation 1

By property, we know that $\int_a^b f(x)dx = \int_a^b f(a + b - x)dx$

thus

$$I = \int_0^{\pi} \frac{(\pi-x)\sin(\pi-x)}{1 + \sin(\pi-x)} dx \dots \text{equation 2}$$

Since $\sin(\pi - x) = \sin x$

Adding equation 1 and equation 2

$$2I = \int_0^{\pi} \frac{x \sin x}{1 + \sin x} dx + \int_0^{\pi} \frac{(\pi - x) \sin x}{1 + \sin x} dx$$

We know

$$\int_m^n [f(x) + g(x)]dx = \int_m^n f(x)dx + \int_m^n g(x)dx$$

Thus

$$2I = \int_0^{\pi} \frac{x \sin x}{1 + \sin x} + \frac{(\pi - x) \sin x}{1 + \sin x} dx$$

$$2I = \int_0^{\pi} \frac{x \sin x + (\pi - x) \sin x}{1 + \sin x} dx$$

$$2I = \int_0^{\pi} \frac{\pi \sin x}{1 + \sin x} dx$$

Adding and subtracting 1

$$2I = \pi \int_0^{\pi} \frac{\sin x + 1 - 1}{1 + \sin x} dx$$

$$2I = \pi \left(\int_0^{\frac{\pi}{2}} \frac{\sin x + 1}{1 + \sin x} + \frac{-1}{1 + \sin x} dx \right)$$

We know

$$\int_{-\infty}^n [f(x) + g(x)] dx = \int_{-\infty}^n f(x) dx + \int_{-\infty}^n g(x) dx$$

$$2I = \pi \left(\int_0^{\pi} \frac{\sin x + 1}{1 + \sin x} dx + \int_0^{\pi} \frac{-1}{1 + \sin x} dx \right)$$

$$\text{Let } I_1 = \int_0^{\pi} \frac{\sin x + 1}{1 + \sin x} dx$$

$$\text{Let } I_2 = \int_0^{\pi} \frac{-1}{1 + \sin x} dx$$

$$2I = \pi(I_1 + I_2) \dots \text{equation 3}$$

Solving I_1 :

$$I_1 = \int_0^{\pi} 1 dx$$

$$I_1 = [x]_0^{\pi}$$

We know $[f(x)]_a^b = f(b) - f(a)$ b and a being the upper and lower limits respectively.

$$I_1 = [\pi - 0]$$

$$I_1 = \pi$$

Solving I_2 :

$$I_2 = \int_0^{\pi} \frac{-1}{1 + \sin x} dx$$

Using trigonometric identity and formula

$$I_2 = \int_0^{\pi} \frac{-1}{\sin^2(\frac{x}{2}) + \cos^2(\frac{x}{2}) + 2 \sin(\frac{x}{2}) \cos(\frac{x}{2})} dx$$

$$I_2 = \int_0^{\pi} \frac{-1}{[\sin(\frac{x}{2}) + \cos(\frac{x}{2})]^2} dx$$

Taking $\cos(x/2)$ common

$$I_2 = \int_0^{\pi} \frac{-1}{\cos^2(\frac{x}{2})[\sin(\frac{x}{2})/\cos(\frac{x}{2}) + 1]^2} dx$$

$$I_2 = \int_0^{\pi} \frac{-1}{\cos^2(\frac{x}{2})[\tan(\frac{x}{2}) + 1]^2} dx$$

$$I_2 = \int_0^{\pi} \frac{-\sec^2(\frac{x}{2})}{[\tan(\frac{x}{2}) + 1]^2} dx$$

$$\text{Let } \tan(\frac{x}{2}) + 1 = y$$

Differentiating both sides, we get,

$$\frac{1}{2} \sec^2(\frac{x}{2}) dx = dy$$

$$\sec^2(\frac{x}{2}) dx = 2dy$$

For $x = 0$

$$\tan(\frac{0}{2}) + 1 = y$$

$$1 = y$$

For $x = \pi$

$$\tan\left(\frac{\pi}{2}\right) + 1 = y$$

$$\infty + 1 = y$$

$$\infty = y$$

Substituting the values

Thus

$$I_2 = \int_1^{\infty} \frac{-2}{y^2} dy$$

$$I_2 = -2 \int_1^{\infty} y^{-2} dy$$

$$I_2 = -2 \left[\frac{y^{-2+1}}{-2+1} \right]_1^{\infty}$$

We know $[f(x)]_a^b = f(b) - f(a)$ b and a being the upper and lower limits respectively

$$I_2 = 2 \left[\frac{1}{\infty} - \frac{1}{1} \right]$$

$$I_2 = -2$$

Substituting values in equation 3

$$2I = \pi(\pi - 2)$$

$$I = \pi\left(\frac{\pi}{2} - 1\right)$$

16. Question

Evaluate the following integral:

$$\int_0^{\pi} \frac{x}{1 + \cos \alpha \sin x} dx, 0 < \alpha < \pi$$

Answer

$$\text{Let } I = \int_0^{\pi} \frac{x}{1 + \cos \alpha \sin x} dx$$

We know that,

$$\int_0^{2a} f(x) dx = \int_0^a f(x) dx + \int_0^a f(2a-x) dx$$

Therefore,

$$I = \int_0^{\frac{\pi}{2}} \frac{x}{1 + \cos \alpha \sin x} dx + \int_0^{\frac{\pi}{2}} \frac{\pi-x}{1 + \cos \alpha \sin x} dx$$

$$2I = \int_0^{\frac{\pi}{2}} \frac{\pi}{1 + \cos \alpha \sin x} dx$$

$$2I = 2\pi \int_0^{\frac{\pi}{2}} \frac{1}{1 + \cos \alpha \sin x} dx$$

$$I = \frac{\pi \alpha}{\sin \alpha}$$

17. Question

Evaluate the following integral:

$$\int_0^{\pi} x \cos^2 x \, dx$$

Answer

Let us assume $I = \int_0^{\pi} x \cos^2 x \, dx$equation 1

By property, we know that $\int_a^b f(x) \, dx = \int_a^b f(a + b - x) \, dx$

Thus

$$I = \int_0^{\pi} (\pi - x) \cos^2(\pi - x) \, dx$$

We know $\cos(\pi - x) = -\cos x$

Thus

$$I = \int_0^{\pi} (\pi - x) \cos^2(x) \, dx \text{equation 2}$$

Adding equation 1 and equation 2

$$2I = \int_0^{\pi} x \cos^2 x \, dx + \int_0^{\pi} (\pi - x) \cos^2 x \, dx$$

We know $\int_m^n [f(x) + g(x)] \, dx = \int_m^n f(x) \, dx + \int_m^n g(x) \, dx$

Thus

$$2I = \int_0^{\pi} (x \cos^2 x + (\pi - x) \cos^2 x) \, dx$$

$$2I = \int_0^{\pi} (\pi \cos^2 x) \, dx$$

We know

$$2 \cos^2 x = 1 + \cos 2x$$

$$2I = \pi \int_0^{\pi} \frac{1 + \cos 2x}{2} \, dx$$

We know $\int_m^n [f(x) + g(x)] \, dx = \int_m^n f(x) \, dx + \int_m^n g(x) \, dx$

Thus

$$2I = \frac{\pi}{2} \int_0^{\pi} 1 \, dx + \frac{\pi}{2} \int_0^{\pi} \cos 2x \, dx$$

since $\int \cos y \, dy = \sin y$

$$2I = \frac{\pi}{2} [x]_0^{\pi} + \frac{\pi}{2} \left[\frac{\sin 2x}{2} \right]_0^{\pi}$$

We know $[f(x)]_a^b = f(b) - f(a)$ b and a being the upper and lower limits respectively

$$2I = \frac{\pi}{2} [\pi - 0] + \frac{\pi}{2} \left[\frac{\sin 2\pi}{2} - \frac{\sin 0}{2} \right]$$

Thus

$$I = \frac{\pi}{4} [\pi]$$

18. Question

Evaluate the following integral:

$$\int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{1}{1 + \cot^2 x} dx$$

Answer

We know that $\tan x = \frac{\cos x}{\sin x}$ $\cot x = \frac{1}{\tan x}$

Substituting the value in equation 1 we have,

$$I = \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{1}{1 + \left(\frac{\cos x}{\sin x}\right)^{\frac{3}{2}}} dx$$

$$I = \int_{\frac{\pi}{2}}^{\frac{\pi}{3}} \frac{\sin^2 x}{\sqrt{5 \cos^2 x + \sin^2 x}} dx \dots \text{equation 2}$$

By property we know that $\int_a^b f(x)dx = \int_a^b f(a + b - x)dx$

Thus in equation 2 $\frac{\pi}{3} + \frac{\pi}{6} = \frac{\pi}{2}$

$$I = \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{\sin^3(\frac{\pi}{2} - x)}{\sin^2(\frac{\pi}{2} - x) + \cos^2(\frac{\pi}{2} - x)} dx$$

$$I = \int_{\frac{\pi}{2}}^{\frac{\pi}{3}} \frac{\cos^{\frac{2}{3}} x}{\sin^{\frac{2}{3}} x + \cos^{\frac{2}{3}} x} dx \dots \text{equation 3}$$

Adding equation 2 and 3

$$2I = \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{\sin^3 x}{\cos^3 x + \sin^3 x} dx + \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{\cos^3 x}{\sin^3 x + \cos^3 x} dx$$

We know

$$\int_{-n}^n [f(x) + g(x)] dx = \int_{-n}^n f(x) dx + \int_{-n}^n g(x) dx$$

Thus

$$2I = \int_{\frac{\pi}{4}}^{\frac{3\pi}{4}} \frac{\sin^2 x + \cos^2 x}{\cos^2 x + \sin^2 x} dx$$

$$2I = \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} 1 \, dx$$

$$2I = [x]_0^{\frac{\pi}{6}}$$

$$2I = \left[\frac{\pi}{3} - \frac{\pi}{6} \right]$$

$$I = \frac{\pi}{12}$$

19. Question

Evaluate the following integral:

$$\int_0^{\frac{\pi}{2}} \frac{\tan^7 x}{\tan^7 x + \cot^7 x} dx$$

Answer

$$\text{Let us assume } I = \int_0^{\frac{\pi}{2}} \frac{\tan^7 x}{\tan^7 x + \cot^7 x} dx \dots\dots \text{equation 1}$$

$$\text{By property, we know that } \int_a^b f(x) dx = \int_a^b f(a + b - x) dx$$

$$\text{We know } \tan\left(\frac{\pi}{2} - x\right) = \cot x$$

$$\cot\left(\frac{\pi}{2} - x\right) = \tan x$$

Thus substituting the values in equation 1

$$I = \int_0^{\frac{\pi}{2}} \frac{\cot^7 x}{\cot^7 x + \tan^7 x} dx \dots\dots \text{equation 2}$$

Adding equation 1 and equation 2

$$2I = \int_0^{\frac{\pi}{2}} \frac{\tan^7 x}{\tan^7 x + \cot^7 x} dx + \int_0^{\frac{\pi}{2}} \frac{\cot^7 x}{\tan^7 x + \cot^7 x} dx$$

$$\text{We know } \int_m^n [f(x) + g(x)] dx = \int_m^n f(x) dx + \int_m^n g(x) dx$$

Thus

$$2I = \int_0^{\frac{\pi}{2}} \frac{\tan^7 x}{\tan^7 x + \cot^7 x} + \frac{\cot^7 x}{\tan^7 x + \cot^7 x} dx$$

$$2I = \int_0^{\frac{\pi}{2}} \frac{\tan^7 x + \cot^7 x}{\tan^7 x + \cot^7 x} dx$$

$$2I = \int_0^{\frac{\pi}{2}} 1. dx$$

$$2I = \int_0^{\frac{\pi}{2}} 1. dx$$

$$2I = [x]_0^{\frac{\pi}{2}}$$



$$2I = \left[\frac{\pi}{2} - 0 \right]$$

$$I = \frac{\pi}{4}$$

20. Question

Evaluate the following integral:

$$\int_2^8 \frac{\sqrt{10-x}}{\sqrt{x} + \sqrt{10-x}} dx$$

Answer

Let us assume $I = \int_2^8 \frac{\sqrt{10-x}}{\sqrt{x} + \sqrt{10-x}} dx \dots\dots\text{equation 1}$

By property, we know that $\int_a^b f(x) dx = \int_a^b f(a + b - x) dx$

thus

$$I = \int_2^8 \frac{\sqrt{10-(8+2-x)}}{\sqrt{(8+2-x)} + \sqrt{10-(8+2-x)}} dx \dots\dots\text{equation 2}$$

Adding equation 1 and 2 $2I = \int_2^8 \frac{\sqrt{10-x}}{\sqrt{x} + \sqrt{10-x}} dx + \int_2^8 \frac{\sqrt{x}}{\sqrt{10-x} + \sqrt{x}} dx$

We know

$$\int_m^n [f(x) + g(x)] dx = \int_m^n f(x) dx + \int_m^n g(x) dx$$

$$2I = \int_2^8 \frac{\sqrt{x} + \sqrt{10-x}}{\sqrt{x} + \sqrt{10-x}} dx$$

$$2I = \int_2^8 1 dx$$

We know $[f(x)]_a^b = f(b) - f(a)$

$$2I = [x]_2^8$$

$$2I = [8 - 2]$$

$$2I = 6$$

$$I = 3$$

21. Question

Evaluate the following integral:

$$\int_0^\pi x \sin x \cos^2 x dx$$

Answer

Let us assume $I = \int_0^\pi x \sin x \cos^2 x dx \dots\dots\text{equation 1}$

By property, we know that $\int_a^b f(x) dx = \int_a^b f(a + b - x) dx$

Thus

Adding equation 1 and equation 1

$$2I = \int_0^{\pi} x \sin x \cos^2 x dx + \int_0^{\pi} (\pi - x) \sin(\pi - x) \cos^2(\pi - x) dx$$

We know $\int_m^n [f(x) + g(x)] dx = \int_m^n f(x) dx + \int_m^n g(x) dx$

$$2I = \int_0^{\pi} x \sin x \cos^2 x + (\pi - x) \sin(\pi - x) \cos^2(\pi - x) dx$$

We know $\sin(\pi - x) = \sin x$

$$\cos(\pi - x) = -\cos x$$

$$2I = \int_0^{\pi} (\pi - x + x) \sin x \cos^2 x dx$$

Let $\cos x = y$

Differentiating both sides

$$-\sin x \, dx = dy$$

$$\sin x \, dx = -dy$$

For $x = 0$

$$\cos x = y$$

$$\cos \theta = y$$

y = 1

for $x = \pi$

$$\cos \pi = -1$$

$$v = -1$$

Substituting the given values

$$2I = \int_{-1}^1 -(\pi)y^2 dy$$

We know that $\int_a^b -f(x) \, dx = \int_b^a f(x) \, dx$

$$2I = \int_{-1}^1 (\pi)y^2 dy$$

$$2l = \pi \left[\frac{y^3}{3} \right]_1$$

We know $[f(x)]^b = f(b) - f(a)$ b and a being the upper and lower limits respectively

$$2I = \pi \left[\frac{1^3}{3} - \frac{(-1)^3}{3} \right]$$

$$2l = \pi \left[\frac{2}{3} \right]$$

$$I = \pi \left[\frac{1}{3} \right]$$

22. Question

Evaluate the following integral:

$$\int_0^{\frac{\pi}{2}} \frac{x \sin x \cos x}{\sin^4 x + \cos^4 x} dx$$

Answer

Let us assume $I = \int_0^{\frac{\pi}{2}} \frac{x \sin x \cos x}{\sin^4 x + \cos^4 x} dx \dots\dots$ equation 1

By property, we know that $\int_a^b f(x) dx = \int_a^b f(a + b - x) dx$

$$I = \int_0^{\frac{\pi}{2}} \frac{\frac{\pi}{2} - x}{\sin^4(\frac{\pi}{2} - x) + \cos^4(\frac{\pi}{2} - x)} dx$$

$$I = \int_0^{\frac{\pi}{2}} \frac{\frac{\pi}{2} - x}{\cos^4 x + \sin^4 x} dx \dots\dots$$
 equation 2

Adding equation 1 and 2

Thus

$$2I = \int_0^{\frac{\pi}{2}} \frac{x \sin x \cos x}{\sin^4 x + \cos^4 x} dx + \int_0^{\frac{\pi}{2}} \frac{\frac{\pi}{2} - x}{\cos^4 x + \sin^4 x} dx$$

We know $\int_m^n [f(x) + g(x)] dx = \int_m^n f(x) dx + \int_m^n g(x) dx$

$$2I = \int_0^{\frac{\pi}{2}} \frac{\frac{\pi}{2} \sin x \cos x}{\sin^4 x + \cos^4 x} dx$$

$$2I = \int_0^{\frac{\pi}{2}} \frac{\frac{\pi}{2} \sin x \cos x}{\cos^4 x (\sin^4 x / \cos^4 x + 1)} dx$$

Since $\tan x = \frac{\sin x}{\cos x}$ and $\frac{1}{\cos x} = \sec x$

$$2I = \int_0^{\frac{\pi}{2}} \frac{\frac{\pi}{2} \sin x}{\cos^3 x (\tan^4 x + 1)} dx$$

$$2I = \int_0^{\frac{\pi}{2}} \frac{\frac{\pi}{2} \tan x \sec^2 x}{(\tan^4 x + 1)} dx \dots\dots$$
 equation 3

Let $\tan^2 x = y$

Differentiating both sides

$$2 \tan x \sec^2 x dx = dy$$

For $x = \frac{\pi}{2}$

$$\tan^2 \frac{\pi}{2} = y$$

$y = \infty$

For $x = 0$

$$\tan^2 0 = y$$



$$y = 0$$

substituting values in equation 3

$$2I = \int_0^{\infty} \frac{\frac{\pi}{2}}{2(y^2 + 1)} dy$$

$$2I = \frac{\pi}{4} [\tan^{-1} y]_0^{\infty} \text{ since } \int \frac{1}{x^2 + 1} dx = \tan^{-1} x$$

$$I = \frac{\pi}{8} [\tan^{-1} \infty - \tan^{-1} 0]$$

$$I = \frac{\pi}{8} \left[\frac{\pi}{2} - 0 \right]$$

$$I = \frac{\pi^2}{16}$$

23. Question

Evaluate the following integral:

$$\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \sin^3 x dx$$

Answer

$$\text{Let us assume } I = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \sin^3 x dx$$

$$I = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \sin^2 x \sin x dx$$

$$I = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} (1 - \cos^2 x) \sin x dx$$

By property, we know that $\int_{-a}^a f(x) dx = \int_0^a f(x) + f(-x) dx$

Thus

$$I = \int_0^{\frac{\pi}{2}} (1 - \cos^2 x) \sin x + (1 - \cos^2(-x)) \sin(-x) dx$$

$$I = \int_0^{\frac{\pi}{2}} (1 - \cos^2 x) \sin x - (1 - \cos^2 x) \sin x dx$$

$$I = 0$$

24. Question

Evaluate the following integral:

$$\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \sin^4 x dx$$

Answer

$$\text{Let us assume } I = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \sin^4 x dx$$

By property, we know that $\int_{-a}^a f(x) dx = \int_0^a f(x) + f(-x) dx$

Thus

$$I = \int_0^{\frac{\pi}{2}} \sin^4 x dx + \sin^4(-x) dx$$

$$I = 2 \int_0^{\frac{\pi}{2}} \sin^4 x dx \dots \text{equation 1}$$

By property, we know that $\int_a^b f(x)dx = \int_a^b f(a + b - x)dx$

Thus

$$I = 2 \int_0^{\frac{\pi}{2}} \sin^4\left(\frac{\pi}{2} - x\right) dx$$

Adding equation 1 and 2

$$2I = 2 \int_0^{\frac{\pi}{2}} \sin^4 x dx + 2 \int_0^{\frac{\pi}{2}} \cos^4 x dx$$

We know $\int_m^n [f(x) + g(x)] dx = \int_m^n f(x) dx + \int_m^n g(x) dx$

$$2I = 2 \int_0^{\frac{\pi}{2}} (\sin^4 x + \cos^4 x) dx$$

$$2I = 2 \int_{\frac{\pi}{2}}^{\pi} (\sin^4 x + \cos^4 x) dx$$

$$2I = 2 \int_{0}^{\frac{\pi}{2}} (\sin^2 x)^2 + (\cos^2 x)^2 dx$$

Since $(a + b)^2 = a^2 + b^2 + 2ab$

$$2I = 2 \int_0^{\frac{\pi}{2}} (\sin^2 x + \cos^2 x)^2 - 2 \sin^2 x \cos^2 x dx$$

$$I = \int_0^{\frac{\pi}{2}} 1 - 2 \sin^2 x \cos^2 x \, dx$$

$$I = \int_{0}^{\frac{\pi}{2}} 1 - [2(2) \sin^2 x \cos^2 x] \frac{1}{2} dx$$

$$I = \int_{0}^{\frac{\pi}{2}} 1 - (\sin^2 2x) \frac{1}{2} dx$$

$$I = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} 1 - (1 - \cos 4x) \frac{1}{2 \times 2} dx$$

$$I = \frac{1}{4} \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} 3 + \cos 4x \ dx$$

$$I = \frac{1}{4} \left[3 + \frac{\sin 4x}{4} \right]^{\frac{\pi}{2}}$$

$$I = \frac{1}{4} \left[\frac{3\pi}{2} + \frac{\sin 2\pi}{4} \right] - [0]$$

$$I = \frac{3}{4} \left[\frac{\pi}{2} \right]$$

25. Question

Evaluate the following integral:

$$\int_{-1}^1 \log \frac{2-x}{2+x} dx$$

Answer

$$\text{Let us assume } I = \int_{-1}^1 \log \frac{2-x}{2+x} dx \dots\dots\text{equation 1}$$

$$\text{By property, we know that } \int_a^{-a} f(x) dx = \int_0^a f(x) dx + \int_0^a f(-x) dx$$

thus

$$I = \int_0^1 \log \frac{2-x}{2+x} dx + \int_0^1 \log \frac{2+x}{2-x} dx$$

$$\text{We know } \int_m^n [f(x) + g(x)] dx = \int_m^n f(x) dx + \int_m^n g(x) dx$$

$$I = \int_0^1 [\log \frac{2-x}{2+x} + \log \frac{2+x}{2-x}] dx$$

Since we know $\log(mn) = \log m + \log n$

$$I = \int_0^1 [\log \frac{2-x}{2+x} * \frac{2+x}{2-x}] dx$$

$$I = \int_0^1 \log 1 dx$$

$$I = \int_0^1 0 dx$$

Thus

$$I = 0$$

26. Question

Evaluate the following integral:

$$\int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \sin^2 x dx$$

Answer

$$\text{Let us assume } I = \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \sin^2 x dx$$

$$\text{By property, we know that } \int_{-a}^a f(x) dx = \int_0^a f(x) + f(-x) dx$$

$$I = \int_0^{\frac{\pi}{4}} \sin^2 x dx + \sin^2(-x) dx$$

$$I = 2 \int_0^{\frac{\pi}{4}} \sin^2 x dx$$

$$I = 2 \int_0^{\frac{\pi}{4}} \frac{1 - \cos 2x}{2} dx$$

$$I = \int_0^{\frac{\pi}{4}} 1 - \cos 2x \, dx$$

$$I = \left[x - \frac{\sin 2x}{2} \right]_0^{\frac{\pi}{4}}$$

$$I = \left[\frac{\pi}{4} - \frac{\sin(2\frac{\pi}{4})}{2} \right] - [0]$$

$$I = \left[\frac{\pi}{4} - \frac{1}{2} \right]$$

27. Question

Evaluate the following integral:

$$\int_0^{\pi} \log(1 - \cos x) dx$$

Answer

Let us assume $I = \int_0^{\pi} \log(1 - \cos x) dx$

$$I = \int_0^{\pi} \log(2 \sin^2 x) dx$$

$$I = \int_0^{\pi} \log(2) + \log(\sin^2 x) dx \text{ since } \log(mn) = \log m + \log n \text{ and } \log(m^n) = n \log m$$

$$I = \int_0^{\pi} \log 2 dx + \int_0^{\pi} \log \sin^2 x dx$$

$$\text{Let } I_1 = \int_0^{\pi} \log \sin x dx$$

We know that $\int_0^{2a} f(x) dx = \int_0^a f(x) dx + \int_0^a f(2a - x) dx$

$$\text{If } f(2a - x) = f(x)$$

than $\int_0^{2a} f(x)dx = 2 \int_0^a f(x)dx$

thus

$$I_1 = 2 \int_0^{\frac{\pi}{2}} \log \sin x \, dx \dots \text{equation 1}$$

since $\log \sin(\pi - x) = \log \sin x$

By property, we know that $\int_a^b f(x)dx = \int_a^b f(a + b - x)dx$

$$I_1 = 2 \int_0^{\frac{\pi}{2}} \log \sin\left(\frac{\pi}{2} - x\right) dx$$

$$I_1 = 2 \int_0^{\frac{\pi}{2}} \log \cos x \, dx \quad \dots \dots \text{equation 2}$$

Adding equation 1 and equation 2

$$2I_1 = 2 \int_0^{\frac{\pi}{2}} \log \sin x \, dx + 2 \int_0^{\frac{\pi}{2}} \log \cos x \, dx$$

We know

$$\int_m^n [f(x) + g(x)]dx = \int_m^n f(x)dx + \int_m^n g(x)dx$$

$$2I_1 = 2 \int_0^{\frac{\pi}{2}} \log \sin x + \log \cos x \, dx$$

We know $\log m + \log n = \log mn$ thus

$$I_1 = \int_{0}^{\frac{\pi}{2}} \log[(2\sin x \cos x)/2] dx$$

$$I_1 = \int_0^{\frac{\pi}{2}} \log[(\sin 2x)/2] dx$$

$$I_1 = \int_{\frac{\pi}{2}}^{\pi} \log \sin 2x - \log 2 \, dx \text{ since } \log(m/n) = \log m - \log n$$

$$I_1 = \int_{\frac{\pi}{2}}^{\pi} \log \sin 2x dx - \int_{\frac{\pi}{2}}^{\pi} \log 2 dx \quad \dots \text{equation 3}$$

$$\text{Let } I_2 = \int_0^{\frac{\pi}{2}} \log \sin 2x dx$$

Let $2x = v$

$$2dx = dy$$

$$dx = dy/2$$

For $x = 0$

$$v = 0$$

for $x = \frac{\pi}{2}$

v = π

thus substituting value in I1

$$I_2 = \frac{1}{2} \int_0^{\pi} \log \sin y dy$$

From equation 3 we get

$$I_2 = \frac{1}{2}(I1)$$

$$I_2 = \frac{I_1}{2}$$

Thus substituting the value of I_2 in equation 3

$$I_1 = I_2 - \int_{\frac{\pi}{2}}^{\pi} \log 2 dx$$

$$\frac{I_1}{2} = -\log 2 \int_0^{\frac{\pi}{2}} 1 dx$$

$$I_1 = -2 \log 2 [x]_0^{\frac{\pi}{2}}$$

$$I_1 = -2 \log 2 \left[\frac{\pi}{2} \right]$$

$$I_1 = -2 \log 2 \left[\frac{\pi}{2} \right]$$

$$I_1 = -\pi \log 2$$

Substituting in equation (a) i.e

$$I = \int_0^{\pi} \log 2 dx + 2 \int_0^{\pi} \log \sin x dx$$

$$I = \int_0^{\pi} \log 2 dx - 2\pi \log 2$$

$$I = \pi \log 2 - 2\pi \log 2$$

$$I = -\pi \log 2$$

28. Question

Evaluate the following integral:

$$\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \log \frac{2 - \sin x}{2 + \sin x} dx$$

Answer

$$\text{Let us assume } I = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \log \frac{2 - \sin x}{2 + \sin x} dx \dots\dots \text{equation 1}$$

$$\text{By property, we know that } \int_a^{-a} f(x) dx = \int_0^a f(x) dx + \int_0^a f(-x) dx$$

thus

$$I = \int_0^{\frac{\pi}{2}} \log \frac{2 - \sin x}{2 + \sin x} dx + \int_0^{\frac{\pi}{2}} \log \frac{2 + \sin x}{2 - \sin x} dx$$

$$\text{We know } \int_m^n [f(x) + g(x)] dx = \int_m^n f(x) dx + \int_m^n g(x) dx$$

$$I = \int_0^{\frac{\pi}{2}} [\log \frac{2 - \sin x}{2 + \sin x} + \log \frac{2 + \sin x}{2 - \sin x}] dx$$

$$\text{Since we know } \log(mn) = \log m + \log n$$

$$I = \int_0^{\frac{\pi}{2}} [\log \frac{2 - \sin x}{2 + \sin x} * \frac{2 + \sin x}{2 - \sin x}] dx$$

$$I = \int_0^{\frac{\pi}{2}} \log 1 dx$$

$$I = \int_0^{\frac{\pi}{2}} 0 \, dx$$

Thus

$$I = 0$$

29. Question

Evaluate the following integral:

$$\int_{-\pi}^{\pi} \frac{2x(1+\sin x)}{1+\cos^2 x} dx$$

Answer

$$\text{Let us assume } I = \int_{-\pi}^{\pi} \frac{2x(1+\sin x)}{1+\cos^2 x} dx \dots\dots\dots\text{equation 1}$$

By property, we know that $\int_a^{-a} f(x) dx = \int_0^a f(x) dx + \int_0^a f(-x) dx$

$$I = \int_0^{\pi} \frac{2x(1+\sin x)}{1+\cos^2 x} dx + \int_0^{\pi} \frac{-2x(1+\sin(-x))}{1+\cos^2(-x)} dx$$

We know that

$$\sin(-x) = -\sin x$$

$$\cos(-x) = \cos x$$

$$\text{We know } \int_m^n [f(x) + g(x)] dx = \int_m^n f(x) dx + \int_m^n g(x) dx$$

Thus substituting the values, we get,

$$I = \int_0^{\pi} \left[\frac{2x(1+\sin x)}{1+\cos^2 x} + \frac{2(-x)(1-\sin x)}{1+\cos^2 x} \right] dx$$

$$I = \int_0^{\pi} \frac{4x(\sin x)}{1+\cos^2 x} dx \dots\dots\dots\text{equation 2}$$

By property, we know that $\int_a^b f(x) dx = \int_a^b f(a+b-x) dx$

$$I = \int_0^{\pi} \frac{4(\pi-x)(\sin(\pi-x))}{1+\cos^2(\pi-x)} dx$$

$$\sin(\pi-x) = \sin x$$

$$\cos(\pi-x) = -\cos x$$

Thus substituting the values

$$I = \int_0^{\pi} \frac{4(\pi-x)(\sin x)}{1+\cos^2 x} dx \dots\dots\dots\text{equation 3}$$

Adding equation 2 and equation 3

$$2I = \int_0^{\pi} \frac{4x(\sin x)}{1+\cos^2 x} dx + \int_0^{\pi} \frac{4(\pi-x)(\sin x)}{1+\cos^2 x} dx$$

$$\text{We know } \int_m^n [f(x) + g(x)] dx = \int_m^n f(x) dx + \int_m^n g(x) dx$$

$$2I = \int_0^{\pi} \left[\frac{4x(\sin x)}{1+\cos^2 x} + \frac{4(\pi-x)(\sin x)}{1+\cos^2 x} \right] dx$$

$$2I = \int_0^{\pi} \frac{4x(\sin x) + 4(\pi - x)(\sin x)}{1 + \cos^2 x} dx$$

$$2I = \int_0^{\pi} \frac{4\pi(\sin x)}{1 + \cos^2 x} dx$$

Let $\cos x = y$

Differentiating both sides

$$-\sin x dx = dy$$

$$\sin x dx = -dy$$

For $x = \pi$

$$\cos \pi = y$$

$$y = -1$$

For $x = 0$

$$\cos 0 = y$$

$$y = 1$$

Thus substituting the given values

$$2I = \int_{-1}^1 \frac{4\pi}{1+y^2} dy \dots \text{equation 4}$$

Now let $y = \tan \theta$

Differentiating both sides

$$dy = \sec^2 \theta d\theta$$

$$\text{For } y = -1$$

$$\tan \theta = -1$$

$$-\frac{\pi}{4} = \theta$$

$$\text{For } y = 1$$

$$\tan \theta = 1$$

$$\theta = \frac{\pi}{4}$$

Substituting the values in equation 4

$$2I = \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \frac{4\pi}{1 + \tan^2 \theta} \sec^2 \theta d\theta$$

$$2I = 4\pi \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} 1 d\theta$$

$$2I = 4\pi [\theta]_{-\frac{\pi}{4}}^{\frac{\pi}{4}}$$

$$2I = 4\pi \left[\frac{\pi}{4} - \left(-\frac{\pi}{4} \right) \right]$$

$$I = \pi^2$$

30. Question

Evaluate the following integral:

$$\int_{-a}^a \log \frac{a - \sin \theta}{a + \sin \theta} dx$$

Answer

$$\text{Let us assume } I = \int_{-a}^a \log \frac{a - \sin \theta}{a + \sin \theta} dx \dots\dots\text{equation 1}$$

$$\text{By property, we know that } \int_a^{-a} f(x) dx = \int_0^a f(x) dx + \int_0^a f(-x) dx$$

thus

$$I = \int_0^a \log \frac{a - \sin \theta}{a + \sin \theta} dx + \int_0^a \log \frac{a + \sin \theta}{a - \sin \theta} dx$$

$$\text{We know } \int_m^n [f(x) + g(x)] dx = \int_m^n f(x) dx + \int_m^n g(x) dx$$

$$I = \int_0^a [\log \frac{a - \sin \theta}{a + \sin \theta} + \log \frac{a + \sin \theta}{a - \sin \theta}] dx$$

Since we know $\log(mn) = \log m + \log n$

$$I = \int_0^a [\log \frac{a - \sin \theta}{a + \sin \theta} * \frac{a + \sin \theta}{a - \sin \theta}] dx$$

$$I = \int_0^a \log 1 dx$$

$$I = \int_0^a 0 dx$$

Thus

$$I = 0$$

31. Question

Evaluate the following integral:

$$\int_{-2}^2 \frac{3x^3 + 2|x| + 1}{x^2 + |x| + 1} dx$$

Answer

$$\int_{-a}^a f(x) dx = 2 \int_0^a f(x) dx$$

And also,

We know that if $f(x)$ is an odd function,

$$\int_{-a}^a f(x) dx = 0$$

As we know the property,

$$\int_{-a}^a f(x)dx = \int_0^a [f(x) + f(-x)]dx$$

Applying this property we get,

$$\int_{-2}^2 \frac{3x^3 + 2|x| + 1}{x^2 + |x| + 1} dx = \int_0^2 \frac{3x^3 + 2|x| + 1}{x^2 + |x| + 1} dx + \int_0^2 \frac{-3x^3 + 2|x| + 1}{x^2 + |x| + 1} dx$$

$$\int_{-2}^2 \frac{3x^3 + 2|x| + 1}{x^2 + |x| + 1} dx = \int_0^2 \frac{2(2x+1)}{x^2 + x + 1} dx$$

$$\text{Let } x^2 + x + 1 = t$$

$$(2x+1)dx = dt$$

And for limits,

$$\text{At } x = 0, t = 1$$

$$\text{At } x = 2, t = 7$$

Therefore, we get,

$$\int_{-2}^2 \frac{3x^3 + 2|x| + 1}{x^2 + |x| + 1} dx = \int_1^7 \frac{2}{t} dt$$

$$\int_{-2}^2 \frac{3x^3 + 2|x| + 1}{x^2 + |x| + 1} dx = 2 [\log(7) - \log(1)]$$

$$\int_{-2}^2 \frac{3x^3 + 2|x| + 1}{x^2 + |x| + 1} dx = 2 \log_e 7$$

32. Question

Evaluate the following integral:

$$\int_{-\frac{3\pi}{2}}^{-\frac{\pi}{2}} \{\sin^2(3\pi+x) + (\pi+x)^3\} dx$$

Answer

$$\text{Let us assume that } I = \int_{-\frac{3\pi}{2}}^{-\frac{\pi}{2}} \{\sin^2(3\pi+x) + (\pi+x)^3\} dx$$

$$\text{We know } \int_m^n [f(x) + g(x)] dx = \int_m^n f(x) dx + \int_m^n g(x) dx$$

$$I = \int_{-\frac{3\pi}{2}}^{-\frac{\pi}{2}} \{\sin^2(3\pi+x)\} dx + \int_{-\frac{3\pi}{2}}^{-\frac{\pi}{2}} (\pi+x)^3 dx$$

$$I = \int_{-\frac{3\pi}{2}}^{-\frac{\pi}{2}} \{\sin^2(3\pi+x)\} dx + \int_{-\frac{3\pi}{2}}^{-\frac{\pi}{2}} (\pi+x)^3 dx$$

$$I = \int_{-\frac{3\pi}{2}}^{-\frac{\pi}{2}} \{1 - \cos 2(3\pi+x)\}/2 dx + \int_{-\frac{3\pi}{2}}^{-\frac{\pi}{2}} (\pi+x)^3 dx$$

$$I = \frac{1}{2} \left[x + \frac{\sin 2(3\pi+x)}{2} \right]_{-\frac{3\pi}{2}}^{-\frac{\pi}{2}} + \left[\frac{(\pi+x)^4}{4} \right]_{-\frac{3\pi}{2}}^{-\frac{\pi}{2}}$$

We know $[f(x)]_a^b = f(b) - f(a)$ b and a being the upper and lower limits respectively thus

$$I = \frac{1}{2} \left[\left\{ -\frac{\pi}{2} + \frac{\sin 2 \left(3\pi - \frac{\pi}{2} \right)}{2} \right\} - \left\{ -\frac{3\pi}{2} + \frac{\sin 2 \left(3\pi - \frac{3\pi}{2} \right)}{2} \right\} \right] \\ + \left[\left\{ \frac{\left(\pi - \frac{\pi}{2} \right)^4}{4} \right\} - \left\{ \frac{\left(\pi - \frac{3\pi}{2} \right)^4}{4} \right\} \right]$$

Thus solving the above equation, we get

$$I = \frac{\pi}{2} + \frac{\left(\frac{\pi}{2} \right)^4}{4} - \frac{\left(-\frac{\pi}{2} \right)^4}{4}$$

$$I = \frac{\pi}{2}$$

33. Question

Evaluate the following integral:

$$\int_0^2 x \sqrt{2-x} dx$$

Answer

$$\text{Let } I = \int_0^2 x \sqrt{2-x} dx \dots \text{equation 1}$$

Put

$$2-x = y^2$$

Differentiating both sides

$$-dx = 2ydy$$

For $x = 2$

$$2-x = y^2$$

$$2-2 = y^2$$

$$y = 0$$

For $x = 0$

$$2-x = y^2$$

$$2-0 = y^2$$

$$y = \sqrt{2}$$

Substituting the values in equation 1

$$I = \int_{\sqrt{2}}^0 -2y(2-y^2)ydy$$

$$I = \int_0^{\sqrt{2}} 2y(2-y^2)ydy$$

$$I = 2 \int_0^{\sqrt{2}} (2y^2 - y^4) dy$$

$$I = 2 \left[\frac{2y^3}{3} - \frac{y^5}{5} \right]_0^{\sqrt{2}}$$

We know $[f(x)]_a^b = f(b) - f(a)$ b and a being the upper and lower limits respectively

thus

$$I = 2 \left[\frac{2(\sqrt{2})^3}{3} - \frac{(\sqrt{2})^5}{5} \right] - [0]$$

Solving this we get

$$I = 2 \left[\frac{(\sqrt{2})^5}{3} - \frac{(\sqrt{2})^5}{5} \right]$$

$$I = 2(\sqrt{2})^5 \left[\frac{1}{3} - \frac{1}{5} \right]$$

$$I = 2(\sqrt{2})^5 \left[\frac{5-3}{15} \right]$$

Thus

$$I = \sqrt{2} \left[\frac{16}{15} \right]$$

34. Question

Evaluate the following integral:

$$\int_0^1 \log\left(\frac{1}{x} - 1\right) dx$$

Answer

$$\text{Let } I = \int_0^1 \log\left(\frac{1}{x} - 1\right) dx$$

$$I = \int_0^1 \log\left(\frac{1-x}{x}\right) dx$$

$$\text{We know } \log\left(\frac{m}{n}\right) = \log m - \log n$$

thus

$$I = \int_0^1 (\log(1-x) - \log(x)) dx \dots\dots \text{equation 2}$$

$$\text{By property, we know that } \int_a^b f(x) dx = \int_a^b f(a + b - x) dx$$

thus

$$I = \int_0^1 (\log(1-(1-x)) - \log(1-x)) dx$$

$$I = \int_0^1 (\log(x) - \log(1-x)) dx \dots\dots \text{equation 3}$$

Adding equation 2 and equation 3 we have

$$2I = \int_0^1 (\log(1-x) - \log(x)) dx + \int_0^1 (\log(x) - \log(1-x)) dx$$

We know $\int_m^n [f(x) + g(x)] dx = \int_m^n f(x) dx + \int_m^n g(x) dx$

Thus on solving we get

$$2I = \int_0^1 0 dx$$

Thus

$$I = 0$$

35. Question

Evaluate the following integral:

$$\int_{-1}^1 |x \cos \pi x| dx$$

Answer

Let $f(x) = |x \cos \pi x|$

Substituting $x = -x$ in $f(x)$

$$f(-x) = |-x \cos(-\pi x)| = |-x \cos(\pi x)| = |x \cos \pi x| = f(x)$$

$$f(x) = f(-x)$$

∴ it is an even function

$$\int_{-1}^1 |x \cos \pi x| dx = 2 \int_0^1 |x \cos \pi x| dx \dots\dots\dots(1)$$

Now,

$$f(x) = |x \cos \pi x| = x \cos \pi x; \text{ for } x \in [0, 1/2]$$

$$= -x \cos \pi x; \text{ for } x \in [1/2, 1]$$

Using interval addition property of integration, we know that

$$\int_a^c f(x) dx = \int_a^b f(x) dx + \int_b^c f(x) dx$$

Equation 1 can be written as

$$2 \left[\int_0^{1/2} x \cos \pi x dx + \int_{1/2}^1 -x \cos \pi x dx \right]$$

Putting the limits in above equation

$$= 2 \{ [(x/\pi) \sin \pi x + (1/\pi^2) \cos \pi x]_0^{1/2} - [(x/\pi) \sin \pi x + (1/\pi^2) \cos \pi x]_{1/2}^1 \}$$

$$= 2 \{ [(1/2\pi) - (1/\pi^2)] - [(-1/\pi^2) - (1/2\pi)] \}$$

$$= 2/\pi$$

36. Question

Evaluate the following integral: $\int_0^\pi \frac{x}{1 + \sin^2 x} + \cos^7 x dx$

Answer

Let us assume $I = \int_0^\pi \frac{x}{1 + \sin^2 x} \cos^7 x dx \dots\dots\dots$ equation 1

By property, we know that $\int_a^b f(x)dx = \int_a^b f(a + b - x)dx$

$$I = \int_0^{\pi} \frac{\pi-x}{1+\sin^2(\pi-x)} + \cos^7(\pi-x) dx. \dots \text{equation 2}$$

Adding equations 1 and 2, we get,

$$2I = \int_0^{\pi} \frac{x}{1 + \sin^2 x} dx + \cos^7 x dx + \int_0^{\pi} \frac{\pi - x}{1 + \sin^2(x)} - \cos^7(x) dx$$

We know $\int_m^n [f(x) + g(x)]dx = \int_m^n f(x)dx + \int_m^n g(x)dx$

Thus

$$2I = \int_0^{\pi} \frac{x}{1 + \sin^2 x} + \cos^7 x + \frac{\pi - x}{1 + \sin^2(x)} - \cos^7(x) dx$$

$$2I = \int_0^{\pi} \frac{\pi}{1 + \sin^2 x} dx$$

We know that $\int_0^{2a} f(x) dx = \int_0^a f(x) dx + \int_a^{2a} f(2a-x) dx$

$$\text{If } f(2a - x) = f(x)$$

than $\int_0^{2a} f(x)dx = 2 \int_0^a f(x)dx$

thus

$$2I = 2 \int_0^{\frac{\pi}{2}} \frac{\pi}{1 + \sin^2 x} dx \text{ since } \sin x = \sin(\pi - x)$$

Now

By property, we know that $\int_a^b f(x)dx = \int_a^b f(a + b - x)dx$

$$2I = 2 \int_0^{\frac{\pi}{2}} \frac{\pi}{1 + \sin^2(\frac{\pi}{2} - x)} dx$$

$$2I = 2 \int_{0}^{\frac{\pi}{2}} \frac{\pi}{1 + \cos^2(x)} dx$$

$$I = \int_{\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{\pi \sec^2 x}{\sec^2(x) + 1} dx$$

since $1/\cos x = \sec x$

$$I = \int_{\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{\pi \sec^2 x}{\tan^2(x) + 2} dx$$

since $\tan^2 x + 1 = \sec^2 x$

Let $\tan x = y$

$$\sec^2 x dx = dy$$

Thus

For $x = \frac{\pi}{3}$

$$\tan \frac{\pi}{2} = y$$

$$y = \infty$$

For $x = 0$

$$\tan 0 = y$$

$$y = 0$$

thus substituting in

$$I = \int_0^{\frac{\pi}{2}} \frac{\pi \sec^2 x}{\tan^2(x) + 2} dx$$

$$I = \int_0^{\infty} \frac{\pi}{y^2 + 2} dy$$

$$I = \int_0^{\infty} \frac{\pi}{2[y^2/(\sqrt{2})^2] + 1} dy$$

$$I = \frac{\pi}{2} [\sqrt{2} \tan^{-1}[y/\sqrt{2}]]_0^{\infty}$$

$$I = \frac{\pi}{\sqrt{2}} [\tan^{-1}(\infty) - \tan^{-1} 0]$$

$$I = \frac{\pi}{\sqrt{2}} \left[\frac{\pi}{2} \right]$$

37. Question

Evaluate the following integral: $\int_0^{\pi} \frac{x}{1 + \sin \alpha \sin x} dx$

Answer

$$\text{Let } I = \int_0^{\pi} \frac{x}{1 + \sin \alpha \sin x} dx$$

We know that,

$$\int_0^{2a} f(x) dx = \int_0^a f(x) dx + \int_0^a f(2a-x) dx$$

Therefore,

$$I = \int_0^{\frac{\pi}{2}} \frac{x}{1 + \sin \alpha \sin x} dx + \int_0^{\frac{\pi}{2}} \frac{\pi-x}{1 + \sin \alpha \sin x} dx$$

$$2I = \int_0^{\frac{\pi}{2}} \frac{\pi}{1 + \sin \alpha \sin x} dx$$

$$2I = 2\pi \int_0^{\frac{\pi}{2}} \frac{1}{1 + \sin \alpha \sin x} dx$$

$$I = \frac{\pi(\frac{\pi}{2} - \alpha)}{\cos \alpha}$$

38. Question

Evaluate the following integral: $\int_0^{2\pi} \sin^{100} x \cos^{101} x dx$

Answer

$$\text{Let us assume } I = \int_0^{2\pi} \sin^{100} x \cos^{101} x dx$$

We know that $\int_0^{2a} f(x)dx = \int_0^a f(x)dx + \int_0^a f(2a-x)dx$

If $f(2a-x) = f(x)$

then $\int_0^{2a} f(x)dx = 2 \int_0^a f(x)dx$

$I = 2 \int_0^\pi \sin^{100} x \cos^{101} x dx$ since $\sin 2\pi - x = -\sin x$ and $\cos 2\pi - x = \cos x$ and $(-\sin x)^{100} = \sin^{100} x$ equation 1

By property, we know that $\int_a^b f(x)dx = \int_a^b f(a+b-x)dx$

$I = 2 \int_0^\pi \sin^{100}(\pi-x) \cos^{101}(\pi-x)dx$

$I = -2 \int_0^\pi \sin^{100} x \cos^{101} x dx$equation 2 since $\cos \pi - x = \cos x$

Adding equation 1 and equation 2

We know $\int_m^n [f(x) + g(x)]dx = \int_m^n f(x)dx + \int_m^n g(x)dx$

thus

$$2I = 2 \int_0^\pi \sin^{100} x \cos^{101} x dx - 2 \int_0^\pi \sin^{100} x \cos^{101} x dx$$

$$2I = 0$$

$$I = 0$$

39. Question

Evaluate the following integral: $\int_0^{\frac{\pi}{2}} \frac{a \sin x + b \cos x}{\sin x + \cos x} dx$

Answer

Let us assume $I = \int_0^{\frac{\pi}{2}} \frac{a \sin x + b \cos x}{\sin x + \cos x} dx$ equation 1

By property, we know that $\int_a^b f(x)dx = \int_a^b f(a+b-x)dx$

Thus

$$I = \int_0^{\frac{\pi}{2}} \frac{a \sin(\frac{\pi}{2}-x) + b \cos(\frac{\pi}{2}-x)}{\sin x + \cos x} dx$$

$$I = \int_0^{\frac{\pi}{2}} \frac{a \cos x + b \sin x}{\sin x + \cos x} dx$$
.....equation 2

Adding the equation 1 and 2

$$2I = \int_0^{\frac{\pi}{2}} \frac{a \sin x + b \cos x}{\sin x + \cos x} dx + \int_0^{\frac{\pi}{2}} \frac{a \cos x + b \sin x}{\sin x + \cos x} dx$$

We know $\int_m^n [f(x) + g(x)]dx = \int_m^n f(x)dx + \int_m^n g(x)dx$

$$2I = \int_0^{\frac{\pi}{2}} \frac{a \sin x + b \cos x + a \cos x + b \sin x}{\sin x + \cos x} dx$$

$$2I = \int_0^{\frac{\pi}{2}} \frac{a \sin x + b \cos x + a \cos x + b \sin x}{\sin x + \cos x} dx$$

$$2I = \int_0^{\frac{\pi}{2}} \frac{a(\sin x + \cos x) + b(\cos x + \sin x)}{\sin x + \cos x} dx$$

$$2I = \int_0^{\frac{\pi}{2}} a + b dx$$

$$2I = a + b[x]_0^{\frac{\pi}{2}}$$

$$I = \frac{(a+b)(\pi)}{2}$$

40. Question

Evaluate the following integrals: $\int_0^{\frac{3}{2}} |x \cos \pi x| dx$

Answer

$$\text{Let us assume } I = \int_0^{\frac{3}{2}} |x \cos \pi x| dx$$

We know $|\cos x| = \cos x$ for $0 < x < \pi/2$ & $|\cos x| = -\cos x$ for $\pi/2 < x < 3\pi/2$

We know that $\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx$ given $a < c < b$

Thus

$$I = \int_0^{\frac{1}{2}} x \cos \pi x dx - \int_{\frac{1}{2}}^{\frac{3}{2}} x \sin \pi x dx$$

By partial integration $\int(u)(v) = (u) \int(v) - \int du \int v$

Thus

$$I = [x \int \cos \pi x dx - \int \frac{dx}{dx} \int \cos \pi x dx]_0^{\frac{1}{2}} - [x \int \cos \pi x dx - \int \frac{dx}{dx} \int \cos \pi x dx]_{\frac{1}{2}}^{\frac{3}{2}}$$

$$I = \left[\frac{x \{\sin \pi x\}}{\pi} + \frac{\cos \pi x}{\pi^2} \right]_0^{\frac{1}{2}} - \left[\frac{x \{\sin \pi x\}}{\pi} + \frac{\cos \pi x}{\pi^2} \right]_{\frac{1}{2}}^{\frac{3}{2}}$$

Since $[f(x)]_a^b = f(b) - f(a)$

$$I = \left[\left[\frac{1/2 \{\sin(\frac{\pi}{2})\}}{\pi} + \frac{\cos(\frac{\pi}{2})}{\pi^2} \right] - \left[\frac{0 \{\sin 0\}}{\pi} + \frac{\cos 0}{\pi^2} \right] \right] - \left[\left[\frac{3/2 \{\sin(\frac{3\pi}{2})\}}{\pi} + \frac{\cos(\frac{3\pi}{2})}{\pi^2} \right] - \left[\frac{1/2 \{\sin(\frac{\pi}{2})\}}{\pi} + \frac{\cos(\frac{\pi}{2})}{\pi^2} \right] \right]$$

$$I = \left[\left[\frac{1}{2\pi} + \frac{0}{\pi^2} \right] - \left[\frac{0 \{0\}}{\pi} + \frac{1}{\pi^2} \right] \right] - \left[\left[\frac{3/2 \{-1\}}{\pi} + \frac{0}{\pi^2} \right] - \left[\frac{1}{2\pi} + \frac{0}{\pi^2} \right] \right]$$

$$I = \left[\left[\frac{1}{2\pi} \right] - \left[\frac{1}{\pi^2} \right] \right] - \left[\left[\frac{-3}{2\pi} \right] - \left[\frac{1}{2\pi} \right] \right]$$

$$I = \frac{5}{2\pi} - \frac{1}{\pi^2}$$

41. Question

Evaluate the following integrals: $\int_0^1 |x \sin \pi x| dx$

Answer

$$\text{Let us assume } I = \int_0^1 |x \sin \pi x| dx$$

We know $|\sin x| = \sin x$ for $0 < x < \pi$ & $|\sin x| = -\sin x$ for $\pi < x < 2\pi$

We know that $\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx$ given $a < c < b$

Thus

$$I = \int_0^1 x \sin \pi x dx$$

By partial integration $\int(u)(v) = (u) \int(v) - \int du \int v$

Thus

$$I = [x \int \sin \pi x dx - \int \frac{dx}{dx} \int \sin \pi x dx]_0^1$$

$$I = \left[\frac{x \{-\cos \pi x\}}{\pi} + \frac{\sin \pi x}{\pi^2} \right]_0^1$$

Since $[f(x)]_a^b = f(b) - f(a)$

$$I = \left[\left[\frac{1 \{-\cos \pi 1\}}{\pi} + \frac{\sin \pi 1}{\pi^2} \right] - \left[\frac{0 \{-\cos \pi 0\}}{\pi} + \frac{\sin \pi 0}{\pi^2} \right] \right]$$

$$I = \left[\left[\frac{1}{\pi} + \frac{0}{\pi^2} \right] - \left[\frac{0 \{-1\}}{\pi} + \frac{0}{\pi^2} \right] \right]$$

$$I = \left[\left[\frac{1}{\pi} \right] \right]$$

$$I = \frac{1}{\pi}$$

42. Question

Evaluate the following integrals: $\int_0^{\frac{3}{2}} |x \sin \pi x| dx$

Answer

$$\text{Let us assume, } I = \int_0^{\frac{3}{2}} |x \sin \pi x| dx$$

We know $|\sin x| = \sin x$ for $0 < x < \pi$ & $|\sin x| = -\sin x$ for $\pi < x < 2\pi$

We know that $\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx$ given $a < c < b$

Thus

$$I = \int_0^1 x \sin \pi x dx - \int_1^{\frac{3}{2}} x \sin \pi x dx$$

By partial integration, $\int(u)(v) = (u)\int(v) - \int du \int v$

Thus,

$$I = \left[x \int \sin \pi x dx - \int \frac{dx}{dx} \int \sin \pi x dx \right]_0^3 - \left[x \int \sin \pi x dx - \int \frac{dx}{dx} \int \sin \pi x dx \right]_1^2$$

$$I = \left[\frac{x\{-\cos nx\}}{\pi} + \frac{\sin nx}{\pi^2} \right]_0^1 - \left[\frac{x\{-\cos nx\}}{\pi} + \frac{\sin nx}{\pi^2} \right]_1^3$$

Since, $[f(x)]_a^b = f(b) - f(a)$

$$I = \left[\left[\frac{1\{-\cos\pi\}}{\pi} + \frac{\sin\pi}{\pi^2} \right] - \left[\frac{0\{-\cos\pi\}}{\pi} + \frac{\sin\pi}{\pi^2} \right] \right] \\ - \left[\left[\frac{3/2\{-\cos\pi/2\}}{\pi} + \frac{\sin\pi/2}{\pi^2} \right] - \left[\frac{1\{-\cos\pi\}}{\pi} + \frac{\sin\pi}{\pi^2} \right] \right]$$

$$I = \left[\left(\frac{1}{\pi} + \frac{0}{\pi^2} \right) - \left(\frac{0\{-1\}}{\pi} + \frac{0}{\pi^2} \right) \right] - \left[\left(\frac{3/2\{0\}}{\pi} + \frac{-1}{\pi^2} \right) - \left(\frac{1}{\pi} + \frac{0}{\pi^2} \right) \right]$$

$$I = \left[\left[\frac{1}{\pi} \right] \right] - \left[\left[\frac{-1}{\pi^2} \right] \right] - \left[\left[\frac{1}{\pi} \right] \right]$$

$$I = \frac{2}{\pi} + \frac{1}{\pi^2}$$

43. Question

If f is an integrable function such that $f(2a - x) = f(x)$, then prove that

$$\int_0^{2a} f(x) dx = 2 \int_0^a f(x) dx$$

Answer

Using interval addition property of integration, we know that

$$\int_a^c f(x) dx = \int_a^b f(x) dx + \int_b^c f(x) dx$$

So L.H.S can be written as,

Let us assume $x = 2a - t$

Differentiating it we get,

$$dx = -dt$$

from above assumption

when $x = 2a \Rightarrow t = 0$

and when $x = a \Rightarrow t = a$

substituting above assumptions in L.H.S

$$\int_a^{2a} f(x)dx = - \int_a^0 f(2a-t)dt$$

Using the property of integration $\int_a^b f(x) dx = - \int_b^a f(x) dx$

$$\int_a^{2a} f(x) dx = \int_0^a f(2a - t) dt$$

Using integration property

$$\int_a^{2a} f(x)dx = \int_a^{2a} f(2a-x)dx$$

Substituting above value in equation 1

$$\int_0^{2a} f(x) dx = \int_0^a f(x) dx + \int_a^{2a} f(2a-x) dx$$

Now using the property $\int_a^b f(x)dx + \int_a^b g(x)dx = \int_a^b \{f(x) + g(x)\}dx$

$$\int_0^{2a} f(x) dx = \int_0^a \{f(x) + f(2a-x)\} dx$$

Since, $f(2a - x) = f(x)$

$$\therefore \int_0^{2a} f(x)dx = \int_0^a \{f(x) + f(x)\}$$

$$\int_0^{2a} f(x)dx = 2 \int_0^a f(x)dx$$

Hence proved.

44. Question

If $f(2a - x) = -f(x)$ prove that $\int_0^{2a} f(x) dx = 0$

Answer

Let us assume $I = \int_0^{2a} f(x)dx$ equation 1

By property, we know that $\int_a^b f(x)dx = \int_a^b f(a + b - x)dx$

Thus

$$I = \int_0^{2a} f(2a - x) dx$$

Given: $f(2a - x) = -f(x)$

Equation 1 becomes

Adding equation 2 and 3

$$2I = \int_0^{2a} f(2a-x)dx + \int_0^{2a} -f(x)dx$$

We know $\int_m^n [f(x) + g(x)]dx = \int_m^n f(x)dx + \int_m^n g(x)dx$

Thus

$$2I = \int_0^{2a} [f(x) - f(x)]dx$$

Thus

$$2I = 0$$

$$I = 0$$

$$\int_0^{2a} f(x)dx = 0$$

45. Question

If f is an integrable function, show that

(i) $\int_{-a}^a f(x^2)dx = 2 \int_0^a f(x^2)dx$

(ii) $\int_{-a}^a xf(x^2)dx = 0$

Answer

(i) Let us check the given function for being even and odd.

$$f((-x)^2) = f(x^2)$$

The function does not change sign and therefore the function is even.

We know that if $f(x)$ is an even function,

$$\int_{-a}^a f(x)dx = 2 \int_0^a f(x)dx$$

Therefore,

$$\int_{-a}^a f(x^2)dx = 2 \int_0^a f(x^2)dx$$

Hence, Proved.

(ii) Let us check the given function for even and odd.

$$\text{Let } g(x) = xf(x^2)$$

$$g(-x) = -x f((-x)^2)$$

$$g(-x) = -xf(x^2)$$

$$g(-x) = -g(x)$$

Therefore, the function is odd.

We know that if $f(x)$ is an odd function,

$$\int_{-a}^a f(x)dx = 0$$

Therefore,

$$\int_{-a}^a xf(x^2)dx = 0$$

Hence, Proved.

46. Question

If $f(x)$ is a continuous function defined on $[0, 2a]$. Then Prove that

$$\int_0^{2a} f(x)dx = \int_0^a \{f(x) + f(2a - x)\}dx$$

Answer

Using interval addition property of integration, we know that

$$\int_a^c f(x)dx = \int_a^b f(x)dx + \int_b^c f(x)dx$$

So L.H.S can be written as,

$$\int_0^{2a} f(x)dx = \int_0^a f(x)dx + \int_a^{2a} f(x)dx \dots \dots (1)$$

Let us assume $x = 2a - t$

Differentiating it we get,

$$dx = -dt$$

from above assumption

$$\text{when } x = 2a \Rightarrow t = 0$$

$$\text{and when } x = a \Rightarrow t = a$$

substituting above assumptions in L.H.S

$$\int_a^{2a} f(x)dx = - \int_a^0 f(2a - t)dt$$

Using the property of integration $\int_a^b f(x)dx = - \int_b^a f(x)dx$

$$\int_a^{2a} f(x)dx = \int_0^a f(2a - t)dt$$

Using integration property

$$\int_a^{2a} f(x)dx = \int_a^{2a} f(2a - x)dx$$

Substituting above value in equation 1

$$\int_0^{2a} f(x)dx = \int_0^a f(x)dx + \int_a^{2a} f(2a - x)dx$$

Now using the property $\int_a^b f(x)dx + \int_a^b g(x)dx = \int_a^b \{f(x) + g(x)\}dx$

$$\int_0^{2a} f(x)dx = \int_0^a \{f(x) + f(2a - x)\}dx$$

$\therefore L.H.S = R.H.S$

Hence, proved.

47. Question

If $f(a + b - x) = f(x)$ prove that:

$$\int_a^b xf(x)dx = (a+b)/2 \int_a^b f(x)dx$$

Answer

LHS

By property, we know that $\int_a^b f(x)dx = \int_a^b f(a + b - x)dx$

thus

$$\int_a^b xf(x)dx = \int_a^b (a + b - x)f(a + b - x)dx$$

Given $f(a + b - x) = f(x)$

$$\int_a^b xf(x)dx = \int_a^b (a + b - x)f(x)dx$$

We know

$$\int_m^n [f(x) + g(x)]dx = \int_m^n f(x)dx + \int_m^n g(x)dx$$

Thus

$$\int_a^b xf(x)dx = \int_a^b (a + b)f(x)dx - \int_a^b xf(x)dx$$

$$2 \int_a^b xf(x)dx = \int_a^b (a + b)f(x)dx$$

$$\int_a^b xf(x)dx = (a + b)/2 \int_a^b f(x)dx$$

Hence proved

48. Question

If $f(x)$ is a continuous function defined on $[-a, a]$, then prove that

$$\int_{-a}^a f(x)dx = \int_0^a \{f(x) + f(-x)\}dx$$

Answer

Using interval addition property of integration, we know that

$$\int_a^c f(x)dx = \int_a^b f(x)dx + \int_b^c f(x)dx$$

So L.H.S can be written as,

$$\int_{-a}^a f(x)dx = \int_{-a}^0 f(x)dx + \int_0^a f(x)dx \dots\dots(1)$$

Now let us take $x = -t$

Differentiating it, we get,

$$dx = -dt$$

from above assumption

when $x = -a \Rightarrow t = a$

and when $x = 0 \Rightarrow t = 0$

Substituting the above assumptions in equation 1 $\int_{-a}^0 f(x)dx = \int_a^0 f(-t)(-dt) = -\int_a^0 f(-t)dt$

Using the property of integration $\int_a^b f(x)dx = -\int_b^a f(x)dx$

$$\int_{-a}^0 f(x)dx = \int_0^a f(-t)dt \dots\dots(2)$$

Using integration property

$$\int_0^a f(-t)dt = \int_0^a f(-x)dx \dots\dots(3)$$

Using equation 2 and 3, now equation 1 can be rewritten as

$$\int_{-a}^a f(x)dx = \int_0^a f(-x)dx + \int_0^a f(x)dx$$

Now using the property $\int_a^b f(x)dx + \int_a^b g(x)dx = \int_a^b \{f(x) + g(x)\}dx$

$$\int_{-a}^a f(x)dx = \int_0^a \{f(-x) + f(x)\}dx$$

$\therefore L.H.S = R.H.S$

Hence proved.

49. Question

Prove that: $\int_0^\pi xf(\sin x)dx = \frac{\pi}{2} \int_0^\pi f(\sin x).dx$

Answer

LHS

Let $I = \int_0^\pi xf(\sin x)dx \dots\dots$ equation 1

By property, we know that $\int_a^b f(x)dx = \int_a^b f(a + b - x)dx$

thus

$$I = \int_0^\pi xf(\sin x).dx = \int_0^\pi (\pi - x)f(\sin(\pi - x)).dx \dots\dots \text{equation 2}$$

Adding equations 1 and equation 2, we get,

$$2I = \int_0^\pi xf(\sin x)dx + \int_0^\pi (\pi - x)f(\sin(\pi - x)).dx$$

Since we know, $\sin(\pi - x) = \sin x$

We know

$$\int_m^n [f(x) + g(x)]dx = \int_m^n f(x)dx + \int_m^n g(x)dx$$

$$2I = \int_0^{\pi} [xf(\sin x) + (\pi - x)f(\sin x)]dx$$

Thus on solving

$$2I = \int_0^{\pi} [(\pi)f(\sin x)]dx$$

We know that by integration property:

$$\int_a^b [(m)f(x)]dx = m \int_a^b [f(x)]dx$$

Thus we have

$$2I = \pi \int_0^{\pi} [f(\sin x)]dx$$

$$I = \frac{\pi}{2} \int_0^{\pi} [f(\sin x)]dx$$

Putting back the value of I we have

$$\int_0^{\pi} xf(\sin x)dx = \frac{\pi}{2} \int_0^{\pi} [f(\sin x)]dx$$

Hence proved

Exercise 20.6

1. Question

Evaluate the following integrals as a limit of sums:

$$\int_0^3 (x + 4)dx$$

Answer

$$\text{To find: } \int_0^3 (x + 4) dx$$

Formula used:

$$\int_a^b f(x) dx = \lim_{h \rightarrow 0} h[f(a) + f(a+h) + f(a+2h) + \dots + f(a+(n-1)h)],$$

where,

$$h = \frac{b-a}{n}$$

Here, $a = 0$ and $b = 3$

Therefore,

$$h = \frac{3-0}{n}$$

$$\Rightarrow nh = 3$$

Let,

$$I = \int_0^3 (x + 4) dx$$

Here, $f(x) = x + 4$ and $a = 0$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[f(0) + f(0+h) + f(0+2h) + \dots + f(0+(n-1)h)]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[f(0) + f(h) + f(2h) + \dots + f((n-1)h)]$$

Now, By putting $x = 0$ in $f(x)$ we get,

$$f(0) = 0 + 4 = 4$$

Similarly, $f(h) = h + 4$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[4 + h + 4 + 2h + 4 + \dots + (n-1)h + 4]$$

In this series, 4 is getting added n times

$$\Rightarrow I = \lim_{h \rightarrow 0} h[4n + h + 2h + \dots + (n-1)h]$$

Now take h common in remaining series

$$\Rightarrow I = \lim_{h \rightarrow 0} h[4n + h(1 + 2 + \dots + (n-1))]$$

$$\left\{ \because \sum_{i=1}^{n-1} i = 1 + 2 + \dots + (n-1) = \frac{n(n-1)}{2} \right\}$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[4n + h \left\{ \frac{n(n-1)}{2} \right\} \right]$$

Put,

$$h = \frac{3}{n}$$

Since,

$$h \rightarrow 0 \text{ and } h = \frac{3}{n} \Rightarrow n \rightarrow \infty$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{3}{n} \left[4n + \frac{3}{n} \left\{ \frac{n(n-1)}{2} \right\} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{3}{n} \left[4n + \frac{3(n-1)}{2} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left\{ 12 + \frac{9(n-1)}{2n} \right\}$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left\{ 12 + \frac{9}{2} \left(1 - \frac{1}{n} \right) \right\}$$

$$\Rightarrow I = 12 + \frac{9}{2} \left(1 - \frac{1}{\infty} \right)$$

$$\Rightarrow I = 12 + \frac{9}{2} (1 - 0)$$

$$\Rightarrow I = 12 + \frac{9}{2}$$

$$\Rightarrow I = \frac{24 + 9}{2}$$

$$\Rightarrow I = \frac{33}{2}$$

Hence, the value of $\int_0^3 (x + 4) dx = \frac{33}{2}$

2. Question

Evaluate the following integrals as a limit of sums:

$$\int_0^2 (x + 3) dx$$

Answer

To find: $\int_0^2 (x + 3) dx$

Formula used:

$$\int_a^b f(x) dx = \lim_{h \rightarrow 0} h[f(a) + f(a+h) + f(a+2h) + \dots + f(a+(n-1)h)],$$

where,

$$h = \frac{b-a}{n}$$

Here, $a = 0$ and $b = 2$

Therefore,

$$h = \frac{2-0}{n}$$

$$\Rightarrow h = \frac{2}{n}$$

Let,

$$I = \int_0^2 (x + 3) dx$$

Here, $f(x) = x + 3$ and $a = 0$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[f(0) + f(0+h) + f(0+2h) + \dots + f(0+(n-1)h)]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[f(0) + f(h) + f(2h) + \dots + f((n-1)h)]$$

Now, By putting $x = 0$ in $f(x)$ we get,

$$f(0) = 0 + 3 = 3$$

Similarly, $f(h) = h + 3$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[3 + h + 3 + 2h + 3 + \dots + (n-1)h + 3]$$

In this series, 3 is getting added n times

$$\Rightarrow I = \lim_{h \rightarrow 0} h[3n + h + 2h + \dots + (n-1)h]$$

Now take h common in remaining series

$$\Rightarrow I = \lim_{h \rightarrow 0} h[3n + h(1 + 2 + \dots + (n-1))]$$

$$\left\{ \therefore \sum_{i=1}^{n-1} i = 1 + 2 + \dots + (n-1) = \frac{n(n-1)}{2} \right\}$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[3n + h \left\{ \frac{n(n-1)}{2} \right\} \right]$$

Put,

$$h = \frac{2}{n}$$

Since,

$$h \rightarrow 0 \text{ and } h = \frac{2}{n} \Rightarrow n \rightarrow \infty$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{2}{n} \left[3n + \frac{2}{n} \left\{ \frac{n(n-1)}{2} \right\} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{2}{n} [3n + (n-1)]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left\{ 6 + \frac{2(n-1)}{n} \right\}$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left\{ 6 + 2 \left(1 - \frac{1}{n} \right) \right\}$$

$$\Rightarrow I = 6 + 2 \left(1 - \frac{1}{\infty} \right)$$

$$\Rightarrow I = 6 + 2(1 - 0)$$

$$\Rightarrow I = 6 + 2$$

$$\Rightarrow I = 8$$

Hence, the value of $\int_0^2 (x+3) dx = 8$

3. Question

Evaluate the following integrals as a limit of sums:

$$\int_1^3 (3x - 2) dx$$

Answer

$$\text{To find: } \int_1^3 (3x - 2) dx$$

Formula used:

$$\int_a^b f(x) dx = \lim_{h \rightarrow 0} h[f(a) + f(a+h) + f(a+2h) + \dots + f(a+(n-1)h)],$$

where,

$$h = \frac{b-a}{n}$$

Here, $a = 1$ and $b = 3$

Therefore,

$$h = \frac{3-1}{n}$$

$$\Rightarrow h = \frac{2}{n}$$

Let,

$$I = \int_1^3 (3x - 2) dx$$

Here, $f(x) = 3x - 2$ and $a = 1$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[f(1) + f(1+h) + f(1+2h) + \dots + f(1+(n-1)h)]$$

Now, By putting $x = 1$ in $f(x)$ we get,

$$f(1) = 3(1) - 2 = 3 - 2 = 1$$

Similarly, $f(1+h)$

$$= 3(1+h) - 2$$

$$= 3 + 3h - 2$$

$$= 3h + 1$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[1 + 3h + 1 + 3(2h) + 1 + \dots + 3(n-1)h + 1]$$

In this series, 1 is getting added n times

$$\Rightarrow I = \lim_{h \rightarrow 0} h[1 \times n + 3h + 3(2h) + \dots + 3(n-1)h]$$

Now take $3h$ common in remaining series

$$\Rightarrow I = \lim_{h \rightarrow 0} h[n + 3h(1 + 2 + \dots + (n-1))]$$

$$\left\{ \because \sum_{i=1}^{n-1} i = 1 + 2 + \dots + (n-1) = \frac{n(n-1)}{2} \right\}$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[n + 3h \left\{ \frac{n(n-1)}{2} \right\} \right]$$

Put,

$$h = \frac{2}{n}$$

Since,

$$h \rightarrow 0 \text{ and } h = \frac{2}{n} \Rightarrow n \rightarrow \infty$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{2}{n} \left[n + \frac{3(2) \left\{ \frac{n(n-1)}{2} \right\}}{n} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{2}{n} \left[n + \frac{6(n-1)}{2} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{2}{n} [n + 3(n-1)]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left\{ 2 + \frac{6(n-1)}{n} \right\}$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left\{ 2 + 6 \left(1 - \frac{1}{n} \right) \right\}$$

$$\Rightarrow I = 2 + 6 \left(1 - \frac{1}{\infty} \right)$$

$$\Rightarrow I = 2 + 6(1 - 0)$$

$$\Rightarrow I = 2 + 6$$

$$\Rightarrow I = 8$$

Hence, the value of $\int_1^3 (3x - 2) dx = 8$

4. Question

Evaluate the following integrals as a limit of sums:

$$\int_{-1}^1 (x + 3) dx$$

Answer

To find: $\int_{-1}^1 (x + 3) dx$

Formula used:

$$\int_a^b f(x) dx = \lim_{h \rightarrow 0} h[f(a) + f(a+h) + f(a+2h) + \dots + f(a+(n-1)h)],$$

where,

$$h = \frac{b-a}{n}$$

Here, $a = -1$ and $b = 1$

Therefore,

$$h = \frac{1 - (-1)}{n}$$

$$\Rightarrow h = \frac{1+1}{n}$$

$$\Rightarrow h = \frac{2}{n}$$

Let,

$$I = \int_{-1}^1 (x + 3) dx$$

Here, $f(x) = x + 3$ and $a = -1$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[f(-1) + f(-1+h) + f(-1+2h) + \dots + f(-1+(n-1)h)]$$

Now, By putting $x = -1$ in $f(x)$ we get,

$$f(-1) = -1 + 3 = 2$$

Similarly, $f(-1 + h)$

$$= -1 + h + 3$$

$$= h + 2$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[2 + h + 2 + 2h + 2 + \dots + (n - 1)h + 2]$$

In this series, 2 is getting added n times

$$\Rightarrow I = \lim_{h \rightarrow 0} h[2n + h + 2h + \dots + (n - 1)h]$$

Now take h common in remaining series

$$\Rightarrow I = \lim_{h \rightarrow 0} h[2n + h(1 + 2 + \dots + (n - 1))]$$

$$\left\{ \because \sum_{i=1}^{n-1} i = 1 + 2 + \dots + (n - 1) = \frac{n(n-1)}{2} \right\}$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[2n + h \left\{ \frac{n(n-1)}{2} \right\} \right]$$

Put,

$$h = \frac{2}{n}$$

Since,

$$h \rightarrow 0 \text{ and } h = \frac{2}{n} \Rightarrow n \rightarrow \infty$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{2}{n} \left[2n + \frac{2}{n} \left\{ \frac{n(n-1)}{2} \right\} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{2}{n} [2n + (n - 1)]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left\{ 4 + \frac{2(n-1)}{n} \right\}$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left\{ 4 + 2 \left(1 - \frac{1}{n} \right) \right\}$$

$$\Rightarrow I = 4 + 2 \left(1 - \frac{1}{\infty} \right)$$

$$\Rightarrow I = 4 + 2(1 - 0)$$

$$\Rightarrow I = 4 + 2$$

$$\Rightarrow I = 6$$

Hence, the value of $\int_{-1}^1 (x + 3) dx = 6$

5. Question

Evaluate the following integrals as a limit of sums:

$$\int_0^5 (x + 1) dx$$

Answer

To find: $\int_0^5 (x + 1) dx$

Formula used:

$$\int_a^b f(x) dx = \lim_{h \rightarrow 0} h[f(a) + f(a+h) + f(a+2h) + \dots + f(a+(n-1)h)],$$

where,

$$h = \frac{b-a}{n}$$

Here, $a = 0$ and $b = 5$

Therefore,

$$h = \frac{5-0}{n}$$

$$\Rightarrow h = \frac{5}{n}$$

Let,

$$I = \int_0^5 (x+1) dx$$

Here, $f(x) = x+1$ and $a = 0$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[f(0) + f(0+h) + f(0+2h) + \dots + f(0+(n-1)h)]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[f(0) + f(h) + f(2h) + \dots + f((n-1)h)]$$

Now, By putting $x = 0$ in $f(x)$ we get,

$$f(0) = 0 + 1 = 1$$

Similarly, $f(h) = h + 1$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[1 + h + 1 + 2h + 1 + \dots + (n-1)h + 1]$$

In this series, 1 is getting added n times

$$\Rightarrow I = \lim_{h \rightarrow 0} h[1 \times n + h + 2h + \dots + (n-1)h]$$

Now take h common in remaining series

$$\Rightarrow I = \lim_{h \rightarrow 0} h[n + h(1 + 2 + \dots + (n-1))]$$

$$\left\{ \because \sum_{i=1}^{n-1} i = 1 + 2 + \dots + (n-1) = \frac{n(n-1)}{2} \right\}$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[n + h \left\{ \frac{n(n-1)}{2} \right\} \right]$$

Put,

$$h = \frac{5}{n}$$

Since,

$$h \rightarrow 0 \text{ and } h = \frac{5}{n} \Rightarrow n \rightarrow \infty$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} n \left[n + \frac{5}{n} \left\{ \frac{n(n-1)}{2} \right\} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} n \left[n + \frac{5(n-1)}{2} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left\{ 5 + \frac{25(n-1)}{2n} \right\}$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left\{ 5 + \frac{25}{2} \left(1 - \frac{1}{n} \right) \right\}$$

$$\Rightarrow I = 5 + \frac{25}{2} \left(1 - \frac{1}{\infty} \right)$$

$$\Rightarrow I = 5 + \frac{25}{2} (1 - 0)$$

$$\Rightarrow I = 5 + \frac{25}{2}$$

$$\Rightarrow I = \frac{10 + 25}{2}$$

$$\Rightarrow I = \frac{35}{2}$$

Hence, the value of $\int_0^5 (x+1) dx = \frac{35}{2}$

6. Question

Evaluate the following integrals as a limit of sums:

$$\int_1^3 (2x+3) dx$$

Answer

To find: $\int_1^3 (2x+3) dx$

Formula used:

$$\int_a^b f(x) dx = \lim_{h \rightarrow 0} h [f(a) + f(a+h) + f(a+2h) + \dots + f(a+(n-1)h)],$$

where,

$$h = \frac{b-a}{n}$$

Here, $a = 1$ and $b = 3$

Therefore,

$$h = \frac{3-1}{n}$$

$$\Rightarrow h = \frac{2}{n}$$

Let,

$$I = \int_1^3 (2x+3) dx$$

Here, $f(x) = 2x + 3$ and $a = 1$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[f(1) + f(1+h) + f(1+2h) + \dots + f(1+(n-1)h)]$$

Now, By putting $x = 1$ in $f(x)$ we get,

$$f(1) = 2(1) + 3 = 2 + 3 = 5$$

Similarly, $f(1+h)$

$$= 2(1+h) + 3$$

$$= 2 + 2h + 3$$

$$= 2h + 5$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[5 + 2h + 5 + 2(2h) + 5 + \dots + 2(n-1)h + 5]$$

In this series, 5 is getting added n times

$$\Rightarrow I = \lim_{h \rightarrow 0} h[5n + 2h(1 + 2 + \dots + (n-1))]$$

$$\left\{ \because \sum_{i=1}^{n-1} i = 1 + 2 + \dots + (n-1) = \frac{n(n-1)}{2} \right\}$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[5n + 2h \left\{ \frac{n(n-1)}{2} \right\} \right]$$

Put,

$$h = \frac{2}{n}$$

Since,

$$h \rightarrow 0 \text{ and } h = \frac{2}{n} \Rightarrow n \rightarrow \infty$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{2}{n} \left[5n + \frac{2(2) \left\{ \frac{n(n-1)}{2} \right\}}{n} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{2}{n} \left[5n + \frac{4(n-1)}{2} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{2}{n} [5n + 2(n-1)]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left\{ 10 + \frac{4(n-1)}{n} \right\}$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left\{ 10 + 4 \left(1 - \frac{1}{n} \right) \right\}$$

$$\Rightarrow I = 10 + 4 \left(1 - \frac{1}{\infty} \right)$$

$$\Rightarrow I = 10 + 4(1 - 0)$$

$$\Rightarrow I = 10 + 4$$

$$\Rightarrow I = 14$$

Hence, the value of $\int_1^3 (2x+3) dx = 14$

7. Question

Evaluate the following integrals as a limit of sums:

$$\int_3^5 (2-x) dx$$

Answer

To find: $\int_3^5 (2-x) dx$

Formula used:

$$\int_a^b f(x) dx = \lim_{h \rightarrow 0} h[f(a) + f(a+h) + f(a+2h) + \dots + f(a+(n-1)h)],$$

where,

$$h = \frac{b-a}{n}$$

Here, $a = 3$ and $b = 5$

Therefore,

$$h = \frac{5-3}{n}$$

$$\Rightarrow h = \frac{2}{n}$$

Let,

$$I = \int_3^5 (2-x) dx$$

Here, $f(x) = 2 - x$ and $a = 3$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[f(3) + f(3+h) + f(3+2h) + \dots + f(3+(n-1)h)]$$

Now, By putting $x = 3$ in $f(x)$ we get,

$$f(3) = 2 - 3 = -1$$

Similarly, $f(3+h)$

$$= 2 - (3+h)$$

$$= 2 - 3 - h$$

$$= -1 - h$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[-1 - 1 - h - 1 - 2h - \dots - 1 - (n-1)h]$$

In this series, -1 is getting added n times

$$\Rightarrow I = \lim_{h \rightarrow 0} h[-1 \times n - h - 2h - \dots - (n-1)h]$$

Now take $-h$ common in remaining series

$$\Rightarrow I = \lim_{h \rightarrow 0} h[-n - h(1 + 2 + \dots + (n-1))]$$

$$\left\{ \because \sum_{i=1}^{n-1} i = 1 + 2 + \dots + (n-1) = \frac{n(n-1)}{2} \right\}$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[-n - h \left\{ \frac{n(n-1)}{2} \right\} \right]$$

Put,

$$h = \frac{2}{n}$$

Since,

$$h \rightarrow 0 \text{ and } h = \frac{2}{n} \Rightarrow n \rightarrow \infty$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{2}{n} \left[-n - \frac{2}{n} \left\{ \frac{n(n-1)}{2} \right\} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{2}{n} [-n - (n-1)]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{2}{n} [-n - n + 1]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{2}{n} \{-2n + 1\}$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left\{ -4 + \frac{2}{n} \right\}$$

$$\Rightarrow I = -4 + \frac{2}{\infty}$$

$$\Rightarrow I = -4 + 0$$

$$\Rightarrow I = -4$$

Hence, the value of $\int_3^5 (2-x) dx = -4$

8. Question

Evaluate the following integrals as a limit of sums:

$$\int_0^2 (x^2 + 1) dx$$

Answer

$$\text{To find: } \int_0^2 (x^2 + 1) dx$$

Formula used:

$$\int_a^b f(x) dx = \lim_{h \rightarrow 0} h [f(a) + f(a+h) + f(a+2h) + \dots + f(a+(n-1)h)],$$

where,

$$h = \frac{b-a}{n}$$

Here, $a = 0$ and $b = 2$

Therefore,

$$h = \frac{2-0}{n}$$

$$\Rightarrow h = \frac{2}{n}$$

Let,

$$I = \int_0^2 (x^2 + 1) dx$$

Here, $f(x) = x^2 + 1$ and $a = 0$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[f(0) + f(0+h) + f(0+2h) + \dots + f(0+(n-1)h)]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[f(0) + f(h) + f(2h) + \dots + f((n-1)h)]$$

Now, By putting $x = 0$ in $f(x)$ we get,

$$f(0) = 0^2 + 1 = 0 + 1 = 1$$

Similarly, $f(h) = h^2 + 1$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[1 + h^2 + 1 + (2h)^2 + 1 + \dots + \{(n-1)h\}^2 + 1]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[1 + h^2 + 1 + h^2(2)^2 + 1 + \dots + h^2(n-1)^2 + 1]$$

In this series, 1 is getting added n times

$$\Rightarrow I = \lim_{h \rightarrow 0} h[1 \times n + h^2 + h^2(2)^2 + \dots + h^2(n-1)^2]$$

Now take h^2 common in remaining series

$$\Rightarrow I = \lim_{h \rightarrow 0} h[n + h^2\{1^2 + 2^2 + \dots + (n-1)^2\}]$$

$$\left\{ \because \sum_{i=1}^{n-1} i^2 = 1^2 + 2^2 + \dots + (n-1)^2 = \frac{n(n-1)(2n-1)}{6} \right\}$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[n + h^2 \left\{ \frac{n(n-1)(2n-1)}{6} \right\} \right]$$

Put,

$$h = \frac{2}{n}$$

Since,

$$h \rightarrow 0 \text{ and } h = \frac{2}{n} \Rightarrow n \rightarrow \infty$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{2}{n} \left[n + \left(\frac{2}{n} \right)^2 \left\{ \frac{n(n-1)(2n-1)}{6} \right\} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{2}{n} \left[n + \frac{4}{n^2} \left\{ \frac{n(n-1)(2n-1)}{6} \right\} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{2}{n} \left[n + \frac{2(n-1)(2n-1)}{3n} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[2 + \frac{4(n-1)(2n-1)}{3n \times n} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[2 + \frac{4}{3} \left(\frac{n-1}{n} \right) \left(\frac{2n-1}{n} \right) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[2 + \frac{4}{3} \left(1 - \frac{1}{n} \right) \left(2 - \frac{1}{n} \right) \right]$$

$$\Rightarrow I = 2 + \frac{4}{3} \left(1 - \frac{1}{\infty} \right) \left(2 - \frac{1}{\infty} \right)$$

$$\Rightarrow I = 2 + \frac{4}{3} (1 - 0)(2 - 0)$$

$$\Rightarrow I = 2 + \frac{4}{3} \times 1 \times 2$$

$$\Rightarrow I = 2 + \frac{8}{3}$$

$$\Rightarrow I = \frac{6+8}{3}$$

$$\Rightarrow I = \frac{14}{3}$$

Hence, the value of $\int_0^2 (x^2 + 1) dx = \frac{14}{3}$

9. Question

Evaluate the following integrals as a limit of sums:

$$\int_1^2 x^2 dx$$

Answer

To find: $\int_1^2 x^2 dx$

Formula used:

$$\int_a^b f(x) dx = \lim_{h \rightarrow 0} h [f(a) + f(a+h) + f(a+2h) + \dots + f(a+(n-1)h)],$$

where,

$$h = \frac{b-a}{n}$$

Here, $a = 1$ and $b = 2$

Therefore,

$$h = \frac{2-1}{n}$$

$$\Rightarrow h = \frac{1}{n}$$

Let,

$$I = \int_1^2 x^2 dx$$

Here, $f(x) = x^2$ and $a = 1$

$$\Rightarrow I = \lim_{h \rightarrow 0} h [f(1) + f(1+h) + f(1+2h) + \dots + f(1+(n-1)h)]$$

Now, by putting $x = 1$ in $f(x)$ we get,

$$f(1) = 1^2 = 1$$

$$f(1 + h)$$

$$= (1 + h)^2$$

$$= h^2 + 1^2 + 2(h)(1)$$

$$= h^2 + 1 + 2(h)$$

Similarly, $f(1 + 2h)$

$$= (1 + 2h)^2$$

$$= (2h)^2 + 1^2 + 2(2h)(1)$$

$$= (2h)^2 + 1 + 2(2h)$$

$$\{\because (x + y)^2 = x^2 + y^2 + 2xy\}$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[1 + h^2 + 1 + 2(h) + (2h)^2 + 1 + 2(2h) + \dots + \{(n - 1)h\}^2 + 1 + 2\{(n - 1)h\}]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[1 + h^2 + 1 + 2h + h^2(2)^2 + 1 + 2(2h) + \dots + h^2(n - 1)^2 + 1 + 2h(n - 1)]$$

In this series, 1 is getting added n times

$$\Rightarrow I = \lim_{h \rightarrow 0} h[1 \times n + h^2 + 2h + h^2(2)^2 + 2(2h) + \dots + h^2(n - 1)^2 + 2h(n - 1)]$$

Now take h^2 and $2h$ common in remaining series

$$\Rightarrow I = \lim_{h \rightarrow 0} h[n + h^2\{1^2 + 2^2 + \dots + (n - 1)^2\} + 2h\{1 + 2 + \dots + (n - 1)\}]$$

$$\left. \begin{aligned} & \because \sum_{i=1}^{n-1} i^2 = 1^2 + 2^2 + \dots + (n - 1)^2 = \frac{n(n - 1)(2n - 1)}{6}; \\ & \sum_{i=1}^{n-1} i = 1 + 2 + \dots + (n - 1) = \frac{n(n - 1)}{2} \end{aligned} \right\}$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[n + h^2 \left\{ \frac{n(n - 1)(2n - 1)}{6} \right\} + 2h \left\{ \frac{n(n - 1)}{2} \right\} \right]$$

Put,

$$h = \frac{1}{n}$$

Since,

$$h \rightarrow 0 \text{ and } h = \frac{1}{n} \Rightarrow n \rightarrow \infty$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{1}{n} \left[n + \left(\frac{1}{n} \right)^2 \left\{ \frac{n(n - 1)(2n - 1)}{6} \right\} + \frac{2(1)}{n} \left\{ \frac{n(n - 1)}{2} \right\} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{1}{n} \left[n + \frac{1}{n^2} \left\{ \frac{n(n - 1)(2n - 1)}{6} \right\} + \frac{2}{n} \left\{ \frac{n(n - 1)}{2} \right\} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{1}{n} \left[n + \frac{(n - 1)(2n - 1)}{6n} + (n - 1) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[1 + \frac{(n - 1)(2n - 1)}{6n \times n} + \frac{1}{n}(n - 1) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[1 + \frac{1}{6} \left(\frac{n-1}{n} \right) \left(\frac{2n-1}{n} \right) + \left(\frac{n-1}{n} \right) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[1 + \frac{1}{6} \left(1 - \frac{1}{n} \right) \left(2 - \frac{1}{n} \right) + \left(1 - \frac{1}{n} \right) \right]$$

$$\Rightarrow I = 1 + \frac{1}{6} \left(1 - \frac{1}{\infty} \right) \left(2 - \frac{1}{\infty} \right) + \left(1 - \frac{1}{\infty} \right)$$

$$\Rightarrow I = 1 + \frac{1}{6} (1 - 0)(2 - 0) + (1 - 0)$$

$$\Rightarrow I = 1 + \frac{1}{6} \times 1 \times 2 + 1$$

$$\Rightarrow I = 2 + \frac{1}{3}$$

$$\Rightarrow I = \frac{6+1}{3}$$

$$\Rightarrow I = \frac{7}{3}$$

Hence, the value of $\int_1^2 x^2 dx = \frac{7}{3}$

10. Question

Evaluate the following integrals as a limit of sums:

$$\int_2^3 (2x^2 + 1) dx$$

Answer

To find: $\int_2^3 (2x^2 + 1) dx$

Formula used:

$$\int_a^b f(x) dx = \lim_{h \rightarrow 0} h [f(a) + f(a+h) + f(a+2h) + \dots + f(a+(n-1)h)],$$

where,

$$h = \frac{b-a}{n}$$

Here, $a = 2$ and $b = 3$

Therefore,

$$h = \frac{3-2}{n}$$

$$\Rightarrow h = \frac{1}{n}$$

Let,

$$I = \int_2^3 (2x^2 + 1) dx$$

Here, $f(x) = 2x^2 + 1$ and $a = 2$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[f(2) + f(2+h) + f(2+2h) + \dots + f(2+(n-1)h)]$$

Now, by putting $x = 2$ in $f(x)$ we get,

$$f(2) = 2(2^2) + 1 = 2(4) + 1 = 8 + 1 = 9$$

$$f(1+h)$$

$$= 2(2+h)^2 + 1$$

$$= 2\{h^2 + 2^2 + 2(h)(2)\} + 1$$

$$= 2(h)^2 + 8 + 2(4h) + 1$$

$$= 2(h)^2 + 9 + 8(h)$$

Similarly, $f(2+2h)$

$$= 2(2+2h)^2 + 1$$

$$= 2\{2(2h)^2 + 2^2 + 2(2h)(2)\} + 1$$

$$= 2(2h)^2 + 8 + 8(2h) + 1$$

$$= 2(2h)^2 + 9 + 8(2h)$$

$$\{\because (x+y)^2 = x^2 + y^2 + 2xy\}$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[9 + 2(h)^2 + 9 + 8(h) + 2(2h)^2 + 9 + 8(2h) + \dots + 2\{(n-1)h\}^2 + 9 + 8\{(n-1)h\}]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[9 + 2h^2 + 9 + 8h + 2h^2(2)^2 + 9 + 8(2h) + \dots + 2h^2(n-1)^2 + 9 + 8h(n-1)]$$

In this series, 9 is getting added n times

$$\Rightarrow I = \lim_{h \rightarrow 0} h[9n + 2h^2\{1^2 + 2^2 + \dots + (n-1)^2\} + 8h\{1 + 2 + \dots + (n-1)\}]$$

Now take $2h^2$ and $4h$ common in remaining series

$$\Rightarrow I = \lim_{h \rightarrow 0} h[9n + 2h^2\{1^2 + 2^2 + \dots + (n-1)^2\} + 8h\{1 + 2 + \dots + (n-1)\}]$$

$$\left\{ \begin{array}{l} \because \sum_{i=1}^{n-1} i^2 = 1^2 + 2^2 + \dots + (n-1)^2 = \frac{n(n-1)(2n-1)}{6}; \\ \sum_{i=1}^{n-1} i = 1 + 2 + \dots + (n-1) = \frac{n(n-1)}{2} \end{array} \right\}$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[9n + 2h^2 \left\{ \frac{n(n-1)(2n-1)}{6} \right\} + 8h \left\{ \frac{n(n-1)}{2} \right\} \right]$$

Put,

$$h = \frac{1}{n}$$

Since,

$$h \rightarrow 0 \text{ and } h = \frac{1}{n} \Rightarrow n \rightarrow \infty$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{1}{n} \left[9n + 2 \left(\frac{1}{n} \right)^2 \left\{ \frac{n(n-1)(2n-1)}{6} \right\} + \frac{8(1)}{n} \left\{ \frac{n(n-1)}{2} \right\} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{1}{n} \left[9n + \frac{2}{n^2} \left\{ \frac{n(n-1)(2n-1)}{6} \right\} + \frac{8}{n} \left\{ \frac{n(n-1)}{2} \right\} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{1}{n} \left[9n + \frac{(n-1)(2n-1)}{3n} + 4(n-1) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[9 + \frac{(n-1)(2n-1)}{3n \times n} + \frac{4}{n}(n-1) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[9 + \frac{1}{3} \left(\frac{n-1}{n} \right) \left(\frac{2n-1}{n} \right) + 4 \left(\frac{n-1}{n} \right) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[9 + \frac{1}{3} \left(1 - \frac{1}{n} \right) \left(2 - \frac{1}{n} \right) + 4 \left(1 - \frac{1}{n} \right) \right]$$

$$\Rightarrow I = 9 + \frac{1}{3} \left(1 - \frac{1}{\infty} \right) \left(2 - \frac{1}{\infty} \right) + 4 \left(1 - \frac{1}{\infty} \right)$$

$$\Rightarrow I = 9 + \frac{1}{3} (1 - 0) (2 - 0) + 4(1 - 0)$$

$$\Rightarrow I = 9 + \frac{1}{3} \times 1 \times 2 + 4$$

$$\Rightarrow I = 13 + \frac{2}{3}$$

$$\Rightarrow I = \frac{39 + 2}{3}$$

$$\Rightarrow I = \frac{41}{3}$$

Hence, the value of $\int_2^3 (2x^2 + 1) dx = \frac{41}{3}$

11. Question

Evaluate the following integrals as a limit of sums:

$$\int_1^2 (x^2 - 1) dx$$

Answer

$$\text{To find: } \int_1^2 (x^2 - 1) dx$$

Formula used:

$$\int_a^b f(x) dx = \lim_{h \rightarrow 0} h [f(a) + f(a+h) + f(a+2h) + \dots + f(a+(n-1)h)],$$

where,

$$h = \frac{b-a}{n}$$

Here, $a = 1$ and $b = 2$

Therefore,

$$h = \frac{2-1}{n}$$

$$\Rightarrow h = \frac{1}{n}$$

Let,

$$I = \int_1^2 (x^2 - 1) dx$$

Here, $f(x) = x^2 - 1$ and $a = 1$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[f(1) + f(1+h) + f(1+2h) + \dots + f(1+(n-1)h)]$$

Now, by putting $x = 1$ in $f(x)$ we get,

$$f(1) = 1^2 - 1 = 1 - 1 = 0$$

$$f(1+h)$$

$$= (1+h)^2 - 1$$

$$= h^2 + 1^2 + 2(h)(1) - 1$$

$$= h^2 + 2(h)$$

Similarly, $f(1+2h)$

$$= (1+2h)^2 - 1$$

$$= (2h)^2 + 1^2 + 2(2h)(1) - 1$$

$$= (2h)^2 + 2(2h)$$

$$\{\because (x+y)^2 = x^2 + y^2 + 2xy\}$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[0 + h^2 + 2(h) + (2h)^2 + 2(2h) \dots \dots + \{(n-1)h\}^2 + 2\{(n-1)h\}]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[h^2 + 2h + h^2(2)^2 + 2(2h) \dots \dots + h^2(n-1)^2 + 2h(n-1)]$$

Now take h^2 and $2h$ common in remaining series

$$\Rightarrow I = \lim_{h \rightarrow 0} h[h^2\{1^2 + 2^2 + \dots + (n-1)^2\} + 2h\{1 + 2 + \dots + (n-1)\}]$$

$$\left. \begin{cases} \because \sum_{i=1}^{n-1} i^2 = 1^2 + 2^2 \dots \dots + (n-1)^2 = \frac{n(n-1)(2n-1)}{6}; \\ \sum_{i=1}^{n-1} i = 1 + 2 \dots \dots + (n-1) = \frac{n(n-1)}{2} \end{cases} \right\}$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[h^2 \left\{ \frac{n(n-1)(2n-1)}{6} \right\} + 2h \left\{ \frac{n(n-1)}{2} \right\} \right]$$

Put,

$$h = \frac{1}{n}$$

Since,

$$h \rightarrow 0 \text{ and } h = \frac{1}{n} \Rightarrow n \rightarrow \infty$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{1}{n} \left[\left(\frac{1}{n} \right)^2 \left\{ \frac{n(n-1)(2n-1)}{6} \right\} + \frac{2(1)}{n} \left\{ \frac{n(n-1)}{2} \right\} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{1}{n^2} \left[\frac{1}{6} \left\{ n(n-1)(2n-1) \right\} + \frac{2}{n} \left\{ n(n-1) \right\} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{1}{n} \left[\frac{(n-1)(2n-1)}{6n} + (n-1) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[\frac{(n-1)(2n-1)}{6n \times n} + \frac{1}{n}(n-1) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[\frac{1}{6} \left(\frac{n-1}{n} \right) \left(\frac{2n-1}{n} \right) + \left(\frac{n-1}{n} \right) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[\frac{1}{6} \left(1 - \frac{1}{n} \right) \left(2 - \frac{1}{n} \right) + \left(1 - \frac{1}{n} \right) \right]$$

$$\Rightarrow I = \frac{1}{6} \left(1 - \frac{1}{\infty} \right) \left(2 - \frac{1}{\infty} \right) + \left(1 - \frac{1}{\infty} \right)$$

$$\Rightarrow I = \frac{1}{6} (1 - 0) (2 - 0) + (1 - 0)$$

$$\Rightarrow I = \frac{1}{6} \times 1 \times 2 + 1$$

$$\Rightarrow I = 1 + \frac{1}{3}$$

$$\Rightarrow I = \frac{3+1}{3}$$

$$\Rightarrow I = \frac{4}{3}$$

Hence, the value of $\int_1^2 (x^2 - 1) dx = \frac{4}{3}$

12. Question

Evaluate the following integrals as a limit of sums:

$$\int_0^2 (x^2 + 4) dx$$

Answer

$$\text{To find: } \int_0^2 (x^2 + 4) dx$$

Formula used:

$$\int_a^b f(x) dx = \lim_{h \rightarrow 0} h [f(a) + f(a+h) + f(a+2h) + \dots + f(a+(n-1)h)],$$

where,

$$h = \frac{b-a}{n}$$

Here, $a = 0$ and $b = 2$

Therefore,

$$h = \frac{2-0}{n}$$

$$\Rightarrow h = \frac{2}{n}$$

Let,

$$I = \int_0^2 (x^2 + 4) dx$$

Here, $f(x) = x^2 + 4$ and $a = 0$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[f(0) + f(0+h) + f(0+2h) + \dots + f(0+(n-1)h)]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[f(0) + f(h) + f(2h) + \dots + f((n-1)h)]$$

Now, by putting $x = 0$ in $f(x)$ we get,

$$f(0) = 0^2 + 4 = 0 + 4 = 4$$

$$f(h)$$

$$= (h)^2 + 4$$

$$= h^2 + 4$$

Similarly, $f(2h)$

$$= (2h)^2 + 4$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[4 + h^2 + 4 + (2h)^2 + 4 + \dots + \{(n-1)h\}^2 + 4]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[4 + h^2 + 4 + h^2(2)^2 + 4 + \dots + h^2(n-1)^2 + 4]$$

In this series, 4 is getting added n times

$$\Rightarrow I = \lim_{h \rightarrow 0} h[4 \times n + h^2 + h^2(2)^2 + \dots + h^2(n-1)^2]$$

Now take h^2 common in remaining series

$$\Rightarrow I = \lim_{h \rightarrow 0} h[4n + h^2\{1^2 + 2^2 + \dots + (n-1)^2\}]$$

$$\left\{ \because \sum_{i=1}^{n-1} i^2 = 1^2 + 2^2 + \dots + (n-1)^2 = \frac{n(n-1)(2n-1)}{6} \right\}$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[4n + h^2 \left\{ \frac{n(n-1)(2n-1)}{6} \right\} \right]$$

Put,

$$h = \frac{2}{n}$$

Since,

$$h \rightarrow 0 \text{ and } h = \frac{2}{n} \Rightarrow n \rightarrow \infty$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{2}{n} \left[4n + \left(\frac{2}{n} \right)^2 \left\{ \frac{n(n-1)(2n-1)}{6} \right\} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{2}{n} \left[4n + \frac{4}{n^2} \left\{ \frac{n(n-1)(2n-1)}{6} \right\} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{2}{n} \left[4n + \frac{2(n-1)(2n-1)}{3n} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[8 + \frac{4(n-1)(2n-1)}{3n \times n} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[8 + \frac{4}{3} \left(\frac{n-1}{n} \right) \left(\frac{2n-1}{n} \right) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[8 + \frac{4}{3} \left(1 - \frac{1}{n} \right) \left(2 - \frac{1}{n} \right) \right]$$

$$\Rightarrow I = 8 + \frac{4}{3} \left(1 - \frac{1}{\infty} \right) \left(2 - \frac{1}{\infty} \right)$$

$$\Rightarrow I = 8 + \frac{4}{3} (1 - 0) (2 - 0)$$

$$\Rightarrow I = 8 + \frac{4}{3} \times 1 \times 2$$

$$\Rightarrow I = 8 + \frac{8}{3}$$

$$\Rightarrow I = \frac{24 + 8}{3}$$

$$\Rightarrow I = \frac{32}{3}$$

Hence, the value of $\int_0^2 (x^2 + 4) dx = \frac{32}{3}$

13. Question

Evaluate the following integrals as a limit of sums:

$$\int_1^4 (x^2 - x) dx$$

Answer

$$\text{To find: } \int_1^4 (x^2 - x) dx$$

Formula used:

$$\int_a^b f(x) dx = \lim_{h \rightarrow 0} h [f(a) + f(a+h) + f(a+2h) + \dots + f(a+(n-1)h)],$$

where,

$$h = \frac{b-a}{n}$$

Here, $a = 1$ and $b = 4$

Therefore,

$$h = \frac{4-1}{n}$$

$$\Rightarrow h = \frac{3}{n}$$

Let,

$$I = \int_1^4 (x^2 - x) dx$$

Here, $f(x) = x^2 - x$ and $a = 1$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[f(1) + f(1+h) + f(1+2h) + \dots + f(1+(n-1)h)]$$

Now, by putting $x = 1$ in $f(x)$ we get,

$$f(1) = 1^2 - 1 = 1 - 1 = 0$$

$$f(1+h)$$

$$= (1+h)^2 - (1+h)$$

$$= h^2 + 1^2 + 2(h)(1) - 1 - h$$

$$= h^2 + 2h - h$$

$$= h^2 + h$$

Similarly, $f(1+2h)$

$$= (1+2h)^2 - (1+2h)$$

$$= (2h)^2 + 1^2 + 2(2h)(1) - 1 - 2h$$

$$= (2h)^2 + 4h - 2h$$

$$= (2h)^2 + 2h$$

$$\{\because (x+y)^2 = x^2 + y^2 + 2xy\}$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[0 + h^2 + h + (2h)^2 + 2h + \dots + \{(n-1)h\}^2 + (n-1)h]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[h^2 + h + h^2(2)^2 + 2h + \dots + h^2(n-1)^2 + h(n-1)]$$

Now take h^2 and h common in remaining series

$$\Rightarrow I = \lim_{h \rightarrow 0} h[h^2\{1^2 + 2^2 + \dots + (n-1)^2\} + h\{1 + 2 + \dots + (n-1)\}]$$

$$\left. \begin{aligned} & \because \sum_{i=1}^{n-1} i^2 = 1^2 + 2^2 + \dots + (n-1)^2 = \frac{n(n-1)(2n-1)}{6}; \\ & \sum_{i=1}^{n-1} i = 1 + 2 + \dots + (n-1) = \frac{n(n-1)}{2} \end{aligned} \right\}$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[h^2 \left\{ \frac{n(n-1)(2n-1)}{6} \right\} + h \left\{ \frac{n(n-1)}{2} \right\} \right]$$

Put,

$$h = \frac{3}{n}$$

Since,

$$h \rightarrow 0 \text{ and } h = \frac{3}{n} \Rightarrow n \rightarrow \infty$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{3}{n} \left[\left(\frac{3}{n} \right)^2 \left\{ \frac{n(n-1)(2n-1)}{6} \right\} + \frac{3}{n} \left\{ \frac{n(n-1)}{2} \right\} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{3}{n} \left[\frac{9}{n^2} \left\{ \frac{n(n-1)(2n-1)}{6} \right\} + \frac{3}{n} \left\{ \frac{n(n-1)}{2} \right\} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{3}{n} \left[\frac{3(n-1)(2n-1)}{2n} + \frac{3}{2}(n-1) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[\frac{9(n-1)(2n-1)}{2n \times n} + \frac{9}{2n}(n-1) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[\frac{9}{2} \left(\frac{n-1}{n} \right) \left(\frac{2n-1}{n} \right) + \frac{9}{2} \left(\frac{n-1}{n} \right) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[\frac{9}{2} \left(1 - \frac{1}{n} \right) \left(2 - \frac{1}{n} \right) + \frac{9}{2} \left(1 - \frac{1}{n} \right) \right]$$

$$\Rightarrow I = \frac{9}{2} \left(1 - \frac{1}{\infty} \right) \left(2 - \frac{1}{\infty} \right) + \frac{9}{2} \left(1 - \frac{1}{\infty} \right)$$

$$\Rightarrow I = \frac{9}{2} (1 - 0) (2 - 0) + \frac{9}{2} (1 - 0)$$

$$\Rightarrow I = \frac{9}{2} \times 1 \times 2 + \frac{9}{2}$$

$$\Rightarrow I = 9 + \frac{9}{2}$$

$$\Rightarrow I = \frac{18 + 9}{2}$$

$$\Rightarrow I = \frac{27}{2}$$

Hence, the value of $\int_1^4 (x^2 - x) dx = \frac{27}{2}$

14. Question

Evaluate the following integrals as a limit of sums:

$$\int_0^1 (3x^2 + 5x) dx$$

Answer

$$\text{To find: } \int_0^1 (3x^2 + 5x) dx$$

Formula used:

$$\int_a^b f(x) dx = \lim_{h \rightarrow 0} h [f(a) + f(a+h) + f(a+2h) + \dots + f(a+(n-1)h)],$$

where,

$$h = \frac{b-a}{n}$$

Here, $a = 0$ and $b = 1$

Therefore,

$$h = \frac{1-0}{n}$$

$$\Rightarrow h = \frac{1}{n}$$

Let,

$$I = \int_0^1 (3x^2 + 5x) dx$$

Here, $f(x) = 3x^2 + 5x$ and $a = 0$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[f(0) + f(0+h) + f(0+2h) + \dots + f(0+(n-1)h)]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[f(0) + f(h) + f(2h) + \dots + f((n-1)h)]$$

Now, by putting $x = 0$ in $f(x)$ we get,

$$f(0) = 3(0)^2 + 5(0) = 0 + 0 = 0$$

$f(h)$

$$= 3(h)^2 + 5(h)$$

$$= 3h^2 + 5h$$

Similarly, $f(2h)$

$$= 3(2h)^2 + 5(2h)$$

$$= 3h^2(2)^2 + 5h(2)$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[0 + 3h^2 + 5h + 3h^2(2)^2 + 5h(2) + \dots + 3h^2(n-1)^2 + 5h(n-1)]$$

Now take $3h^2$ and $5h$ common in remaining series

$$\Rightarrow I = \lim_{h \rightarrow 0} h[3h^2\{1^2 + 2^2 + \dots + (n-1)^2\} + 5h\{1 + 2 + \dots + (n-1)\}]$$

$$\left. \begin{aligned} & \sum_{i=1}^{n-1} i^2 = 1^2 + 2^2 + \dots + (n-1)^2 = \frac{n(n-1)(2n-1)}{6}; \\ & \sum_{i=1}^{n-1} i = 1 + 2 + \dots + (n-1) = \frac{n(n-1)}{2} \end{aligned} \right\}$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[3h^2 \left\{ \frac{n(n-1)(2n-1)}{6} \right\} + 5h \left\{ \frac{n(n-1)}{2} \right\} \right]$$

Put,

$$h = \frac{1}{n}$$

Since,

$$h \rightarrow 0 \text{ and } h = \frac{1}{n} \Rightarrow n \rightarrow \infty$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{1}{n} \left[3 \times \left(\frac{1}{n} \right)^2 \left\{ \frac{n(n-1)(2n-1)}{6} \right\} + \frac{5(1)}{n} \left\{ \frac{n(n-1)}{2} \right\} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{1}{n} \left[\frac{3}{n^2} \left\{ \frac{n(n-1)(2n-1)}{6} \right\} + \frac{5}{n} \left\{ \frac{n(n-1)}{2} \right\} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{1}{n} \left[\frac{(n-1)(2n-1)}{2n} + \frac{5}{2}(n-1) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[\frac{(n-1)(2n-1)}{2n \times n} + \frac{5}{2n}(n-1) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[\frac{1}{2} \left(\frac{n-1}{n} \right) \left(\frac{2n-1}{n} \right) + \frac{5}{2} \left(\frac{n-1}{n} \right) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[\frac{1}{2} \left(1 - \frac{1}{n} \right) \left(2 - \frac{1}{n} \right) + \frac{5}{2} \left(1 - \frac{1}{n} \right) \right]$$

$$\Rightarrow I = \frac{1}{2} \left(1 - \frac{1}{\infty}\right) \left(2 - \frac{1}{\infty}\right) + \frac{5}{2} \left(1 - \frac{1}{\infty}\right)$$

$$\Rightarrow I = \frac{1}{2}(1-0)(2-0) + \frac{5}{2}(1-0)$$

$$\Rightarrow I = \frac{1}{2} \times 1 \times 2 + \frac{5}{2}$$

$$\Rightarrow I = 1 + \frac{5}{2}$$

$$\Rightarrow I = \frac{2+5}{2}$$

$$\Rightarrow I = \frac{7}{2}$$

Hence, the value of $\int_0^1 (3x^2 - 5x) dx = \frac{7}{2}$

15. Question

Evaluate the following integrals as a limit of sums:

$$\int_0^2 e^x dx$$

Answer

To find: $\int_0^2 e^x dx$

Formula used:

$$\int_a^b f(x) dx = \lim_{h \rightarrow 0} h[f(a) + f(a+h) + f(a+2h) + \dots + f(a+(n-1)h)],$$

where,

$$h = \frac{b-a}{n}$$

Here, $a = 0$ and $b = 2$

Therefore,

$$h = \frac{2-0}{n}$$

$$\Rightarrow h = \frac{2}{n}$$

Let,

$$I = \int_0^2 e^x dx$$

Here, $f(x) = e^x$ and $a = 0$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[f(0) + f(0+h) + f(0+2h) + \dots + f(0+(n-1)h)]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[f(0) + f(h) + f(2h) + \dots + f((n-1)h)]$$

Now, by putting $x = 0$ in $f(x)$ we get,

$$f(0) = e^0 = 1$$

$$f(h)$$

$$= (e)^h$$

$$= e^h$$

Similarly, $f(2h)$

$$= e^{2h}$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h [1 + e^h + e^{2h} + \dots + e^{(n-1)h}]$$

This is G.P. (Geometric Progression) of n terms whose first term (a) is 1

$$\text{and common ratio}(r) = \frac{e^h}{1} = e^h$$

Sum of n terms of a G.P. is given by,

$$S_n = \frac{a(r^n - 1)}{r - 1}, r > 1$$

Therefore,

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[\frac{1((e^h)^n - 1)}{e^h - 1} \right]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[\frac{e^{nh} - 1}{e^h - 1} \right]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[\frac{e^2 - 1}{e^h - 1} \right]$$

$$\left\{ \because h = \frac{2}{n} \Rightarrow nh = 2 \right\}$$

$$\Rightarrow I = (e^2 - 1) \lim_{h \rightarrow 0} \left[\frac{1}{\frac{e^h - 1}{h}} \right]$$

$$\Rightarrow I = (e^2 - 1) \lim_{h \rightarrow 0} \frac{1}{\frac{e^h - 1}{h}}$$

$$\left\{ \because \lim_{x \rightarrow 0} \frac{e^x - 1}{x} = 1 \right\}$$

$$\Rightarrow I = e^2 - 1$$

$$\text{Hence, the value of } \int_0^2 e^x dx = e^2 - 1$$

16. Question

Evaluate the following integrals as a limit of sums:

$$\int_a^b e^x dx$$

Answer

To find: $\int_a^b e^x dx$

Formula used:

$$\int_a^b f(x) dx = \lim_{h \rightarrow 0} h[f(a) + f(a+h) + f(a+2h) + \dots + f(a+(n-1)h)],$$

where,

$$h = \frac{b-a}{n}$$

$$\Rightarrow nh = b - a$$

Let,

$$I = \int_a^b e^x dx$$

Here, $f(x) = e^x$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[f(a) + f(a+h) + f(a+2h) + \dots + f(a+(n-1)h)]$$

Now, by putting $x = a$ in $f(x)$ we get,

$$f(a) = e^a$$

$$f(a+h)$$

$$= (e)^{a+h}$$

$$= e^{a+h}$$

Similarly, $f(a+2h)$

$$= e^{a+2h}$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[e^a + e^{a+h} + e^{a+2h} + \dots + e^{a+(n-1)h}]$$

This is G.P. (Geometric Progression) of n terms whose first term(a) is 1

$$\text{and common ratio}(r) = \frac{e^{a+h}}{e^a} = \frac{e^a \times e^h}{e^a} = e^h$$

Sum of n terms of a G.P. is given by,

$$S_n = \frac{a(r^n - 1)}{r - 1}, r > 1$$

Therefore,

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[\frac{e^a \{(e^h)^n - 1\}}{e^h - 1} \right]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[e^a \left\{ \frac{e^{nh} - 1}{e^h - 1} \right\} \right]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[\frac{e^a (e^{b-a} - 1)}{e^h - 1} \right]$$

$$\left\{ \because h = \frac{2}{n} \Rightarrow nh = 2 \right\}$$

$$\Rightarrow I = e^a (e^{b-a} - 1) \lim_{h \rightarrow 0} \left[\frac{1}{\frac{e^h - 1}{h}} \right]$$

$$\Rightarrow I = e^a \left(\frac{e^b}{e^a} - 1 \right) \frac{1}{\lim_{h \rightarrow 0} \frac{e^h - 1}{h}}$$

$$\left\{ \because \lim_{x \rightarrow 0} \frac{e^x - 1}{x} = 1 \right\}$$

$$\Rightarrow I = e^b - e^a$$

Hence, the value of $\int_a^b e^x dx = e^b - e^a$

17. Question

Evaluate the following integrals as a limit of sums:

$$\int_a^b \cos x dx$$

Answer

$$\text{To find: } \int_a^b \cos x dx$$

Formula used:

$$\int_a^b f(x) dx = \lim_{h \rightarrow 0} h[f(a) + f(a+h) + f(a+2h) + \dots + f(a+(n-1)h)],$$

where,

$$h = \frac{b-a}{n}$$

$$\Rightarrow nh = b - a$$

Let,

$$I = \int_a^b \cos x dx$$

Here, $f(x) = \cos x$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[f(a) + f(a+h) + f(a+2h) + \dots + f(a+(n-1)h)]$$

Now, by putting $x = a$ in $f(x)$ we get,

$$f(a) = \cos a$$

$$f(a+h)$$

$$= \cos(a+h)$$

$$\text{Similarly, } f(a+2h)$$

$$= \cos(a+2h)$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[\cos a + \cos(a+h) + \cos(a+2h) + \dots + \cos(a+(n-1)h)]$$

We know,

$$\cos A + \cos(A+B) + \cos(A+2B) + \dots + \cos\{A+(n-1)B\}$$

$$= \frac{\cos\left\{A + \frac{(n-1)B}{2}\right\} \sin \frac{nB}{2}}{\sin \frac{B}{2}}$$

Therefore,

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[\frac{\cos\left\{a + \frac{(n-1)h}{2}\right\} \sin \frac{nh}{2}}{\sin \frac{h}{2}} \right]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[\frac{\cos\left\{a + \frac{nh}{2} - \frac{h}{2}\right\} \sin \frac{nh}{2}}{\sin \frac{h}{2}} \right]$$

$$\{\because nh = b - a\}$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[\frac{\cos\left\{a + \frac{b-a}{2} - \frac{h}{2}\right\} \sin \frac{b-a}{2}}{\sin \frac{h}{2}} \right]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[\frac{\cos\left\{a + \frac{b}{2} - \frac{a}{2} - \frac{h}{2}\right\} \sin \frac{b-a}{2}}{\sin \frac{h}{2}} \right]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[\frac{\cos\left\{\frac{a}{2} + \frac{b}{2} - \frac{h}{2}\right\} \sin \frac{b-a}{2}}{\sin \frac{h}{2}} \right]$$

$$\Rightarrow I = \sin \frac{b-a}{2} \times \lim_{h \rightarrow 0} \left[\frac{2 \times \frac{h}{2} \times \cos\left\{\frac{a}{2} + \frac{b}{2} - \frac{h}{2}\right\}}{\sin \frac{h}{2}} \right]$$

$$\Rightarrow I = 2 \sin \frac{b-a}{2} \times \lim_{h \rightarrow 0} \left[\frac{\cos\left\{\frac{a+b}{2} - \frac{h}{2}\right\}}{\frac{\sin \frac{h}{2}}{\frac{h}{2}}} \right]$$

$$\Rightarrow I = 2 \sin \frac{b-a}{2} \times \frac{\lim_{h \rightarrow 0} \left[\cos\left\{\frac{a+b}{2} - \frac{h}{2}\right\} \right]}{\lim_{h \rightarrow 0} \left[\frac{\sin \frac{h}{2}}{\frac{h}{2}} \right]}$$

$$\text{As, } h \rightarrow 0 \Rightarrow \frac{h}{2} \rightarrow 0$$

$$\Rightarrow I = 2 \sin \frac{b-a}{2} \times \frac{\lim_{\frac{h}{2} \rightarrow 0} \left[\cos\left\{\frac{a+b}{2} - \frac{h}{2}\right\} \right]}{\lim_{\frac{h}{2} \rightarrow 0} \left[\frac{\sin \frac{h}{2}}{\frac{h}{2}} \right]}$$

$$\left\{ \because \lim_{x \rightarrow 0} \left[\frac{\sin x}{x} \right] = 1 \right\}$$

$$\Rightarrow I = 2 \sin \frac{b-a}{2} \times \frac{\cos \left\{ \frac{a+b}{2} - 0 \right\}}{1}$$

$$\Rightarrow I = 2 \cos \frac{a+b}{2} \sin \frac{b-a}{2}$$

$\{ \because 2 \cos A \sin B = \sin(A+B) - \sin(A-B) \}$

$$\Rightarrow I = \sin \left\{ \frac{a+b}{2} + \frac{b-a}{2} \right\} - \sin \left\{ \frac{a+b}{2} - \frac{b-a}{2} \right\}$$

$$\Rightarrow I = \sin \left\{ \frac{a+b+b-a}{2} \right\} - \sin \left\{ \frac{a+b-b+a}{2} \right\}$$

$$\Rightarrow I = \sin \left\{ \frac{2b}{2} \right\} - \sin \left\{ \frac{2a}{2} \right\}$$

$$\Rightarrow I = \sin b - \sin a$$

Hence, the value of $\int_a^b \cos x \, dx = \sin b - \sin a$

18. Question

Evaluate the following integrals as a limit of sums:

$$\int_0^{\pi/2} \sin x \, dx$$

Answer

$$\text{To find: } \int_0^{\frac{\pi}{2}} \sin x \, dx$$

Formula used:

$$\int_a^b f(x) \, dx = \lim_{h \rightarrow 0} h[f(a) + f(a+h) + f(a+2h) + \dots + f(a+(n-1)h)],$$

where,

$$h = \frac{b-a}{n}$$

$$\text{Here, } a = 0 \text{ and } b = \frac{\pi}{2}$$

$$\Rightarrow h = \frac{\frac{\pi}{2} - 0}{n}$$

$$\Rightarrow h = \frac{\pi}{2n}$$

$$\Rightarrow nh = \frac{\pi}{2}$$

Let,

$$I = \int_0^{\frac{\pi}{2}} \sin x \, dx$$

Here, $f(x) = \sin x$ and $a = 0$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[f(0) + f(0+h) + f(0+2h) + \dots + f(0+(n-1)h)]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h [f(0) + f(h) + f(2h) + \dots + f((n-1)h)]$$

Now, by putting $x = 0$ in $f(x)$ we get,

$$f(0) = \sin 0$$

$$f(h)$$

$$= \sin h$$

$$\text{Similarly, } f(2h)$$

$$= \sin 2h$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h [\sin 0 + \sin h + \sin 2h + \dots + \sin(n-1)h]$$

We know,

$$\begin{aligned} & \sin A + \sin(A+B) + \sin(A+2B) + \dots + \sin\{A+(n-1)B\} \\ &= \frac{\sin\left\{A + \frac{(n-1)B}{2}\right\} \sin \frac{nB}{2}}{\sin \frac{B}{2}} \end{aligned}$$

Here $A = 0$ and $B = h$

Therefore,

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[\frac{\sin\left\{0 + \frac{(n-1)h}{2}\right\} \sin \frac{nh}{2}}{\sin \frac{h}{2}} \right]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[\frac{\sin\left\{\frac{nh}{2} - \frac{h}{2}\right\} \sin \frac{nh}{2}}{\sin \frac{h}{2}} \right]$$

$$\left\{ \because nh = \frac{\pi}{2} \right\}$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[\frac{\sin\left\{\frac{\pi}{2} - \frac{h}{2}\right\} \sin \frac{\pi}{2}}{\sin \frac{h}{2}} \right]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[\frac{\sin\left\{\frac{\pi}{4} - \frac{h}{2}\right\} \sin \frac{\pi}{4}}{\sin \frac{h}{2}} \right]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[\frac{\sin\left\{\frac{\pi}{4} - \frac{h}{2}\right\} \sin \frac{\pi}{4}}{\sin \frac{h}{2}} \right]$$

$$\Rightarrow I = \sin \frac{\pi}{4} \times \lim_{h \rightarrow 0} \left[\frac{2 \times \frac{h}{2} \times \sin\left\{\frac{\pi}{4} - \frac{h}{2}\right\}}{\sin \frac{h}{2}} \right]$$

$$\left\{ \because \sin \frac{\pi}{4} = \sin 45^\circ = \frac{1}{\sqrt{2}} \right\}$$

$$\Rightarrow I = 2 \times \frac{1}{\sqrt{2}} \times \lim_{h \rightarrow 0} \left[\frac{\sin \left\{ \frac{\pi}{4} - \frac{h}{2} \right\}}{\frac{\sin \frac{h}{2}}{\frac{h}{2}}} \right]$$

As, $h \rightarrow 0 \Rightarrow \frac{h}{2} \rightarrow 0$

$$\Rightarrow I = \sqrt{2} \times \frac{\lim_{h \rightarrow 0} \left[\sin \left\{ \frac{\pi}{4} - \frac{h}{2} \right\} \right]}{\lim_{h \rightarrow 0} \left[\frac{\sin \frac{h}{2}}{\frac{h}{2}} \right]}$$

$$\left\{ \because \lim_{x \rightarrow 0} \left[\frac{\sin x}{x} \right] = 1 \right\}$$

$$\Rightarrow I = \sqrt{2} \times \frac{\sin \left\{ \frac{\pi}{4} - 0 \right\}}{1}$$

$$\Rightarrow I = \sqrt{2} \times \sin \frac{\pi}{4}$$

$$\left\{ \because \sin \frac{\pi}{4} = \sin 45^\circ = \frac{1}{\sqrt{2}} \right\}$$

$$\Rightarrow I = \sqrt{2} \times \frac{1}{\sqrt{2}}$$

$$\Rightarrow I = 1$$

Hence, the value of $\int_0^{\frac{\pi}{2}} \sin x \, dx = 1$

19. Question

Evaluate the following integrals as a limit of sums:

$$\int_0^{\pi/2} \cos x \, dx$$

Answer

$$\text{To find: } \int_0^{\frac{\pi}{2}} \cos x \, dx$$

Formula used:

$$\int_a^b f(x) \, dx = \lim_{h \rightarrow 0} h[f(a) + f(a+h) + f(a+2h) + \dots + f(a+(n-1)h)],$$

where,

$$h = \frac{b-a}{n}$$

$$\text{Here, } a = 0 \text{ and } b = \frac{\pi}{2}$$

$$\Rightarrow h = \frac{\frac{\pi}{2} - 0}{n}$$

$$\Rightarrow h = \frac{\pi}{2n}$$

$$\Rightarrow nh = \frac{\pi}{2}$$

Let,

$$I = \int_0^{\frac{\pi}{2}} \cos x \, dx$$

Here, $f(x) = \cos x$ and $a = 0$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[f(0) + f(0+h) + f(0+2h) + \dots + f(0+(n-1)h)]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[f(0) + f(h) + f(2h) + \dots + f((n-1)h)]$$

Now, by putting $x = 0$ in $f(x)$ we get,

$$f(0) = \cos 0$$

$$f(h)$$

$$= \cos h$$

Similarly, $f(2h)$

$$= \cos 2h$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[\cos 0 + \cosh + \cos 2h + \dots + \cos(n-1)h]$$

We know,

$$\begin{aligned} & \cos A + \cos(A+B) + \cos(A+2B) + \dots + \cos(A+(n-1)B) \\ &= \frac{\cos\left\{A + \frac{(n-1)B}{2}\right\} \sin \frac{nB}{2}}{\sin \frac{B}{2}} \end{aligned}$$

Here $A = 0$ and $B = h$

Therefore,

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[\frac{\cos\left\{0 + \frac{(n-1)h}{2}\right\} \sin \frac{nh}{2}}{\sin \frac{h}{2}} \right]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[\frac{\cos\left\{\frac{nh}{2} - \frac{h}{2}\right\} \sin \frac{nh}{2}}{\sin \frac{h}{2}} \right]$$

$$\left\{ \because nh = \frac{\pi}{2} \right\}$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[\frac{\cos\left\{\frac{\pi}{2} - \frac{h}{2}\right\} \sin \frac{\pi}{2}}{\sin \frac{h}{2}} \right]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[\frac{\cos\left(\frac{\pi}{4} - \frac{h}{2}\right) \sin\frac{\pi}{4}}{\sin\frac{h}{2}} \right]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[\frac{\cos\left(\frac{\pi}{4} - \frac{h}{2}\right) \sin\frac{\pi}{4}}{\sin\frac{h}{2}} \right]$$

$$\Rightarrow I = \sin\frac{\pi}{4} \times \lim_{h \rightarrow 0} \left[\frac{2 \times \frac{h}{2} \times \cos\left(\frac{\pi}{4} - \frac{h}{2}\right)}{\sin\frac{h}{2}} \right]$$

$$\left\{ \because \sin\frac{\pi}{4} = \sin 45^\circ = \frac{1}{\sqrt{2}} \right\}$$

$$\Rightarrow I = 2 \times \frac{1}{\sqrt{2}} \times \lim_{h \rightarrow 0} \left[\frac{\cos\left(\frac{\pi}{4} - \frac{h}{2}\right)}{\frac{\sin\frac{h}{2}}{\frac{h}{2}}} \right]$$

$$\text{As, } h \rightarrow 0 \Rightarrow \frac{h}{2} \rightarrow 0$$

$$\Rightarrow I = \sqrt{2} \times \frac{\lim_{\frac{h}{2} \rightarrow 0} \left[\cos\left(\frac{\pi}{4} - \frac{h}{2}\right) \right]}{\lim_{\frac{h}{2} \rightarrow 0} \left[\frac{\sin\frac{h}{2}}{\frac{h}{2}} \right]}$$

$$\left\{ \because \lim_{x \rightarrow 0} \left[\frac{\sin x}{x} \right] = 1 \right\}$$

$$\Rightarrow I = \sqrt{2} \times \frac{\cos\left(\frac{\pi}{4} - 0\right)}{1}$$

$$\Rightarrow I = \sqrt{2} \times \cos\frac{\pi}{4}$$

$$\left\{ \because \cos\frac{\pi}{4} = \cos 45^\circ = \frac{1}{\sqrt{2}} \right\}$$

$$\Rightarrow I = \sqrt{2} \times \frac{1}{\sqrt{2}}$$

$$\Rightarrow I = 1$$

Hence, the value of $\int_0^{\frac{\pi}{2}} \cos x \, dx = 1$

20. Question

Evaluate the following integrals as a limit of sums:

$$\int_1^4 (3x^2 + 2x) \, dx$$

Answer

To find: $\int_1^4 (3x^2 + 2x) dx$

Formula used:

$$\int_a^b f(x) dx = \lim_{h \rightarrow 0} h[f(a) + f(a+h) + f(a+2h) + \dots + f(a+(n-1)h)],$$

where,

$$h = \frac{b-a}{n}$$

Here, $a = 1$ and $b = 4$

Therefore,

$$h = \frac{4-1}{n}$$

$$\Rightarrow h = \frac{3}{n}$$

Let,

$$I = \int_1^4 (3x^2 + 2x) dx$$

Here, $f(x) = 3x^2 + 2x$ and $a = 1$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[f(1) + f(1+h) + f(1+2h) + \dots + f(1+(n-1)h)]$$

Now, by putting $x = 1$ in $f(x)$ we get,

$$f(1) = 3(1)^2 + 2(1) = 3 + 2 = 5$$

$$f(1+h)$$

$$= 3(1+h)^2 + 2(1+h)$$

$$= 3\{h^2 + 1^2 + 2(h)(1)\} + 2 + 2h$$

$$= 3h^2 + 3 + 6h + 2 + 2h$$

$$= 3h^2 + 8h + 5$$

Similarly, $f(1+2h)$

$$= 3(1+2h)^2 + 2(1+2h)$$

$$= 3\{(2h)^2 + 1^2 + 2(2h)(1)\} + 2 + 4h$$

$$= 3(2h)^2 + 3 + 6(2h) + 2 + 2(2h)$$

$$= 3(2h)^2 + 8(2h) + 5$$

$$\{\because (x+y)^2 = x^2 + y^2 + 2xy\}$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[5 + 3h^2 + 8h + 5 + 3(2h)^2 + 8(2h) + 5 + \dots + 3\{(n-1)h\}^2 + 8(n-1)h + 5]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[5 + 3h^2 + 8h + 5 + 3h^2(2)^2 + 8h(2) + 5 + \dots + 3h^2(n-1)^2 + 8h(n-1) + 5]$$

Since 5 is repeating n times in series

$$\Rightarrow I = \lim_{h \rightarrow 0} h[5n + 3h^2 + 8h + 3h^2(2)^2 + 8h(2) + \dots + 3h^2(n-1)^2 + 8h(n-1)]$$

Now take $3h^2$ and $8h$ common in remaining series

$$\Rightarrow I = \lim_{h \rightarrow 0} h [5n + 3h^2 \{1^2 + 2^2 + \dots + (n-1)^2\} + 8h \{1 + 2 + \dots + (n-1)\}]$$

$$\left\{ \begin{array}{l} \because \sum_{i=1}^{n-1} i^2 = 1^2 + 2^2 + \dots + (n-1)^2 = \frac{n(n-1)(2n-1)}{6}; \\ \sum_{i=1}^{n-1} i = 1 + 2 + \dots + (n-1) = \frac{n(n-1)}{2} \end{array} \right\}$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[5n + 3h^2 \left\{ \frac{n(n-1)(2n-1)}{6} \right\} + 8h \left\{ \frac{n(n-1)}{2} \right\} \right]$$

Put,

$$h = \frac{3}{n}$$

Since,

$$h \rightarrow 0 \text{ and } h = \frac{3}{n} \Rightarrow n \rightarrow \infty$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{3}{n} \left[5n + 3 \left(\frac{3}{n} \right)^2 \left\{ \frac{n(n-1)(2n-1)}{6} \right\} + \frac{3(8)}{n} \left\{ \frac{n(n-1)}{2} \right\} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{3}{n} \left[5n + \frac{27}{n^2} \left\{ \frac{n(n-1)(2n-1)}{6} \right\} + \frac{24}{n} \left\{ \frac{n(n-1)}{2} \right\} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{3}{n} \left[5n + \frac{9(n-1)(2n-1)}{2n} + 12(n-1) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[15 + \frac{27(n-1)(2n-1)}{2n \times n} + \frac{36}{n}(n-1) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[15 + \frac{27}{2} \left(\frac{n-1}{n} \right) \left(\frac{2n-1}{n} \right) + 36 \left(\frac{n-1}{n} \right) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[15 + \frac{27}{2} \left(1 - \frac{1}{n} \right) \left(2 - \frac{1}{n} \right) + 36 \left(1 - \frac{1}{n} \right) \right]$$

$$\Rightarrow I = 15 + \frac{27}{2} \left(1 - \frac{1}{\infty} \right) \left(2 - \frac{1}{\infty} \right) + 36 \left(1 - \frac{1}{\infty} \right)$$

$$\Rightarrow I = 15 + \frac{27}{2} (1 - 0)(2 - 0) + 36(1 - 0)$$

$$\Rightarrow I = 15 + \frac{27}{2} \times 1 \times 2 + 36$$

$$\Rightarrow I = 15 + 27 + 36$$

$$\Rightarrow I = 78$$

Hence, the value of $\int_1^4 (3x^2 + 2x) dx = 78$

21. Question

Evaluate the following integrals as a limit of sums:

$$\int_0^2 (3x^2 - 2) dx$$

Answer

To find: $\int_0^2 (3x^2 - 2) dx$

Formula used:

$$\int_a^b f(x) dx = \lim_{h \rightarrow 0} h [f(a) + f(a+h) + f(a+2h) + \dots + f(a+(n-1)h)],$$

where,

$$h = \frac{b-a}{n}$$

Here, $a = 0$ and $b = 2$

Therefore,

$$h = \frac{2-0}{n}$$

$$\Rightarrow h = \frac{2}{n}$$

Let,

$$I = \int_0^2 (3x^2 - 2) dx$$

Here, $f(x) = 3x^2 - 2$ and $a = 0$

$$\Rightarrow I = \lim_{h \rightarrow 0} h [f(0) + f(0+h) + f(0+2h) + \dots + f(0+(n-1)h)]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h [f(0) + f(h) + f(2h) + \dots + f((n-1)h)]$$

Now, by putting $x = 0$ in $f(x)$ we get,

$$f(0) = 3(0)^2 - 2 = 0 - 2 = -2$$

$f(h)$

$$= 3(h)^2 - 2$$

Similarly, $f(2h)$

$$= 3(2h)^2 - 2$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h [-2 + 3h^2 - 2 + 3(2h)^2 - 2 + \dots + 3\{(n-1)h\}^2 - 2]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h [-2 + 3h^2 - 2 + 3h^2(2)^2 - 2 + \dots + 3h^2(n-1)^2 - 2]$$

Since -2 is repeating n times in series

$$\Rightarrow I = \lim_{h \rightarrow 0} h [-2n + 3h^2 + 3h^2(2)^2 + \dots + 3h^2(n-1)^2]$$

Now take $3h^2$ common in remaining series

$$\Rightarrow I = \lim_{h \rightarrow 0} h [-2n + 3h^2 \{1^2 + 2^2 + \dots + (n-1)^2\}]$$

$$\left\{ \because \sum_{i=1}^{n-1} i^2 = 1^2 + 2^2 + \dots + (n-1)^2 = \frac{n(n-1)(2n-1)}{6} \right\}$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[-2n + 3h^2 \left\{ \frac{n(n-1)(2n-1)}{6} \right\} \right]$$

Put,

$$h = \frac{2}{n}$$

Since,

$$h \rightarrow 0 \text{ and } h = \frac{2}{n} \Rightarrow n \rightarrow \infty$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{2}{n} \left[-2n + 3 \left(\frac{2}{n} \right)^2 \left\{ \frac{n(n-1)(2n-1)}{6} \right\} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{2}{n} \left[-2n + \frac{12}{n^2} \left\{ \frac{n(n-1)(2n-1)}{6} \right\} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{2}{n} \left[-2n + \frac{2(n-1)(2n-1)}{n} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[-4 + \frac{4(n-1)(2n-1)}{n \times n} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[-4 + 4 \left(\frac{n-1}{n} \right) \left(\frac{2n-1}{n} \right) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[-4 + 4 \left(1 - \frac{1}{n} \right) \left(2 - \frac{1}{n} \right) \right]$$

$$\Rightarrow I = -4 + 4 \left(1 - \frac{1}{\infty} \right) \left(2 - \frac{1}{\infty} \right)$$

$$\Rightarrow I = -4 + 4(1-0)(2-0)$$

$$\Rightarrow I = -4 + 4 \times 1 \times 2$$

$$\Rightarrow I = -4 + 8$$

$$\Rightarrow I = 4$$

Hence, the value of $\int_0^2 (3x^2 - 2) dx = 4$

22. Question

Evaluate the following integrals as a limit of sums:

$$\int_0^2 (x^2 + 2) dx$$

Answer

$$\text{To find: } \int_0^2 (x^2 + 2) dx$$

Formula used:

$$\int_a^b f(x) dx = \lim_{h \rightarrow 0} h [f(a) + f(a+h) + f(a+2h) + \dots + f(a+(n-1)h)],$$

where,

$$h = \frac{b-a}{n}$$

Here, $a = 0$ and $b = 2$

Therefore,

$$h = \frac{2 - 0}{n}$$

$$\Rightarrow h = \frac{2}{n}$$

Let,

$$I = \int_0^2 (x^2 + 2) dx$$

Here, $f(x) = x^2 + 2$ and $a = 0$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[f(0) + f(0+h) + f(0+2h) + \dots + f(0+(n-1)h)]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[f(0) + f(h) + f(2h) + \dots + f((n-1)h)]$$

Now, by putting $x = 0$ in $f(x)$ we get,

$$f(0) = (0)^2 + 2 = 0 + 2 = 2$$

$$f(h)$$

$$= (h)^2 + 2$$

Similarly, $f(2h)$

$$= (2h)^2 + 2$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[2 + h^2 + 2 + (2h)^2 + 2 + \dots + \{(n-1)h\}^2 + 2]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[2 + h^2 + 2 + h^2(2)^2 + 2 + \dots + h^2(n-1)^2 + 2]$$

Since 2 is repeating n times in series

$$\Rightarrow I = \lim_{h \rightarrow 0} h[2n + h^2(2)^2 + \dots + h^2(n-1)^2]$$

Now take h^2 common in remaining series

$$\Rightarrow I = \lim_{h \rightarrow 0} h[2n + h^2\{1^2 + 2^2 + \dots + (n-1)^2\}]$$

$$\left\{ \because \sum_{i=1}^{n-1} i^2 = 1^2 + 2^2 + \dots + (n-1)^2 = \frac{n(n-1)(2n-1)}{6} \right\}$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[2n + h^2 \left\{ \frac{n(n-1)(2n-1)}{6} \right\} \right]$$

Put,

$$h = \frac{2}{n}$$

Since,

$$h \rightarrow 0 \text{ and } h = \frac{2}{n} \Rightarrow n \rightarrow \infty$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{2}{n} \left[2n + \left(\frac{2}{n} \right)^2 \left\{ \frac{n(n-1)(2n-1)}{6} \right\} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{2}{n} \left[2n + \frac{4}{n^2} \left\{ \frac{n(n-1)(2n-1)}{6} \right\} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{2}{n} \left[2n + \frac{2(n-1)(2n-1)}{3n} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[4 + \frac{4(n-1)(2n-1)}{3n \times n} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[4 + \frac{4}{3} \left(\frac{n-1}{n} \right) \left(\frac{2n-1}{n} \right) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[4 + \frac{4}{3} \left(1 - \frac{1}{n} \right) \left(2 - \frac{1}{n} \right) \right]$$

$$\Rightarrow I = 4 + \frac{4}{3} \left(1 - \frac{1}{\infty} \right) \left(2 - \frac{1}{\infty} \right)$$

$$\Rightarrow I = 4 + \frac{4}{3} (1 - 0) (2 - 0)$$

$$\Rightarrow I = 4 + \frac{4}{3} \times 1 \times 2$$

$$\Rightarrow I = 4 + \frac{8}{3}$$

$$\Rightarrow I = \frac{12 + 8}{3}$$

$$\Rightarrow I = \frac{20}{3}$$

Hence, the value of $\int_0^2 (x^2 + 2) dx = \frac{20}{3}$

23. Question

Evaluate the following integrals as a limit of sums:

$$\int_0^4 (x + e^{2x}) dx$$

Answer

$$\text{To find: } \int_0^4 (x + e^{2x}) dx$$

Formula used:

$$\int_a^b f(x) dx = \lim_{h \rightarrow 0} h [f(a) + f(a+h) + f(a+2h) + \dots + f(a+(n-1)h)],$$

where,

$$h = \frac{b-a}{n}$$

Here, $a = 0$ and $b = 4$

Therefore,

$$h = \frac{4-0}{n}$$

$$\Rightarrow h = \frac{4}{n}$$

Let,

$$I = \int_0^4 (x + e^{2x}) dx$$

Here, $f(x) = x + e^{2x}$ and $a = 0$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[f(0) + f(0+h) + f(0+2h) + \dots + f(0+(n-1)h)]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[f(0) + f(h) + f(2h) + \dots + f((n-1)h)]$$

Now, by putting $x = 0$ in $f(x)$ we get,

$$f(0) = 0 + e^{2(0)} = 0 + e^0 = 0 + 1 = 1$$

$f(h)$

$$= h + (e)^{2h}$$

$$= h + e^{2h}$$

Similarly, $f(2h)$

$$= 2h + (e)^{2(2h)}$$

$$= 2h + e^{4h}$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[1 + h + e^{2h} + 2h + e^{4h} + \dots + (n-1)h + e^{2(n-1)h}]$$

Take h common in some of the terms of series

$$\Rightarrow I = \lim_{h \rightarrow 0} h[1 + e^{2h} + e^{4h} + \dots + e^{2(n-1)h} + h\{1 + 2 + \dots + (n-1)\}]$$

This is G.P. (Geometric Progression) of n terms whose first term(a) is 1

$$\text{and common ratio}(r) = \frac{e^{2h}}{1} = e^{2h}$$

Sum of n terms of a G.P. is given by,

$$S_n = \frac{a(r^n - 1)}{r - 1}, r > 1$$

and

$$\sum_{i=1}^{n-1} i = 1 + 2 + \dots + (n-1) = \frac{n(n-1)}{2}$$

Therefore,

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[\frac{1\{(e^{2h})^n - 1\}}{e^{2h} - 1} + h \left\{ \frac{n(n-1)}{2} \right\} \right]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[\frac{e^{2nh} - 1}{e^{2h} - 1} + h \left\{ \frac{n(n-1)}{2} \right\} \right]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} \left[\frac{2h}{2} \times \frac{e^8 - 1}{e^{2h} - 1} + h^2 \left\{ \frac{n(n-1)}{2} \right\} \right]$$

$$\left\{ \because h = \frac{4}{n} \Rightarrow nh = 4 \right\}$$

$$\Rightarrow I = \frac{e^8 - 1}{2} \lim_{h \rightarrow 0} \left[\frac{1}{\frac{e^{2h} - 1}{2h}} \right] + \lim_{h \rightarrow 0} \left[\left(\frac{4}{n} \right)^2 \times \left\{ \frac{n(n-1)}{2} \right\} \right]$$

As, $h \rightarrow 0 \Rightarrow 2h \rightarrow 0$ and $\frac{4}{n} \rightarrow 0 \Rightarrow n \rightarrow \infty$

$$\Rightarrow I = \frac{e^8 - 1}{2} \times \frac{1}{\lim_{2h \rightarrow 0} \frac{e^{2h} - 1}{2h}} + \lim_{n \rightarrow \infty} \left[\frac{16}{n^2} \times \left\{ \frac{n(n-1)}{2} \right\} \right]$$

$$\left\{ \because \lim_{x \rightarrow 0} \frac{e^x - 1}{x} = 1 \right\}$$

$$\Rightarrow I = \frac{e^8 - 1}{2} \times \frac{1}{1} + \lim_{n \rightarrow \infty} \left[8 \times \left(1 - \frac{1}{n} \right) \right]$$

$$\Rightarrow I = \frac{e^8 - 1}{2} + 8 \times \left(1 - \frac{1}{\infty} \right)$$

$$\Rightarrow I = \frac{e^8 - 1}{2} + 8 \times (1 - 0)$$

$$\Rightarrow I = \frac{e^8 - 1 + 16}{2}$$

$$\Rightarrow I = \frac{e^8 + 15}{2}$$

Hence, the value of $\int_0^4 (x + e^x) dx = \frac{e^8 + 15}{2}$

24. Question

Evaluate the following integrals as a limit of sums:

$$\int_0^2 (x^2 + x) dx$$

Answer

To find: $\int_0^2 (x^2 + x) dx$

Formula used:

$$\int_a^b f(x) dx = \lim_{h \rightarrow 0} h[f(a) + f(a+h) + f(a+2h) + \dots + f(a+(n-1)h)],$$

where,

$$h = \frac{b-a}{n}$$

Here, $a = 0$ and $b = 2$

Therefore,

$$h = \frac{2-0}{n}$$

$$\Rightarrow h = \frac{2}{n}$$

Let,

$$I = \int_0^2 (x^2 + x) dx$$

Here, $f(x) = x^2 + x$ and $a = 0$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[f(0) + f(0+h) + f(0+2h) + \dots + f(0+(n-1)h)]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[f(0) + f(h) + f(2h) + \dots + f((n-1)h)]$$

Now, by putting $x = 0$ in $f(x)$ we get,

$$f(0) = 0^2 + 0 = 0 + 0 = 0$$

$$f(h)$$

$$= (h)^2 + (h)$$

$$= h^2 + h$$

$$\text{Similarly, } f(2h)$$

$$= (2h)^2 + (2h)$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[h^2 + h + (2h)^2 + 2h + \dots + \{(n-1)h\}^2 + (n-1)h]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[h^2 + h + h^2(2)^2 + h(2) + \dots + h^2(n-1)^2 + h(n-1)]$$

Now take h^2 and h common in remaining series

$$\Rightarrow I = \lim_{h \rightarrow 0} h[h^2\{1^2 + 2^2 + \dots + (n-1)^2\} + h\{1 + 2 + \dots + (n-1)\}]$$

$$\left. \begin{aligned} & \left\{ \sum_{i=1}^{n-1} i^2 = 1^2 + 2^2 + \dots + (n-1)^2 = \frac{n(n-1)(2n-1)}{6}; \right. \\ & \left. \sum_{i=1}^{n-1} i = 1 + 2 + \dots + (n-1) = \frac{n(n-1)}{2} \right\} \end{aligned} \right.$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[h^2 \left\{ \frac{n(n-1)(2n-1)}{6} \right\} + h \left\{ \frac{n(n-1)}{2} \right\} \right]$$

Put,

$$h = \frac{2}{n}$$

Since,

$$h \rightarrow 0 \text{ and } h = \frac{2}{n} \Rightarrow n \rightarrow \infty$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{2}{n} \left[\left(\frac{2}{n} \right)^2 \left\{ \frac{n(n-1)(2n-1)}{6} \right\} + \frac{2}{n} \left\{ \frac{n(n-1)}{2} \right\} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{2}{n} \left[\frac{4}{n^2} \left\{ \frac{n(n-1)(2n-1)}{6} \right\} + \frac{2}{n} \left\{ \frac{n(n-1)}{2} \right\} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{2}{n} \left[\frac{2(n-1)(2n-1)}{3n} + (n-1) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[\frac{4(n-1)(2n-1)}{3n \times n} + \frac{2}{n}(n-1) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[\frac{4}{3} \left(\frac{n-1}{n} \right) \left(\frac{2n-1}{n} \right) + 2 \left(\frac{n-1}{n} \right) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[\frac{4}{3} \left(1 - \frac{1}{n} \right) \left(2 - \frac{1}{n} \right) + 2 \left(1 - \frac{1}{n} \right) \right]$$

$$\Rightarrow I = \frac{4}{3} \left(1 - \frac{1}{\infty}\right) \left(2 - \frac{1}{\infty}\right) + 2 \left(1 - \frac{1}{\infty}\right)$$

$$\Rightarrow I = \frac{4}{3} (1 - 0)(2 - 0) + 2(1 - 0)$$

$$\Rightarrow I = \frac{4}{3} \times 1 \times 2 + 2$$

$$\Rightarrow I = \frac{8}{3} + 2$$

$$\Rightarrow I = \frac{8+6}{3}$$

$$\Rightarrow I = \frac{14}{3}$$

Hence, the value of $\int_0^2 (x^2 + x) dx = \frac{14}{3}$

25. Question

Evaluate the following integrals as a limit of sums:

$$\int_0^2 (x^2 + 2x + 1) dx$$

Answer

To find: $\int_0^2 (x^2 + 2x + 1) dx$

Formula used:

$$\int_a^b f(x) dx = \lim_{h \rightarrow 0} h[f(a) + f(a+h) + f(a+2h) + \dots + f(a+(n-1)h)],$$

where,

$$h = \frac{b-a}{n}$$

Here, $a = 0$ and $b = 2$

Therefore,

$$h = \frac{2-0}{n}$$

$$\Rightarrow h = \frac{2}{n}$$

Let,

$$I = \int_0^2 (x^2 + 2x + 1) dx$$

Here, $f(x) = x^2 + 2x + 1$ and $a = 0$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[f(0) + f(0+h) + f(0+2h) + \dots + f(0+(n-1)h)]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[f(0) + f(h) + f(2h) + \dots + f((n-1)h)]$$

Now, by putting $x = 0$ in $f(x)$ we get,

$$f(0) = 0^2 + 2(0) + 1 = 0 + 0 + 1 = 1$$

$f(h)$

$$= (h)^2 + 2(h) + 1$$

$$= h^2 + 2h + 1$$

Similarly, $f(2h)$

$$= (2h)^2 + 2(2h) + 1$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[1 + h^2 + 2h + 1 + (2h)^2 + 2(2h) + 1 + \dots + \{(n-1)h\}^2 + 2(n-1)h + 1]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[1 + h^2 + 2h(1) + 1 + h^2(2)^2 + 2h(2) + 1 + \dots + h^2(n-1)^2 + 2h(n-1) + 1]$$

Since 1 is repeating n times in the series

$$\Rightarrow I = \lim_{h \rightarrow 0} h[1 \times n + h^2 + 2h(1) + h^2(2)^2 + 2h(2) + \dots + h^2(n-1)^2 + 2h(n-1)]$$

Now take h^2 and $2h$ common in remaining series

$$\Rightarrow I = \lim_{h \rightarrow 0} h[n + h^2\{1^2 + 2^2 + \dots + (n-1)^2\} + 2h\{1 + 2 + \dots + (n-1)\}]$$

$$\left. \begin{aligned} & \left. \sum_{i=1}^{n-1} i^2 = 1^2 + 2^2 + \dots + (n-1)^2 = \frac{n(n-1)(2n-1)}{6} ; \right. \\ & \left. \sum_{i=1}^{n-1} i = 1 + 2 + \dots + (n-1) = \frac{n(n-1)}{2} \right. \end{aligned} \right\}$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[n + h^2 \left\{ \frac{n(n-1)(2n-1)}{6} \right\} + 2h \left\{ \frac{n(n-1)}{2} \right\} \right]$$

Put,

$$h = \frac{2}{n}$$

Since,

$$h \rightarrow 0 \text{ and } h = \frac{2}{n} \Rightarrow n \rightarrow \infty$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{2}{n} \left[n + \left(\frac{2}{n} \right)^2 \left\{ \frac{n(n-1)(2n-1)}{6} \right\} + \frac{2(2)}{n} \left\{ \frac{n(n-1)}{2} \right\} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{2}{n} \left[n + \frac{4}{n^2} \left\{ \frac{n(n-1)(2n-1)}{6} \right\} + \frac{2(2)}{n} \left\{ \frac{n(n-1)}{2} \right\} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{2}{n} \left[n + \frac{2(n-1)(2n-1)}{3n} + 2(n-1) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[2 + \frac{4(n-1)(2n-1)}{3n \times n} + \frac{4}{n}(n-1) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[2 + \frac{4}{3} \left(\frac{n-1}{n} \right) \left(\frac{2n-1}{n} \right) + 4 \left(\frac{n-1}{n} \right) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[2 + \frac{4}{3} \left(1 - \frac{1}{n} \right) \left(2 - \frac{1}{n} \right) + 4 \left(1 - \frac{1}{n} \right) \right]$$

$$\Rightarrow I = 2 + \frac{4}{3} \left(1 - \frac{1}{\infty}\right) \left(2 - \frac{1}{\infty}\right) + 4 \left(1 - \frac{1}{\infty}\right)$$

$$\Rightarrow I = 2 + \frac{4}{3} (1 - 0)(2 - 0) + 4(1 - 0)$$

$$\Rightarrow I = 2 + \frac{4}{3} \times 1 \times 2 + 4$$

$$\Rightarrow I = 6 + \frac{8}{3}$$

$$\Rightarrow I = \frac{18 + 8}{3}$$

$$\Rightarrow I = \frac{26}{3}$$

Hence, the value of $\int_0^2 (x^2 + 2x + 1) dx = \frac{26}{3}$

26. Question

Evaluate the following integrals as a limit of sums:

$$\int_0^3 (2x^2 + 3x + 5) dx$$

Answer

To find: $\int_0^3 (2x^2 + 3x + 5) dx$

Formula used:

$$\int_a^b f(x) dx = \lim_{h \rightarrow 0} h [f(a) + f(a+h) + f(a+2h) + \dots + f(a+(n-1)h)],$$

where,

$$h = \frac{b-a}{n}$$

Here, $a = 0$ and $b = 3$

Therefore,

$$h = \frac{3-0}{n}$$

$$\Rightarrow h = \frac{3}{n}$$

Let,

$$I = \int_0^3 (2x^2 + 3x + 5) dx$$

Here, $f(x) = 2x^2 + 3x + 5$ and $a = 0$

$$\Rightarrow I = \lim_{h \rightarrow 0} h [f(0) + f(0+h) + f(0+2h) + \dots + f(0+(n-1)h)]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h [f(0) + f(h) + f(2h) + \dots + f((n-1)h)]$$

Now, by putting $x = 0$ in $f(x)$ we get,

$$f(0) = 2(0)^2 + 3(0) + 5 = 0 + 0 + 5 = 5$$

$f(h)$

$$= 2(h)^2 + 3(h) + 5$$

$$= 2h^2 + 3h + 5$$

Similarly, $f(2h)$

$$= 2(2h)^2 + 3(2h) + 5$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[5 + 2h^2 + 3h + 5 + 2(2h)^2 + 3(2h) + 5 + \dots + 2\{(n-1)h\}^2 + 3(n-1)h + 5]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[5 + 2h^2(1)^2 + 3h(1) + 5 + 2h^2(2)^2 + 3h(2) + 5 + \dots + 2h^2(n-1)^2 + 3h(n-1) + 5]$$

Since 5 is repeating n times in the series

$$\Rightarrow I = \lim_{h \rightarrow 0} h[5 \times n + 2h^2(1)^2 + 3h(1) + 2h^2(2)^2 + 3h(2) + \dots + 2h^2(n-1)^2 + 3h(n-1)]$$

Now take h^2 and $2h$ common in remaining series

$$\Rightarrow I = \lim_{h \rightarrow 0} h[5n + 2h^2\{1^2 + 2^2 + \dots + (n-1)^2\} + 3h\{1 + 2 + \dots + (n-1)\}]$$

$$\left. \begin{cases} \sum_{i=1}^{n-1} i^2 = 1^2 + 2^2 + \dots + (n-1)^2 = \frac{n(n-1)(2n-1)}{6}; \\ \sum_{i=1}^{n-1} i = 1 + 2 + \dots + (n-1) = \frac{n(n-1)}{2} \end{cases} \right\}$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[5n + 2h^2 \left\{ \frac{n(n-1)(2n-1)}{6} \right\} + 3h \left\{ \frac{n(n-1)}{2} \right\} \right]$$

Put,

$$h = \frac{3}{n}$$

Since,

$$h \rightarrow 0 \text{ and } h = \frac{3}{n} \Rightarrow n \rightarrow \infty$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{3}{n} \left[5n + 2 \left(\frac{3}{n} \right)^2 \left\{ \frac{n(n-1)(2n-1)}{6} \right\} + \frac{3(3)}{n} \left\{ \frac{n(n-1)}{2} \right\} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{3}{n} \left[5n + \frac{2(9)}{n^2} \left\{ \frac{n(n-1)(2n-1)}{6} \right\} + \frac{9}{n} \left\{ \frac{n(n-1)}{2} \right\} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{3}{n} \left[5n + \frac{18(n-1)(2n-1)}{6n} + \frac{9}{2}(n-1) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[15 + \frac{54(n-1)(2n-1)}{6n \times n} + \frac{27}{2}(n-1) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[15 + 9 \left(\frac{n-1}{n} \right) \left(\frac{2n-1}{n} \right) + \frac{27}{2} \left(\frac{n-1}{n} \right) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[15 + 9 \left(1 - \frac{1}{n} \right) \left(2 - \frac{1}{n} \right) + \frac{27}{2} \left(1 - \frac{1}{n} \right) \right]$$

$$\Rightarrow I = 15 + 9 \left(1 - \frac{1}{\infty}\right) \left(2 - \frac{1}{\infty}\right) + \frac{27}{2} \left(1 - \frac{1}{\infty}\right)$$

$$\Rightarrow I = 15 + 9(1 - 0)(2 - 0) + \frac{27}{2}(1 - 0)$$

$$\Rightarrow I = 15 + 9 \times 1 \times 2 + \frac{27}{2}$$

$$\Rightarrow I = 15 + 18 + \frac{27}{2}$$

$$\Rightarrow I = 33 + \frac{27}{2}$$

$$\Rightarrow I = \frac{66 + 27}{2}$$

$$\Rightarrow I = \frac{93}{2}$$

Hence, the value of $\int_0^3 (2x^2 + 3x + 5) dx = \frac{93}{2}$

27. Question

Evaluate the following integrals as a limit of sums:

$$\int_a^b x dx$$

Answer

$$\text{To find: } \int_a^b x dx$$

We know,

$$\int_a^b f(x) dx = \lim_{h \rightarrow 0} h[f(a) + f(a+h) + f(a+2h) + \dots + f(a+(n-1)h)],$$

where,

$$h = \frac{b-a}{n}$$

Let,

$$I = \int_a^b x dx$$

Here, $f(x) = x$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[f(a) + f(a+h) + f(a+2h) + \dots + f(a+(n-1)h)]$$

Now, By putting $x = a$ in $f(x)$ we get,

$$f(a) = a$$

Similarly, $f(a+h) = a+h$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[a + a + h + a + 2h + \dots + a + (n-1)h]$$

In this series, a is getting added n times

$$\Rightarrow I = \lim_{h \rightarrow 0} h[an + h + 2h + \dots + (n-1)h]$$



Now take h common in remaining series

$$\Rightarrow I = \lim_{h \rightarrow 0} h [a_n + h(1 + 2 + \dots + (n - 1))]$$

$$\left\{ \because \sum_{i=1}^{n-1} i = 1 + 2 + \dots + (n - 1) = \frac{n(n-1)}{2} \right\}$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[a_n + h \left\{ \frac{n(n-1)}{2} \right\} \right]$$

Put,

$$h = \frac{b-a}{n}$$

Since,

$$h \rightarrow 0 \text{ and } h = \frac{b-a}{n} \Rightarrow n \rightarrow \infty$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{b-a}{n} \left[a_n + \frac{b-a}{n} \left\{ \frac{n(n-1)}{2} \right\} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{b-a}{n} \left[a_n + \left(\frac{b-a}{2} \right) (n-1) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left\{ a(b-a) + \left(\frac{b-a}{2} \right) \frac{(b-a)(n-1)}{n} \right\}$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left\{ a(b-a) + \left(\frac{(b-a)^2}{2} \right) \left(1 - \frac{1}{n} \right) \right\}$$

$$\Rightarrow I = a(b-a) + \left(\frac{(b-a)^2}{2} \right) \left(1 - \frac{1}{\infty} \right)$$

$$\Rightarrow I = a(b-a) + \left(\frac{(b-a)^2}{2} \right) (1 - 0)$$

$$\Rightarrow I = a(b-a) + \frac{(b-a)^2}{2}$$

$$\Rightarrow I = (b-a) \left(a + \frac{b-a}{2} \right)$$

$$\Rightarrow I = (b-a) \left(\frac{2a+b-a}{2} \right)$$

$$\Rightarrow I = (b-a) \left(\frac{b+a}{2} \right)$$

$$\Rightarrow I = \frac{b^2 - a^2}{2}$$

$$\text{Hence, the value of } \int_a^b x \, dx = \frac{b^2 - a^2}{2}$$

28. Question

Evaluate the following integrals as a limit of sums:

$$\int_0^5 (x+1) \, dx$$

Answer

To find: $\int_0^5 (x + 1) dx$

We know,

$$\int_a^b f(x) dx = \lim_{h \rightarrow 0} h[f(a) + f(a+h) + f(a+2h) + \dots + f(a+(n-1)h)],$$

where,

$$h = \frac{b-a}{n}$$

Here, $a = 0$ and $b = 5$

Therefore,

$$h = \frac{5-0}{n}$$

$$\Rightarrow h = \frac{5}{n}$$

Let,

$$I = \int_0^5 (x + 1) dx$$

Here, $f(x) = x + 1$ and $a = 0$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[f(0) + f(0+h) + f(0+2h) + \dots + f(0+(n-1)h)]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[f(0) + f(h) + f(2h) + \dots + f((n-1)h)]$$

Now, By putting $x = 0$ in $f(x)$ we get,

$$f(0) = 0 + 1 = 1$$

Similarly, $f(h) = h + 1$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[1 + h + 1 + 2h + 1 + \dots + (n-1)h + 1]$$

In this series, 1 is getting added n times

$$\Rightarrow I = \lim_{h \rightarrow 0} h[1 \times n + h + 2h + \dots + (n-1)h]$$

Now take h common in remaining series

$$\Rightarrow I = \lim_{h \rightarrow 0} h[n + h(1 + 2 + \dots + (n-1))]$$

$$\left\{ \because \sum_{i=1}^{n-1} i = 1 + 2 + \dots + (n-1) = \frac{n(n-1)}{2} \right\}$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[n + h \left\{ \frac{n(n-1)}{2} \right\} \right]$$

Put,

$$h = \frac{5}{n}$$

Since,

$$h \rightarrow 0 \text{ and } h = \frac{5}{n} \Rightarrow n \rightarrow \infty$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{5}{n} \left[n + \frac{5}{n} \left\{ \frac{n(n-1)}{2} \right\} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{5}{n} \left[n + \frac{5(n-1)}{2} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left\{ 5 + \frac{25(n-1)}{2n} \right\}$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left\{ 5 + \frac{25}{2} \left(1 - \frac{1}{n} \right) \right\}$$

$$\Rightarrow I = 5 + \frac{25}{2} \left(1 - \frac{1}{\infty} \right)$$

$$\Rightarrow I = 5 + \frac{25}{2} (1 - 0)$$

$$\Rightarrow I = 5 + \frac{25}{2}$$

$$\Rightarrow I = \frac{10 + 25}{2}$$

$$\Rightarrow I = \frac{35}{2}$$

Hence, the value of $\int_0^5 (x+1) dx = \frac{35}{2}$

29. Question

Evaluate the following integrals as a limit of sums:

$$\int_2^3 x^2 dx$$

Answer

$$\text{To find: } \int_2^3 x^2 dx$$

Formula used:

$$\int_a^b f(x) dx = \lim_{h \rightarrow 0} h [f(a) + f(a+h) + f(a+2h) + \dots + f(a+(n-1)h)],$$

where,

$$h = \frac{b-a}{n}$$

Here, $a = 2$ and $b = 3$

Therefore,

$$h = \frac{3-2}{n}$$

$$\Rightarrow h = \frac{1}{n}$$

Let,

$$I = \int_2^3 x^2 dx$$

Here, $f(x) = x^2$ and $a = 2$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[f(2) + f(2+h) + f(2+2h) + \dots + f(2+(n-1)h)]$$

Now, by putting $x = 2$ in $f(x)$ we get,

$$f(2) = 2^2 = 4$$

$$f(2+h)$$

$$= (2+h)^2$$

$$= h^2 + 2^2 + 2(h)(2)$$

$$= h^2 + 4 + 4(h)$$

Similarly, $f(2+2h)$

$$= (2+2h)^2$$

$$= (2h)^2 + 2^2 + 2(2h)(2)$$

$$= (2h)^2 + 4 + 4(2h)$$

$$\{\because (x+y)^2 = x^2 + y^2 + 2xy\}$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[4 + h^2 + 4 + 4(h) + (2h)^2 + 4 + 4(2h) + \dots + \{(n-1)h\}^2 + 4 + 4\{(n-1)h\}]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[4 + h^2 + 4 + 4h + h^2(2)^2 + 4 + 4(2h) + \dots + h^2(n-1)^2 + 4 + 4h(n-1)]$$

In this series, 4 is getting added n times

$$\Rightarrow I = \lim_{h \rightarrow 0} h[4 \times n + h^2 + 4h + h^2(2)^2 + 4(2h) + \dots + h^2(n-1)^2 + 4h(n-1)]$$

Now take h^2 and $4h$ common in remaining series

$$\Rightarrow I = \lim_{h \rightarrow 0} h[4n + h^2\{1^2 + 2^2 + \dots + (n-1)^2\} + 4h\{1 + 2 + \dots + (n-1)\}]$$

$$\left\{ \begin{array}{l} \because \sum_{i=1}^{n-1} i^2 = 1^2 + 2^2 + \dots + (n-1)^2 = \frac{n(n-1)(2n-1)}{6}; \\ \sum_{i=1}^{n-1} i = 1 + 2 + \dots + (n-1) = \frac{n(n-1)}{2} \end{array} \right\}$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[4n + h^2 \left\{ \frac{n(n-1)(2n-1)}{6} \right\} + 4h \left\{ \frac{n(n-1)}{2} \right\} \right]$$

Put,

$$h = \frac{1}{n}$$

Since,

$$h \rightarrow 0 \text{ and } h = \frac{1}{n} \Rightarrow n \rightarrow \infty$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} n \left[4n + \left(\frac{1}{n} \right)^2 \left\{ \frac{n(n-1)(2n-1)}{6} \right\} + \frac{4(1)}{n} \left\{ \frac{n(n-1)}{2} \right\} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{1}{n} \left[4n + \frac{1}{n^2} \left\{ \frac{n(n-1)(2n-1)}{6} \right\} + \frac{4}{n} \left\{ \frac{n(n-1)}{2} \right\} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{1}{n} \left[4n + \frac{(n-1)(2n-1)}{6n} + 2(n-1) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[4 + \frac{(n-1)(2n-1)}{6n \times n} + \frac{2}{n}(n-1) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[4 + \frac{1}{6} \left(\frac{n-1}{n} \right) \left(\frac{2n-1}{n} \right) + 2 \left(\frac{n-1}{n} \right) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[4 + \frac{1}{6} \left(1 - \frac{1}{n} \right) \left(2 - \frac{1}{n} \right) + 2 \left(1 - \frac{1}{n} \right) \right]$$

$$\Rightarrow I = 4 + \frac{1}{6} \left(1 - \frac{1}{\infty} \right) \left(2 - \frac{1}{\infty} \right) + 2 \left(1 - \frac{1}{\infty} \right)$$

$$\Rightarrow I = 4 + \frac{1}{6} (1 - 0)(2 - 0) + 2(1 - 0)$$

$$\Rightarrow I = 4 + \frac{1}{6} \times 1 \times 2 + 2$$

$$\Rightarrow I = 6 + \frac{1}{3}$$

$$\Rightarrow I = \frac{18 + 1}{3}$$

$$\Rightarrow I = \frac{19}{3}$$

Hence, the value of $\int_2^3 x^2 dx = \frac{19}{3}$

30. Question

Evaluate the following integrals as a limit of sums:

$$\int_1^3 (x^2 + x) dx$$

Answer

$$\text{To find: } \int_1^3 (x^2 + x) dx$$

Formula used:

$$\int_a^b f(x) dx = \lim_{h \rightarrow 0} h [f(a) + f(a+h) + f(a+2h) + \dots + f(a+(n-1)h)],$$

where,

$$h = \frac{b-a}{n}$$

Here, $a = 1$ and $b = 3$

Therefore,

$$h = \frac{3-1}{n}$$

$$\Rightarrow h = \frac{2}{n}$$

Let,

$$I = \int_1^3 (x^2 + x) dx$$

Here, $f(x) = x^2 + x$ and $a = 1$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[f(1) + f(1+h) + f(1+2h) + \dots + f(1+(n-1)h)]$$

Now, by putting $x = 1$ in $f(x)$ we get,

$$f(1) = 1^2 + 1 = 1 + 1 = 2$$

$$f(1+h)$$

$$= (1+h)^2 + (1+h)$$

$$= h^2 + 1^2 + 2(h)(1) + 1 + h$$

$$= h^2 + 2h + h + 1 + 1$$

$$= h^2 + 3h + 2$$

Similarly, $f(1+2h)$

$$= (1+2h)^2 + (1+2h)$$

$$= (2h)^2 + 1^2 + 2(2h)(1) + 1 + 2h$$

$$= (2h)^2 + 4h + 2h + 1 + 1$$

$$= (2h)^2 + 6h + 2$$

$$\{\because (x+y)^2 = x^2 + y^2 + 2xy\}$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[2 + h^2 + 3h + 2 + (2h)^2 + 6h + 2 + \dots + \{(n-1)h\}^2 + 3(n-1)h + 2]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[2 + h^2 + 3h(1) + 2 + h^2(2)^2 + 3h(2) + 2 + \dots + h^2(n-1)^2 + 3h(n-1) + 2]$$

In this series, 2 is getting added n times

$$\Rightarrow I = \lim_{h \rightarrow 0} h[2n + h^2 + 3h(1) + h^2(2)^2 + 3h(2) + \dots + h^2(n-1)^2 + 3h(n-1)]$$

Now take h^2 and $3h$ common in remaining series

$$\Rightarrow I = \lim_{h \rightarrow 0} h[2n + h^2\{1^2 + 2^2 + \dots + (n-1)^2\} + 3h\{1 + 2 + \dots + (n-1)\}]$$

$$\left\{ \begin{array}{l} \because \sum_{i=1}^{n-1} i^2 = 1^2 + 2^2 + \dots + (n-1)^2 = \frac{n(n-1)(2n-1)}{6}; \\ \sum_{i=1}^{n-1} i = 1 + 2 + \dots + (n-1) = \frac{n(n-1)}{2} \end{array} \right\}$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[2n + h^2 \left\{ \frac{n(n-1)(2n-1)}{6} \right\} + 3h \left\{ \frac{n(n-1)}{2} \right\} \right]$$

Put,

$$h = \frac{2}{n}$$

Since,

$$h \rightarrow 0 \text{ and } h = \frac{2}{n} \Rightarrow n \rightarrow \infty$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{2}{n} \left[2n + \left(\frac{2}{n} \right)^2 \left\{ \frac{n(n-1)(2n-1)}{6} \right\} + \frac{2(3)}{n} \left\{ \frac{n(n-1)}{2} \right\} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{2}{n} \left[2n + \frac{4}{n^2} \left\{ \frac{n(n-1)(2n-1)}{6} \right\} + \frac{6}{n} \left\{ \frac{n(n-1)}{2} \right\} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{2}{n} \left[2n + \frac{2(n-1)(2n-1)}{3n} + 3(n-1) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[4 + \frac{4(n-1)(2n-1)}{3n \times n} + \frac{6}{n}(n-1) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[4 + \frac{4}{3} \left(\frac{n-1}{n} \right) \left(\frac{2n-1}{n} \right) + 6 \left(\frac{n-1}{n} \right) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[4 + \frac{4}{3} \left(1 - \frac{1}{n} \right) \left(2 - \frac{1}{n} \right) + 6 \left(1 - \frac{1}{n} \right) \right]$$

$$\Rightarrow I = 4 + \frac{4}{3} \left(1 - \frac{1}{\infty} \right) \left(2 - \frac{1}{\infty} \right) + 6 \left(1 - \frac{1}{\infty} \right)$$

$$\Rightarrow I = 4 + \frac{4}{3} (1 - 0) (2 - 0) + 6(1 - 0)$$

$$\Rightarrow I = 4 + \frac{4}{3} \times 1 \times 2 + 6$$

$$\Rightarrow I = 10 + \frac{8}{3}$$

$$\Rightarrow I = \frac{30 + 8}{3}$$

$$\Rightarrow I = \frac{38}{3}$$

Hence, the value of $\int_1^3 (x^2 + x) dx = \frac{38}{3}$

31. Question

Evaluate the following integrals as a limit of sums:

$$\int_0^2 (x^2 - x) dx$$

Answer

$$\text{To find: } \int_0^2 (x^2 - x) dx$$

Formula used:

$$\int_a^b f(x) dx = \lim_{h \rightarrow 0} h [f(a) + f(a+h) + f(a+2h) + \dots + f(a+(n-1)h)],$$

where,

$$h = \frac{b-a}{n}$$

Here, $a = 0$ and $b = 2$

Therefore,

$$h = \frac{2 - 0}{n}$$

$$\Rightarrow h = \frac{2}{n}$$

Let,

$$I = \int_0^2 (x^2 - x) dx$$

Here, $f(x) = x^2 - x$ and $a = 0$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[f(0) + f(0+h) + f(0+2h) + \dots + f(0+(n-1)h)]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[f(0) + f(h) + f(2h) + \dots + f((n-1)h)]$$

Now, by putting $x = 0$ in $f(x)$ we get,

$$f(0) = 0^2 - 0 = 0 - 0 = 0$$

$f(h)$

$$= (h)^2 - (h)$$

$$= h^2 - h$$

Similarly, $f(2h)$

$$= (2h)^2 - (2h)$$

$$= (2h)^2 - 2h$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[0 + h^2 - h + (2h)^2 - 2h + \dots + \{(n-1)h\}^2 - (n-1)h]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[h^2 - h + h^2(2)^2 - 2h + \dots + h^2(n-1)^2 - h(n-1)]$$

Now take h^2 and $-h$ common in remaining series

$$\Rightarrow I = \lim_{h \rightarrow 0} h[h^2\{1^2 + 2^2 + \dots + (n-1)^2\} - h\{1 + 2 + \dots + (n-1)\}]$$

$$\left\{ \begin{array}{l} \because \sum_{i=1}^{n-1} i^2 = 1^2 + 2^2 + \dots + (n-1)^2 = \frac{n(n-1)(2n-1)}{6}; \\ \sum_{i=1}^{n-1} i = 1 + 2 + \dots + (n-1) = \frac{n(n-1)}{2} \end{array} \right\}$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[h^2 \left\{ \frac{n(n-1)(2n-1)}{6} \right\} - h \left\{ \frac{n(n-1)}{2} \right\} \right]$$

Put,

$$h = \frac{2}{n}$$

Since,

$$h \rightarrow 0 \text{ and } h = \frac{2}{n} \Rightarrow n \rightarrow \infty$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{2}{n} \left[\left(\frac{2}{n} \right)^2 \left\{ \frac{n(n-1)(2n-1)}{6} \right\} - \frac{2}{n} \left\{ \frac{n(n-1)}{2} \right\} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{2}{n} \left[\frac{4}{n^2} \left\{ \frac{n(n-1)(2n-1)}{6} \right\} - \frac{2}{n} \left\{ \frac{n(n-1)}{2} \right\} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{2}{n} \left[\frac{2(n-1)(2n-1)}{3n} - (n-1) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[\frac{4(n-1)(2n-1)}{3n \times n} - \frac{2}{n}(n-1) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[\frac{4}{3} \left(\frac{n-1}{n} \right) \left(\frac{2n-1}{n} \right) - 2 \left(\frac{n-1}{n} \right) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[\frac{4}{3} \left(1 - \frac{1}{n} \right) \left(2 - \frac{1}{n} \right) - 2 \left(1 - \frac{1}{n} \right) \right]$$

$$\Rightarrow I = \frac{4}{3} \left(1 - \frac{1}{\infty} \right) \left(2 - \frac{1}{\infty} \right) - 2 \left(1 - \frac{1}{\infty} \right)$$

$$\Rightarrow I = \frac{4}{3} (1 - 0) (2 - 0) - 2(1 - 0)$$

$$\Rightarrow I = \frac{4}{3} \times 1 \times 2 - 2$$

$$\Rightarrow I = \frac{8}{3} - 2$$

$$\Rightarrow I = \frac{8-6}{3}$$

$$\Rightarrow I = \frac{2}{3}$$

Hence, the value of $\int_0^2 (x^2 - x) dx = \frac{2}{3}$

32. Question

Evaluate the following integrals as a limit of sums:

$$\int_1^3 (2x^2 + 5x) dx$$

Answer

$$\text{To find: } \int_1^3 (2x^2 + 5x) dx$$

Formula used:

$$\int_a^b f(x) dx = \lim_{h \rightarrow 0} h [f(a) + f(a+h) + f(a+2h) + \dots + f(a+(n-1)h)],$$

where,

$$h = \frac{b-a}{n}$$

Here, $a = 1$ and $b = 3$

Therefore,

$$h = \frac{3-1}{n}$$

$$\Rightarrow h = \frac{2}{n}$$

Let,

$$I = \int_1^3 (2x^2 + 5x) dx$$

Here, $f(x) = 2x^2 + 5x$ and $a = 1$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[f(1) + f(1+h) + f(1+2h) + \dots + f(1+(n-1)h)]$$

Now, by putting $x = 1$ in $f(x)$ we get,

$$f(1) = 2(1)^2 + 5(1) = 2 + 5 = 7$$

$$f(1+h)$$

$$= 2(1+h)^2 + 5(1+h)$$

$$= 2\{h^2 + 1^2 + 2(h)(1)\} + 5 + 5h$$

$$= 2h^2 + 4h + 2 + 5 + 5h$$

$$= 2h^2 + 9h + 7$$

$$\text{Similarly, } f(1+2h)$$

$$= 2(1+2h)^2 + 5(1+2h)$$

$$= 2\{(2h)^2 + 1^2 + 2(2h)(1)\} + 5 + 10h$$

$$= 2(2h)^2 + 2 + 8h + 5 + 10h$$

$$= 2(2h)^2 + 18h + 7$$

$$= 2(2h)^2 + 9(2h) + 7$$

$$\{\because (x+y)^2 = x^2 + y^2 + 2xy\}$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[7 + 2h^2 + 9h + 7 + 2(2h)^2 + 9(2h) + 7 + \dots + 2\{(n-1)h\}^2 + 9(n-1)h + 7]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[7 + 2h^2(1) + 9h(1) + 7 + 2h^2(2)^2 + 9h(2) + 7 + \dots + 2h^2(n-1)^2 + 9h(n-1) + 7]$$

In this series, 7 is getting added n times

$$\Rightarrow I = \lim_{h \rightarrow 0} h[7n + 2h^2(1) + 9h(1) + 2h^2(2)^2 + 9h(2) + \dots + 2h^2(n-1)^2 + 9h(n-1)]$$

Now take $2h^2$ and $9h$ common in remaining series

$$\Rightarrow I = \lim_{h \rightarrow 0} h[7n + 2h^2\{1^2 + 2^2 + \dots + (n-1)^2\} + 9h\{1 + 2 + \dots + (n-1)\}]$$

$$\left\{ \begin{array}{l} \because \sum_{i=1}^{n-1} i^2 = 1^2 + 2^2 + \dots + (n-1)^2 = \frac{n(n-1)(2n-1)}{6}; \\ \sum_{i=1}^{n-1} i = 1 + 2 + \dots + (n-1) = \frac{n(n-1)}{2} \end{array} \right\}$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[7n + 2h^2 \left\{ \frac{n(n-1)(2n-1)}{6} \right\} + 9h \left\{ \frac{n(n-1)}{2} \right\} \right]$$

Put,

$$h = \frac{2}{n}$$

Since,

$$h \rightarrow 0 \text{ and } h = \frac{2}{n} \Rightarrow n \rightarrow \infty$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{2}{n} \left[7n + 2 \left(\frac{2}{n} \right)^2 \left\{ \frac{n(n-1)(2n-1)}{6} \right\} + \frac{2(9)}{n} \left\{ \frac{n(n-1)}{2} \right\} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{2}{n} \left[7n + \frac{8}{n^2} \left\{ \frac{n(n-1)(2n-1)}{6} \right\} + \frac{18}{n} \left\{ \frac{n(n-1)}{2} \right\} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{2}{n} \left[7n + \frac{4(n-1)(2n-1)}{3n} + 9(n-1) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[14 + \frac{8(n-1)(2n-1)}{3n \times n} + \frac{18}{n}(n-1) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[14 + \frac{8}{3} \left(\frac{n-1}{n} \right) \left(\frac{2n-1}{n} \right) + 18 \left(\frac{n-1}{n} \right) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[14 + \frac{8}{3} \left(1 - \frac{1}{n} \right) \left(2 - \frac{1}{n} \right) + 18 \left(1 - \frac{1}{n} \right) \right]$$

$$\Rightarrow I = 14 + \frac{8}{3} \left(1 - \frac{1}{\infty} \right) \left(2 - \frac{1}{\infty} \right) + 18 \left(1 - \frac{1}{\infty} \right)$$

$$\Rightarrow I = 14 + \frac{8}{3} (1 - 0)(2 - 0) + 18(1 - 0)$$

$$\Rightarrow I = 14 + \frac{8}{3} \times 1 \times 2 + 18$$

$$\Rightarrow I = 32 + \frac{16}{3}$$

$$\Rightarrow I = \frac{96 + 16}{3}$$

$$\Rightarrow I = \frac{112}{3}$$

Hence, the value of $\int_1^3 (2x^2 + 5x) dx = \frac{112}{3}$

33. Question

Evaluate the following integrals as a limit of sums:

$$\int_1^3 (3x^2 + 1) dx$$

Answer

$$\text{To find: } \int_1^3 (3x^2 + 1) dx$$

Formula used:

$$\int_a^b f(x) dx = \lim_{h \rightarrow 0} h[f(a) + f(a+h) + f(a+2h) + \dots + f(a+(n-1)h)],$$

where,

$$h = \frac{b-a}{n}$$

Here, $a = 1$ and $b = 3$

Therefore,

$$h = \frac{3-1}{n}$$

$$\Rightarrow h = \frac{2}{n}$$

Let,

$$I = \int_1^3 (3x^2 + 1) dx$$

Here, $f(x) = 3x^2 + 1$ and $a = 1$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[f(1) + f(1+h) + f(1+2h) + \dots + f(1+(n-1)h)]$$

Now, by putting $x = 1$ in $f(x)$ we get,

$$f(1) = 3(1^2) + 1 = 3(1) + 1 = 3 + 1 = 4$$

$$f(1+h)$$

$$= 3(1+h)^2 + 1$$

$$= 3\{h^2 + 1^2 + 2(h)(1)\} + 1$$

$$= 3(h)^2 + 3 + 3(2h) + 1$$

$$= 3(h)^2 + 4 + 6h$$

Similarly, $f(1+2h)$

$$= 3(1+2h)^2 + 1$$

$$= 3\{2(2h)^2 + 1^2 + 2(2h)(1)\} + 1$$

$$= 3(2h)^2 + 3 + 3(4h) + 1$$

$$= 3(2h)^2 + 4 + 12h$$

$$\{\because (x+y)^2 = x^2 + y^2 + 2xy\}$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[4 + 3(h)^2 + 4 + 6h + 3(2h)^2 + 4 + 12h + \dots + 3\{(n-1)h\}^2 + 4 + 6\{(n-1)h\}]$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h[4 + 3h^2(1^2) + 4 + 6h + 3h^2(2^2) + 4 + 6(2h) + \dots + 3h^2(n-1)^2 + 4 + 6h(n-1)]$$

In this series, 4 is getting added n times

$$\Rightarrow I = \lim_{h \rightarrow 0} h[4 \times n + 3h^2(1^2) + 6h + 3h^2(2^2) + 6(2h) + \dots + 3h^2(n-1)^2 + 6h(n-1)]$$

Now take $3h^2$ and $6h$ common in remaining series

$$\Rightarrow I = \lim_{h \rightarrow 0} h[4n + 3h^2\{1^2 + 2^2 + \dots + (n-1)^2\} + 6h\{1 + 2 + \dots + (n-1)\}]$$

$$\left\{ \begin{array}{l} \because \sum_{i=1}^{n-1} i^2 = 1^2 + 2^2 + \dots + (n-1)^2 = \frac{n(n-1)(2n-1)}{6}; \\ \quad \sum_{i=1}^{n-1} i = 1 + 2 + \dots + (n-1) = \frac{n(n-1)}{2} \end{array} \right\}$$

$$\Rightarrow I = \lim_{h \rightarrow 0} h \left[4n + 3h^2 \left\{ \frac{n(n-1)(2n-1)}{6} \right\} + 6h \left\{ \frac{n(n-1)}{2} \right\} \right]$$

Put,

$$h = \frac{2}{n}$$

Since,

$$h \rightarrow 0 \text{ and } h = \frac{2}{n} \Rightarrow n \rightarrow \infty$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{2}{n} \left[4n + 3 \left(\frac{2}{n} \right)^2 \left\{ \frac{n(n-1)(2n-1)}{6} \right\} + \frac{6(2)}{n} \left\{ \frac{n(n-1)}{2} \right\} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{2}{n} \left[4n + \frac{12}{n^2} \left\{ \frac{n(n-1)(2n-1)}{6} \right\} + \frac{12}{n} \left\{ \frac{n(n-1)}{2} \right\} \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \frac{2}{n} \left[4n + \frac{2(n-1)(2n-1)}{n} + 6(n-1) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[8 + \frac{4(n-1)(2n-1)}{n \times n} + \frac{12}{n}(n-1) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[8 + 4 \left(\frac{n-1}{n} \right) \left(\frac{2n-1}{n} \right) + 12 \left(\frac{n-1}{n} \right) \right]$$

$$\Rightarrow I = \lim_{n \rightarrow \infty} \left[8 + 4 \left(1 - \frac{1}{n} \right) \left(2 - \frac{1}{n} \right) + 12 \left(1 - \frac{1}{n} \right) \right]$$

$$\Rightarrow I = 8 + 4 \left(1 - \frac{1}{\infty} \right) \left(2 - \frac{1}{\infty} \right) + 12 \left(1 - \frac{1}{\infty} \right)$$

$$\Rightarrow I = 8 + 4(1-0)(2-0) + 12(1-0)$$

$$\Rightarrow I = 8 + 4 \times 1 \times 2 + 12$$

$$\Rightarrow I = 20 + 8$$

$$\Rightarrow I = 28$$

Hence, the value of $\int_1^3 (3x^2 + 1) dx = 28$

Very short answer

1. Question

$$\text{Evaluate } \int_0^{\pi/2} \sin^2 x dx$$

Answer

$$\text{Let } I = \int_0^{\pi/2} \sin^2 x dx - (1)$$

Using the property that $\int_a^b f(x)dx = \int_a^b f(a+b-x)dx$

$$I = \int_0^{\frac{\pi}{2}} \sin^2\left(\frac{\pi}{2} - x\right) dx$$

$$= \int_0^{\frac{\pi}{2}} \cos^2 x dx \quad \text{-(2)}$$

Adding (1) and (2), we get

$$2I = \int_0^{\frac{\pi}{2}} \sin^2 x dx + \int_0^{\frac{\pi}{2}} \cos^2 x dx = \int_0^{\frac{\pi}{2}} dx = \frac{\pi}{2}$$

$$\text{Hence, } I = \frac{\pi}{4}$$

2. Question

Evaluate $\int_0^{\pi/2} \cos^2 x dx$

Answer

$$\text{Let } I = \int_0^{\frac{\pi}{2}} \cos^2 x dx \quad \text{-(1)}$$

Using the property that $\int_a^b f(x)dx = \int_a^b f(a+b-x)dx$

$$I = \int_0^{\frac{\pi}{2}} \cos^2\left(\frac{\pi}{2} - x\right) dx = \int_0^{\frac{\pi}{2}} \sin^2 x dx \quad \text{-(2)}$$

Adding (1) and (2), we get

$$2I = \int_0^{\frac{\pi}{2}} \sin^2 x dx + \int_0^{\frac{\pi}{2}} \cos^2 x dx = \int_0^{\frac{\pi}{2}} dx = \frac{\pi}{2}$$

$$\text{Hence, } I = \frac{\pi}{4}$$

3. Question

Evaluate $\int_{-\pi/2}^{\pi/2} \sin^2 x dx$

Answer

$$\text{Let } I = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \sin^2 x dx$$

$$\text{Since } \cos 2x = 1 - 2\sin^2 x \Rightarrow \sin^2 x = \frac{1 - \cos 2x}{2}$$

$$I = \frac{1}{2} \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} (1 - \cos 2x) dx$$

$$= \frac{x}{2} \left| \frac{\pi}{2} - \frac{\sin 2x}{4} \right|_{-\frac{\pi}{2}}^{\frac{\pi}{2}}$$

$$= \frac{\pi}{2} + \frac{1}{4} (\sin \pi - \sin(-\pi))$$

$$= \frac{\pi}{2}$$

4. Question

Evaluate $\int_{-\pi/2}^{\pi/2} \cos^2 x \, dx$

Answer

Let $I = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \cos^2 x \, dx$

Since $\cos 2x = 2\cos^2 x - 1$

$$\Rightarrow \cos^2 x = \frac{1 + \cos 2x}{2}$$

$$I = \frac{1}{2} \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} (1 + \cos 2x) \, dx$$

$$= \frac{x}{2} \left| \frac{\pi}{2} + \frac{\sin 2x}{4} \right|_{-\frac{\pi}{2}}^{\frac{\pi}{2}}$$

$$= \frac{\pi}{2} - \frac{1}{4} (\sin \pi - \sin(-\pi))$$

$$= \frac{\pi}{2}$$

5. Question

Evaluate $\int_{-\pi/2}^{\pi/2} \sin^3 x \, dx$

Answer

Let $I = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \sin^3 x \, dx$

$$f(x) = \sin^3 x$$

$$f(-x) = \sin^3(-x) = -\sin^3 x$$

Hence, $f(x)$ is an odd function.

Since, $\int_{-a}^a f(x) \, dx = 0$ if $f(x)$ is an odd function.

Therefore, $I=0$.

6. Question

Evaluate $\int_{-\pi/2}^{\pi/2} x \cos^2 x \, dx$

Answer

Let $I = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} x \cos^2 x \, dx$

$$f(x) = x \cos^2 x$$

$$f(-x) = (-x) \cos^2(-x)$$

$$= -x \cos^2 x$$

=-f(x)

Hence, f(x) is an odd function.

Since, $\int_{-a}^a f(x) dx = 0$ if f(x) is an odd function.

Therefore, I=0.

7. Question

Evaluate $\int_0^{\pi/4} \tan^2 x dx$

Answer

Let $I = \int_0^{\pi/4} \tan^2 x dx = \int_0^{\pi/4} (\sec^2 x - 1) dx$

Let $\tan x=t$

$\Rightarrow \sec^2 x dx = dt$

When $x=0, t=0$ and when $x = \frac{\pi}{4}, t = 1$

Hence, $I = \int_0^1 dt + \int_0^{\pi/4} -dx = 1 - \frac{\pi}{4}$

8. Question

Evaluate $\int_0^1 \frac{1}{x^2+1} dx$

Answer

Let $I = \int_0^1 \frac{1}{1+x^2} dx$

Substituting $x=\tan\theta \Rightarrow dx=\sec^2\theta d\theta$ (By differentiating both sides)

Also, when $x=0, \theta=0$ and $x=1, \theta=\frac{\pi}{4}$

We get $I = \int_0^{\pi/4} \frac{1}{1+\tan^2\theta} \sec^2\theta d\theta$

Since $\sec^2\theta=1+\tan^2\theta$

We get $I = \int_0^{\pi/4} d\theta$

$$= \frac{\pi}{4}$$

9. Question

Evaluate $\int_{-2}^1 \frac{|x|}{x} dx$

Answer

Let $I = \int_{-2}^1 \frac{|x|}{x} dx = \int_{-2}^0 \frac{|x|}{x} dx + \int_0^1 \frac{|x|}{x} dx$

$|x|=-x, \text{ if } x<0$

And $|x|=x, \text{ if } x\geq 0$

Hence, $I = \int_{-2}^0 -dx + \int_0^1 dx$

$$I = -2 + 1 = -1$$

10. Question

Evaluate $\int_0^{\infty} e^{-x} dx$

Answer

$$\text{Let } I = \int_0^{\infty} e^{-x} dx$$

$$= -e^{-x} \Big|_0^{\infty}$$

$$= -(0-1)$$

$$= 1$$

11. Question

Evaluate $\int_0^4 \frac{1}{\sqrt{16-x^2}} dx$

Answer

$$\text{Let } I = \int_0^4 \frac{1}{\sqrt{16-x^2}} dx$$

$$\text{Substituting } x=4\sin\theta \Rightarrow dx=4\cos\theta d\theta$$

$$\text{Also, When } x=0, \theta=0 \text{ and } x=4, \theta=\frac{\pi}{2}$$

$$I = \int_0^{\frac{\pi}{2}} \frac{1}{\sqrt{16-16\sin^2\theta}} 4\cos\theta d\theta$$

$$= \int_0^{\frac{\pi}{2}} d\theta = \frac{\pi}{2}$$

12. Question

Evluate $\int_0^3 \frac{1}{x^2+9} dx$

Answer

$$\text{Substituting } x=3\tan\theta \Rightarrow dx=3\sec^2\theta d\theta \text{ (By differentiating both sides)}$$

$$\text{Also, when } x=0, \theta=0 \text{ and } x=3, \theta=\frac{\pi}{4}$$

$$\text{We get } I = \int_0^{\frac{\pi}{4}} \frac{1}{9+9\tan^2\theta} 3\sec^2\theta d\theta$$

$$\text{Since } \sec^2\theta=1+\tan^2\theta$$

$$\text{We get } I = \int_0^{\frac{\pi}{4}} \frac{d\theta}{3}$$

$$= \frac{\pi}{12}$$

13. Question

Evaluate $\int_0^{\pi/2} \sqrt{1 - \cos 2x} dx$

Answer

Let $I = \int_0^{\pi/2} \sqrt{1 - \cos 2x} dx$

Since, $\cos 2x = 1 - 2\sin^2 x \Rightarrow 2\sin^2 x = 1 - \cos 2x$

Hence, $I = \int_0^{\pi/2} \sqrt{2\sin x dx} = -\sqrt{2}\cos x \Big|_0^{\pi/2}$

$$= -\sqrt{2}(0 - 1) = \sqrt{2}$$

14. Question

Evaluate $\int_0^{\pi/2} \log \tan x dx$

Answer

Let $I = \int_0^{\pi/2} \log \tan x dx$

Using the property that $\int_a^b f(x) dx = \int_a^b f(a + b - x) dx$

$$I = \int_0^{\pi/2} \log \tan \left(\frac{\pi}{2} - x \right) dx = \int_0^{\pi/2} \log \cot x dx$$

$$= \int_0^{\pi/2} \log \frac{1}{\tan x} dx = - \int_0^{\pi/2} \log \tan x dx$$

$$= -I$$

$$(\text{Since } \log_a \frac{1}{b} = -\log_a b)$$

Since $I = -I$, therefore $I = 0$

15. Question

Evaluate $\int_0^{\pi/2} \log \left(\frac{3 + 5\cos x}{3 + 5\sin x} \right) dx$

Answer

Let $I = \int_0^{\pi/2} \log \frac{3 + 5\cos x}{3 + 5\sin x} dx$

Using the property that $\int_a^b f(x) dx = \int_a^b f(a + b - x) dx$

$$I = \int_0^{\pi/2} \log \frac{3 + 5\cos \left(\frac{\pi}{2} - x \right)}{3 + 5\sin \left(\frac{\pi}{2} - x \right)} dx$$

$$= \int_0^{\pi/2} \log \frac{3 + 5\sin x}{3 + 5\cos x} dx$$

$$= - \int_0^{\pi/2} \log \frac{3 + 5\cos x}{3 + 5\sin x} dx$$

$$= -I \text{ (Since } \log_a \frac{1}{b} = -\log_a b)$$

Since $I = -I$, therefore $I = 0$

16. Question

$$\text{Evaluate } \int_0^{\pi/2} \frac{\sin^n x}{\sin^n x + \cos^n x} dx, n \in \mathbb{N}.$$

Answer

$$\text{Let } I = \int_0^{\pi/2} \frac{\sin^n x}{\sin^n x + \cos^n x} dx \quad (1)$$

Using the property that $\int_a^b f(x) dx = \int_a^b f(a+b-x) dx$

$$\text{We get } I = \int_0^{\pi/2} \frac{\sin^n x}{\sin^n x + \cos^n x} dx$$

$$= \int_0^{\pi/2} \frac{\sin^n(\frac{\pi}{2}-x)}{\sin^n(\frac{\pi}{2}-x) + \cos^n(\frac{\pi}{2}-x)} dx$$

Since $\sin(\frac{\pi}{2}-x) = \cos x$ and $\cos(\frac{\pi}{2}-x) = \sin x$

$$\text{We get } I = \int_0^{\pi/2} \frac{\cos^n x}{\sin^n x + \cos^n x} dx \quad (2)$$

Add (1) and (2)

$$2I = \int_0^{\pi/2} \frac{\sin^n x + \cos^n x}{\sin^n x + \cos^n x} dx$$

$$= \int_0^{\pi/2} dx$$

$$= \frac{\pi}{2}$$

$$I = \frac{\pi}{4}$$

17. Question

$$\text{Evaluate } \int_0^{\pi} \cos^5 x dx$$

Answer

$$\text{Let } I = \int_0^{\pi} \cos^5 x dx$$

Consider $\cos^5 x = \cos^4 x \times \cos x$

$$= (\cos^2 x)^2 \times \cos x$$

$$= (1 - \sin^2 x)^2 \cos x$$

Let $\sin x = y \Rightarrow \cos x dx = dy$ (Differentiating both sides)

Also, when $x=0, y=0$ and $x=\pi, y=0$

$$\text{Hence, } I \text{ become } \int_0^0 (1 - y^2)^2 dy$$

Since $\int_a^a f(x)dx = 0$, We get $I = \int_0^\pi \cos^5 x dx$

$$= \int_0^0 (1 - y^2)^2 dy$$

=0

18. Question

Evaluate $\int_{-\pi/2}^{\pi/2} \log\left(\frac{a - \sin \theta}{a + \sin \theta}\right) d\theta$

Answer

Let $I = \int_{-\pi/2}^{\pi/2} \log\left(\frac{(a - \sin \theta)}{(a + \sin \theta)}\right) d\theta$

Let $f(\theta) = \log\left(\frac{(a - \sin \theta)}{(a + \sin \theta)}\right)$

Then $f(-\theta) = \log\left(\frac{(a - \sin(-\theta))}{(a + \sin(-\theta))}\right)$

$$= \log\left(\frac{(a + \sin \theta)}{(a - \sin \theta)}\right)$$

$$= -\log\left(\frac{(a - \sin \theta)}{(a + \sin \theta)}\right)$$

$$= -f(\theta)$$

(Since $\sin(-\theta) = -\sin(\theta)$ and $\log_a b = -\log_a \frac{1}{b}$)

From this, we infer that $f(\theta)$ is an odd function.

Using $\int_{-a}^a f(x)dx = 0$ if $f(x)$ is an odd function, we get that $I=0$

19. Question

Evaluate $\int_{-1}^1 x|x| dx$

Answer

Let $I = \int_{-1}^1 x|x| dx$

$|x| = -x$, if $x < 0$

And $|x| = x$, if $x \geq 0$

Therefore $f(x) = x|x| = -x^2$, if $x < 0$

And $f(x) = x|x| = x^2$, if $x \geq 0$

Consider $x \geq 0 \Rightarrow f(x) = x^2$

Then $-x < 0 \Rightarrow f(-x) = -(-x)^2 = -f(x)$

Now Consider $x < 0 \Rightarrow f(x) = -x^2$

Then $-x \geq 0 \Rightarrow f(-x) = -(-x)^2 = x^2 = -f(x)$

Hence $f(x)$ is an odd function. An odd function is a function which satisfies the property $f(-x) = -f(x)$, $\forall x \in$ Domain of $f(x)$



There is a property of integration of odd functions which states that

$$\int_{-a}^a f(x) dx = 0 \text{ if } f(x) \text{ is an odd function.}$$

$$\text{Therefore } I = \int_{-1}^1 x|x| dx = 0$$

20. Question

$$\text{Evaluate } \int_a^b \frac{f(x)}{f(x) + f(a+b-x)} dx$$

Answer

$$\text{Let } I = \int_a^b \frac{f(x)}{f(x) + f(a+b-x)} dx - (1)$$

$$\text{Using the property that } \int_a^b f(x) dx = \int_a^b f(a+b-x) dx$$

$$\text{We get } I = \int_a^b \frac{f(a+b-x)}{f(a+b-x) + f(a+b-(a+b-x))} dx$$

$$I = \int_a^b \frac{f(a+b-x)}{f(a+b-x) + f(x)} dx - (2)$$

Adding (1) and (2) we get

$$2I = \int_a^b \frac{f(x) + f(a+b-x)}{f(x) + f(a+b-x)} dx$$

$$= \int_a^b dx$$

$$= b - a$$

$$I = \frac{b - a}{2}$$

21. Question

$$\text{Evaluate } \int_0^1 \frac{1}{1+x^2} dx$$

Answer

$$\text{Let } I = \int_0^1 \frac{1}{1+x^2} dx$$

Substituting $x=\tan\theta \Rightarrow dx=\sec^2\theta d\theta$ (By differentiating both sides)

$$\text{Also, when } x=0, \theta=0 \text{ and } x=1, \theta=\frac{\pi}{4}$$

$$\text{We get } I = \int_0^{\frac{\pi}{4}} \frac{1}{1+\tan^2\theta} \sec^2\theta d\theta$$

$$\text{Since } \sec^2\theta=1+\tan^2\theta$$

$$\text{We get } I = \int_0^{\frac{\pi}{4}} d\theta$$

$$= \frac{\pi}{4}$$

22. Question

Evaluate $\int_0^{\pi/4} \tan x \, dx$

Answer

$$\text{Let } I = \int_0^{\pi/4} \tan x \, dx = \int_0^{\pi/4} \frac{\sin x}{\cos x} \, dx$$

Substituting $\cos x = y \Rightarrow -\sin x \, dx = dy$ (By differentiating both sides)

Also, when $x=0, y=1$ and $x=\frac{\pi}{4}, y=\frac{1}{\sqrt{2}}$

$$\text{We get } I = - \int_1^{\frac{1}{\sqrt{2}}} \frac{1}{y} \, dy$$

$$\text{We get } I = -\log_e y \Big|_1^{\frac{1}{\sqrt{2}}} \text{ (Check Q23. For proof)}$$

$$= -(\log_e \frac{1}{\sqrt{2}}) - \log_e 1$$

Since $\log_a 1 = 0$ and $-\log_a b = \log_a \frac{1}{b}$, We get

$$I = \log_e \sqrt{2}$$

23. Question

Evaluate $\int_2^3 \frac{1}{x} \, dx$

Answer

$$\text{Let } I = \int_2^3 \frac{1}{x} \, dx$$

Substitute $x = e^y \Rightarrow dx = e^y \, dy$ (Differentiating both sides)

Since $x = e^y \Rightarrow y = \log_e x$ and when $x = 2, y = \log_e 2$, and when $x = 3, y = \log_e 3$

$$\text{We get } I = \int_{\log_e 2}^{\log_e 3} \frac{1}{e^y} e^y \, dy$$

$$I = \int_{\log_e 2}^{\log_e 3} dy$$

$$= \log_e 3 - \log_e 2$$

$$= \log_e \frac{3}{2} \text{ (Since } \log_a b - \log_a c = \log_a \frac{b}{c})$$

24. Question

Evaluate $\int_0^2 \sqrt{4-x^2} \, dx$

Answer

$$\text{Let } I = \int_0^2 \sqrt{4-x^2} \, dx$$

Substituting $x = 2\sin\theta \Rightarrow dx = 2\sin\theta d\theta$

Also, When $x=0, \theta=0$ and $x=2, \theta=\frac{\pi}{2}$

We get $I = \int_0^{\frac{\pi}{2}} \sqrt{4 - (2\sin\theta)^2} 2\cos\theta d\theta$

$$I = \int_0^{\frac{\pi}{2}} \sqrt{4 - 4\sin^2\theta} 2\cos\theta d\theta$$

$$= \int_0^{\frac{\pi}{2}} \sqrt{1 - \sin^2\theta} 4\cos\theta d\theta$$

$$= \int_0^{\frac{\pi}{2}} 4\cos^2\theta d\theta$$

Since $\cos 2\theta = 2\cos^2\theta - 1 \Rightarrow \cos^2\theta = \frac{1+\cos 2\theta}{2}$

$$I = \int_0^{\frac{\pi}{2}} (2 + 2\cos 2\theta) d\theta$$

$$= 2\theta \left| \frac{1}{2} + \sin 2\theta \right|_0^{\frac{\pi}{2}}$$

$$= \pi + \sin \pi - \sin 0$$

$$= \pi$$

25. Question

Evaluate $\int_0^1 \frac{2x}{1+x^2} dx$

Answer

$$\text{Let } I = \int_0^1 \frac{2x}{1+x^2} dx$$

Substituting $1+x^2=t$

$$\Rightarrow 2x dx = dt$$

Also, When $x=0, t=1$ and $x=1, t=2$

We get $I = \int_1^2 \frac{1}{t} dt$ (For proof check Q23.)

$$= \log_e t \Big|_1^2 = \log_e 2 - \log_e 1 = \log_e 2 \quad (\text{Since } \log_e 1 = 0)$$

26. Question

Evaluate $\int_0^1 x e^{x^2} dx$

Answer

$$\text{Let } I = \int_0^1 x e^{x^2} dx$$

Substitute $x^2=y$

$$\Rightarrow 2x dx = dy$$

Also, when $x=0, y=0$ and $x=1, y=1$

We get $I = \int_0^1 \frac{e^y}{2} dy$

$$\text{Since } \int e^y dy = e^y$$

We get $I = \frac{e^y}{2} \Big|_0^1 = \frac{e^1 - e^0}{2} = \frac{e-1}{2}$

27. Question

Evaluate $\int_0^{\pi/4} \sin 2x \, dx$

Answer

Let $I = \int_0^{\pi/4} \sin 2x \, dx$

Substitute $2x=y \Rightarrow 2dx=dy$

Also, when $x=0, y=0$ and $x=\frac{\pi}{4}, y=\frac{\pi}{2}$

We get $I = \int_0^{\pi/2} \frac{\sin y}{2} dy$

$$= -\frac{1}{2} \cos y \Big|_0^{\pi/2} = -\frac{1}{2} (\cos(\frac{\pi}{2}) - \cos 0) = \frac{1}{2}$$

28. Question

Evaluate $\int_e^{e^2} \frac{1}{x \log x} dx$

Answer

Let $I = \int_e^{e^2} \frac{1}{x \log_e x} dx$

Substitute $\log_e x = y \Rightarrow \frac{dx}{x} = dy$

Also, When $x=e, y=1$ and $x=e^2, y=2$

We get $I = \int_1^2 \frac{1}{y} dy$ (Check Q23. For proof)

$$= \log_e y \Big|_1^2 = \log_e 2 - \log_e 1 = \log_e 2$$

29. Question

Evaluate $\int_e^{\pi/2} e^x (\sin x - \cos x) dx$

Answer

Let $I = \int_e^{\pi/2} e^x (\sin x - \cos x) dx$

Substitute $-e^x \cos x = t \Rightarrow e^x (-\cos x + \sin x) dx = dt$

(Differentiating both sides by using multiplication rule)

Also, When $x=e, t=-e^e \cos e$ and $x=\frac{\pi}{2}, t=0$

We get $I = \int_{-e^e \cos e}^0 dt = t \Big|_{-e^e \cos e}^0 = 0 - (-e^e \cos e) = e^e \cos e$

30. Question

Evaluate $\int_2^4 \frac{x}{2x^2 + 1} dx$

Answer

$$\text{Let } I = \int_2^4 \frac{x}{1+x^2} dx$$

Substitute $1+x^2=t \Rightarrow 2xdx=dt$

Also, When $x=2, t=5$ and $x=4, t=17$

$$\text{We get } I = \frac{1}{2} \int_5^{17} \frac{1}{t} dt$$

$$= \frac{1}{2} \log_e t \Big|_5^{17} = \frac{1}{2} (\log_e 17 - \log_e 5) = \frac{1}{2} \log_e \frac{17}{5}$$

$$(\text{Since } \log_a b - \log_a c = \log_a \frac{b}{c})$$

31. Question

If $\int_0^1 (3x^2 + 2x + k) dx = 0$, find the value of k.

Answer

To find the value of K, First we have to integrate above integral for which we have to apply simple formulas of integration $\int x^2 dx$ and $\int x dx$, so

$$\int_0^1 (3x^2 + 2x + k) dx = [3(x^3/3) + 2(x^2/2) + k.x]_0^1 = 0$$

Put the upper limit and lower limit in above equation-

$$\Rightarrow \left[3\left(\frac{1}{3}\right) + 2\left(\frac{1}{2}\right) + k.1 \right] - [3.(0) + 2.(0) + k.(0)] = 0$$

$$=(1+1+k) = 0$$

$$K = -2$$

32. Question

If $\int_0^a 3x^2 dx = 8$, write the value of a.

Answer

Doing integration yields-

$$\left[3\frac{x^3}{3} \right]_0^a = 8$$

$$a^3 = 8$$

$$a = \sqrt[3]{8}$$

$$a=2$$

33. Question

If $f(x) = \int_0^x t \sin t dt$, then write the value of $f'(x)$.

Answer

Doing integration yields-

$$f(x) = t \cdot (-\cos t) \Big|_0^x - \int_0^x \frac{d}{dt}(t) \cdot (-\cos t) dt$$

$$= -(x \cos x - 0) + \int_0^x \cos t dt$$

$$= -x \cos x + [\sin t] \Big|_0^x$$

then finally $f(x) = -x \cos x + \sin x$

To calculate derivative of the above function $f(x)$ we have to apply formula of derivation of products of two functions-

$$F(x) = -x \cos(x) + \sin(x)$$

$$F'(x) = -[x(-\sin x) + \cos x \cdot 1] + \cos x ; \{ \text{by formula } d/dx(f \cdot g) = f \cdot g' + g \cdot f' \}$$

$$F'(x) = -(-x \sin x) - \cos x + \cos x$$

$$F'(x) = x \sin(x)$$

34. Question

$$\text{If } \int_0^a \frac{1}{4+x^2} dx = \frac{\pi}{8}, \text{ find the value of } a.$$

Answer

Doing integration yields-

$$\left[\frac{1}{2} \tan^{-1} \left(\frac{x}{2} \right) \right] \Big|_0^a = \frac{\pi}{8}$$

$$\frac{1}{2} \left(\tan^{-1} \frac{a}{2} - \tan^{-1} 0 \right) = \frac{\pi}{8}$$

$$\tan^{-1} \frac{a}{2} - 0 = \frac{\pi}{4}$$

$$a = \tan \frac{\pi}{4} + 1$$

$$a = 2$$

35. Question

Write the coefficient a, b, c of which the value of the integral $\int_{-3}^3 (ax^2 - bx + c) dx$ is independent.

Answer

Doing integration yields-

$$\left(a \cdot \frac{x^3}{3} - b \cdot \frac{x^2}{2} + c \cdot x \right) \Big|_{-3}^3$$

$$\text{By substituting upper and lower limit in above equation} = \left(9a - b \cdot \frac{9}{2} + 3c \right) - \left(9a - b \cdot \frac{9}{2} - 3c \right)$$

$$= 18a + bc$$

so now we can say this is independent of variable b

36. Question

$$\text{Evaluate } \int_2^3 3^x dx$$

Answer

Doing integration yields-

$$\begin{aligned}
 &= \left[\frac{3^x}{\log 3} \right]_2^3 \\
 &= \frac{1}{\log 3} (3^3 - 3^2) \\
 &= \frac{18}{\log 3}
 \end{aligned}$$

37. Question

$$I = \int_0^2 [x] dx$$

Answer

we know that

$$\begin{aligned}
 [x] &= \begin{cases} 0, & 0 < x < 1 \\ 1, & 1 < x < 2 \end{cases} \\
 I &= \int_0^1 [x] dx + \int_1^2 [x] dx = \int_0^1 0 dx + \int_1^2 1 dx \\
 &= 0 + (x)_1^2 \\
 &= (2-1) \\
 &= 1
 \end{aligned}$$

38. Question

$$I = \int_0^{15} [x] dx$$

Answer

we know that

$$[x] = \begin{cases} 0, & 0 < x < 1 \\ 1, & 1 < x < 2 \end{cases}$$

$$I = \int_0^1 [x] dx + \int_1^2 [x] dx + \int_2^3 [x] dx + \dots + \int_{14}^{15} [x] dx$$

$$= (0) + (1) + (2) + \dots + 14$$

$$= 105 \text{ ans}$$

39. Question

$$I = \int_0^1 \{x\} dx$$

Answer

we all know that-

$$\{x\} = x, 0 < x < 1$$

$$I = \int_0^1 x dx = \left[\frac{x^2}{2} \right]_0^1 = \frac{1}{2} \text{ ans}$$

40. Question

$$I = \int_0^1 e^{\{x\}} dx$$

Answer

we all know that-

$$\{x\} = x, 0 < x < 1$$

$$I = \int_0^1 e^x dx = (e^1 - e^0) = (e - 1)$$

41. Question

$$I = \int_0^2 x[x] dx$$

Answer

we know that

$$[x] = \begin{cases} 0, & 0 < x < 1 \\ 1, & 1 < x < 2 \end{cases}$$

$$I = \int_0^1 x[x] dx + \int_1^2 x[x] dx = \int_0^1 0 dx + \int_1^2 x \cdot 1 dx$$

$$= 0 + \left[\frac{x^2}{2} \right]_1^2 = \frac{2^2}{2} - \frac{1^2}{2}$$

$$= \frac{3}{2}$$

42. Question

$$I = \int_0^1 2^{x-[x]} dx$$

Answer

we know that

$$[x] = \begin{cases} 0, & 0 < x < 1 \\ 1, & 1 < x < 2 \end{cases}$$

$$I = \int_0^1 2^{(x-0)} dx = \int_0^1 2^x dx = \left[\frac{2^x}{\log 2} \right]_0^1 = \frac{1}{\log 2} [2^1 - 2^0]$$

$$= \frac{1}{\log 2}$$

43. Question

$$I = \int_0^2 \log_e [x] dx$$

Answer

we know that

$$[x] = \begin{cases} 0, & 0 < x < 1 \\ 1, & 1 < x < 2 \end{cases}$$

$$\begin{aligned} I &= \int_0^1 \log[x] dx + \int_1^2 \log[x] dx = \int_0^1 0 dx + \int_1^2 \log 1. dx \\ &= 0 \end{aligned}$$

44. Question

$$I = \int_0^{\sqrt{2}} [x^2] dx$$

Answer

we know that

$$[x] = \begin{cases} 0, & 0 < x < 1 \\ 1, & 1 < x < 2 \end{cases}$$

$$\begin{aligned} I &= \int_0^1 [x^2] dx + \int_1^{\sqrt{2}} [x^2] dx = \int_0^1 0 dx + \int_1^{\sqrt{2}} 1. dx \\ &= (\sqrt{2} - 1) \end{aligned}$$

45. Question

$$I = \int_0^{\pi/4} \sin\{x\} dx$$

Answer

we all know that-

$$\{x\} = x, 0 < x < 1$$

$$\begin{aligned} I &= \int_0^{\pi/4} \sin(x) dx = (-\cos x) \Big|_0^{\pi/4} = -(\cos \frac{\pi}{4} - \cos 0) \\ &= \left(1 - \frac{1}{\sqrt{2}}\right) \end{aligned}$$

MCQ

1. Question

$$\int_0^1 \sqrt{x(1-x)} dx \text{ equals}$$

- A. $\pi/2$
- B. $\pi/4$
- C. $\pi/6$
- D. $\pi/8$

Answer

Let, $x = \sin^2 t$

Differentiating both sides with respect to t

$$\frac{dx}{dt} = 2 \sin t \cos t$$

$$\Rightarrow dx = 2 \sin t \cos t dt$$

At $x = 0, t = 0$

At $x = 1, t = \frac{\pi}{2}$

$$y = \int_0^{\frac{\pi}{2}} \sqrt{\sin^2 t (1 - \sin^2 t)} \times 2 \sin t \cos t dt$$

$$= \int_0^{\frac{\pi}{2}} 2 \sin^2 t \cos^2 t dt$$

$$= \frac{1}{2} \int_0^{\frac{\pi}{2}} \sin^2 2t dt$$

$$= \frac{1}{2} \int_0^{\frac{\pi}{2}} \frac{1 - \cos 2(2t)}{2} dt$$

$$= \frac{1}{2} \left(\frac{t}{2} - \frac{\sin 4t}{8} \right)_0^{\frac{\pi}{2}}$$

$$= \frac{1}{2} \left[\left(\frac{\pi}{4} - 0 \right) - (0 - 0) \right]$$

$$y = \frac{\pi}{8}$$

2. Question

$$\int_0^{\pi} \frac{1}{1 + \sin x} dx$$
 equals

A. 0

B. 1/2

C. 2

D. 3/2

Answer

$$y = \int_0^{\pi} \frac{1}{1 + \sin x} dx$$

Multiply by $1 - \sin x$ in numerator and denominator

$$y = \int_0^{\pi} \frac{1 - \sin x}{(1 + \sin x)(1 - \sin x)} dx$$

$$= \int_0^{\pi} \frac{1 - \sin x}{1 - \sin^2 x} dx$$

$$= \int_0^{\pi} \frac{1 - \sin x}{\cos^2 x} dx$$

$$= \int_0^\pi \frac{1}{\cos^2 x} - \frac{\sin x}{\cos^2 x} dx$$

$$= \int_0^\pi \sec^2 x - \tan x \sec x dx$$

$$= (\tan x - \sec x)_0^\pi$$

$$y = [(0 - (-1)) - (0 - 1)]$$

$$y = 2$$

3. Question

$$\int_0^\pi \frac{x \tan x}{\sec x + \cos x} dx$$

A. $\frac{\pi^2}{4}$

B. $\frac{\pi^2}{2}$

C. $\frac{3\pi^2}{2}$

D. $\frac{\pi^2}{3}$

Answer

In this question we can use the king rule

$$y = \int_0^\pi \frac{x \tan x}{\sec x + \cos x} dx \dots (1)$$

$$= \int_0^\pi \frac{(\pi - x) \tan(\pi - x)}{\sec(\pi - x) + \cos(\pi - x)} dx$$

$$= \int_0^\pi \frac{-\pi \tan x}{-(\sec x + \cos x)} - \frac{-x \tan x}{-(\sec x + \cos x)} dx \dots (2)$$

On adding eq(1) and eq(2)

$$2y = \int_0^\pi \frac{\pi \tan x}{\sec x + \cos x} dx$$

$$= \int_0^\pi \frac{\pi \sin x}{1 + \cos^2 x} dx$$

Let, $\cos x = t$

Differentiating both side with respect to x

$$\frac{dt}{dx} = -\sin x \Rightarrow -dt = \sin x dx$$

At $x = 0, t = 1$

At $x = \pi, t = -1$

$$2y = \int_1^{-1} \frac{-\pi}{1 + t^2} dt$$

$$= -\pi(\tan^{-1} t)_1^{-1}$$

$$y = \frac{-\pi}{2} \left[-\frac{\pi}{4} - \frac{\pi}{4} \right]$$

$$= \frac{\pi^2}{4}$$

4. Question

The value of $\int_0^{2\pi} \sqrt{1 + \sin \frac{x}{2}} dx$ is

- A. 0
- B. 2
- C. 8
- D. 4

Answer

$$y = \int_0^{2\pi} \sqrt{\sin^2 \frac{x}{4} + \cos^2 \frac{x}{4} + 2 \sin \frac{x}{4} \cos \frac{x}{4}} dx$$

$$= \int_0^{2\pi} \sin \frac{x}{4} + \cos \frac{x}{4} dx$$

$$= 4 \left(-\cos \frac{x}{4} + \sin \frac{x}{4} \right)_0^{2\pi}$$

$$= 4[(0 + 1) - (-1 - 0)]$$

$$y = 8$$

5. Question

The value of the integral $\int_0^{\frac{\pi}{2}} \frac{\sqrt{\cos x}}{\sqrt{\cos x} + \sqrt{\sin x}} dx$ is

- A. 0
- B. $\pi/2$
- C. $\pi/4$
- D. none of these

Answer

Mistake: limit should be 0 to $\pi/2$

Right sol. In this question we apply the king rule

$$y = \int_0^{\frac{\pi}{2}} \frac{\sqrt{\cos x}}{\sqrt{\cos x} + \sqrt{\sin x}} dx \dots (1)$$

$$= \int_0^{\frac{\pi}{2}} \frac{\sqrt{\cos(\frac{\pi}{2} - x)}}{\sqrt{\cos(\frac{\pi}{2} - x)} + \sqrt{\sin(\frac{\pi}{2} - x)}} dx$$

$$= \int_0^{\frac{\pi}{2}} \frac{\sqrt{\sin x}}{\sqrt{\sin x} + \sqrt{\cos x}} dx \dots (2)$$

On adding eq(1) and eq(2)

$$2y = \int_0^{\frac{\pi}{2}} \frac{\sqrt{\sin x} + \sqrt{\cos x}}{\sqrt{\sin x} + \sqrt{\cos x}} dx$$

$$= \int_0^{\frac{\pi}{2}} dx$$

$$y = \frac{1}{2} (x) \Big|_0^{\frac{\pi}{2}}$$

$$y = \frac{\pi}{4}$$

6. Question

$$\int_0^{\infty} \frac{1}{1+e^x} dx \text{ equals}$$

- A. $\log 2 - 1$
- B. $\log 2$
- C. $\log 4 - 1$
- D. $-\log 2$

Answer

Take e^x out from the denominator

$$y = \int_0^{\infty} \frac{1}{e^x(e^{-x}+1)} dx$$

$$y = \int_0^{\infty} \frac{e^{-x}}{e^{-x}+1} dx$$

Let, $e^{-x} + 1 = t$

Differentiating both side with respect to t

$$\frac{dt}{dx} = -e^{-x} \Rightarrow -dt = e^{-x} dx$$

At $x = 0, t = 2$

At $x = \infty, t = 1$

$$y = - \int_2^1 \frac{1}{t} dt$$

$$y = -(\log t) \Big|_2^1$$

$$y = -(0 - \log 2)$$

$$y = \log 2$$

7. Question

$$\int_0^{\pi/4} \frac{\sin \sqrt{x}}{\sqrt{x}} dx \text{ equals}$$

- A. 2
- B. 1
- C. $\pi/4$

D. $\pi^2/8$

Answer

Let, $\sqrt{x} = t$

Differentiating both side with respect to x

$$\frac{dt}{dx} = \frac{1}{2\sqrt{x}} \Rightarrow 2dt = \frac{1}{\sqrt{x}} dx$$

At $x = 0, t = 0$

At $x = \pi^2/4, t = \pi/2$

$$y = 2 \int_0^{\frac{\pi}{2}} \sin t \ dt$$

$$y = 2(-\cos t) \Big|_0^{\frac{\pi}{2}}$$

$$y = 2[0 - (-1)]$$

$$y = 2$$

8. Question

$$\int_0^{\pi/2} \frac{\cos x}{(2 + \sin x)(1 + \sin x)} dx \text{ equals}$$

A. $\log\left(\frac{2}{3}\right)$

B. $\log\left(\frac{3}{2}\right)$

C. $\log\left(\frac{3}{4}\right)$

D. $\log\left(\frac{4}{3}\right)$

Answer

Let, $\sin x = t$

Differentiating both side with respect to x

$$\frac{dt}{dx} = \cos x \Rightarrow dt = \cos x dx$$

At $x = 0, t = 0$

At $x = \pi/2, t = 1$

$$y = \int_0^1 \frac{1}{(2+t)(1+t)} dt$$

By using the concept of partial fraction

$$\frac{1}{(2+t)(1+t)} = \frac{A}{(2+t)} + \frac{B}{(1+t)}$$

$$1 = A(1+t) + B(2+t)$$

$$1 = (A + 2B) + t(A + B)$$

$$A + 2B = 1, A + B = 0$$

$$A = -1, B = 1$$

$$y = \int_0^1 \frac{-1}{(2+t)} + \frac{1}{(1+t)} dt$$

$$y = [-\log(2+t) + \log(1+t)]_0^1$$

$$y = [(-\log 3 + \log 2) - (-\log 2 + \log 1)]$$

$$y = \log\left(\frac{4}{3}\right)$$

9. Question

$$\int_0^{\pi/2} \frac{1}{2+\cos x} dx \text{ equals}$$

A. $\frac{1}{3} \tan^{-1}\left(\frac{1}{\sqrt{3}}\right)$

B. $\frac{2}{\sqrt{3}} \tan^{-1}\left(\frac{1}{\sqrt{3}}\right)$

C. $\sqrt{3} \tan^{-1}(\sqrt{3})$

D. $2\sqrt{3} \tan^{-1}\sqrt{3}$

Answer

$$y = \int_0^{\frac{\pi}{2}} \frac{1}{2 + \frac{1 - \tan^2 \frac{x}{2}}{1 + \tan^2 \frac{x}{2}}} dx$$

$$y = \int_0^{\frac{\pi}{2}} \frac{1 + \tan^2 \frac{x}{2}}{2 \left(1 + \tan^2 \frac{x}{2}\right) + 1 - \tan^2 \frac{x}{2}} dx$$

$$y = \int_0^{\frac{\pi}{2}} \frac{\sec^2 \frac{x}{2}}{3 + \tan^2 \frac{x}{2}} dx$$

Let, $\tan \frac{x}{2} = t$

Differentiating both side with respect to x

$$\frac{dt}{dx} = \frac{1}{2} \sec^2 \frac{x}{2} \Rightarrow 2 dt = \sec^2 \frac{x}{2} dx$$

At $x = 0, t = 0$

At $x = \pi/2, t = 1$

$$y = \int_0^1 \frac{2}{(\sqrt{3})^2 + t^2} dt$$

$$y = 2 \left(\frac{1}{\sqrt{3}} \tan^{-1} \frac{t}{\sqrt{3}} \right)_0^1$$

$$y = \frac{2}{\sqrt{3}} \left(\tan^{-1} \frac{1}{\sqrt{3}} - 0 \right)$$

$$y = \frac{2}{\sqrt{3}} \tan^{-1} \left(\frac{1}{\sqrt{3}} \right)$$

10. Question

$$\int_0^1 \sqrt{\frac{1-x}{1+x}} dx =$$

A. $\frac{\pi}{2}$

B. $\frac{\pi}{2} - 1$

C. $\frac{\pi}{2} + 1$

D. $\pi + 1$

Answer

Right sol. Let, $x = \sin t$

Differentiating both side with respect to t

$$\frac{dx}{dt} = \cos t \Rightarrow dx = \cos t dt$$

At $x = 0, t = 0$

At $x = 1, t = \pi/2$

$$y = \int_0^{\frac{\pi}{2}} \sqrt{\frac{1-\sin t}{1+\sin t}} \cos t dt$$

Multiply by $1 - \sin t$ in numerator and denominator

$$y = \int_0^{\frac{\pi}{2}} \sqrt{\frac{(1-\sin t)(1-\sin t)}{(1+\sin t)(1-\sin t)}} \cos t dt$$

$$= \int_0^{\frac{\pi}{2}} \frac{1-\sin t}{\cos t} \cos t dt$$

$$= \int_0^{\frac{\pi}{2}} (1-\sin t) dt$$

$$= (t + \cos t) \Big|_0^{\frac{\pi}{2}}$$

$$= \left[\left(\frac{\pi}{2} + 0 \right) - (0 + 1) \right]$$

$$y = \frac{\pi}{2} - 1$$

11. Question

$$\int_0^{\pi} \frac{1}{a+b \cos x} dx =$$

A. $\frac{\pi}{\sqrt{a^2 - b^2}}$

B. $\frac{\pi}{ab}$

C. $\frac{\pi}{a^2 + b^2}$

D. $(a+b)\pi$

Answer

$$y = \int_0^\pi \frac{1}{a+b \frac{1-\tan^2 \frac{x}{2}}{1+\tan^2 \frac{x}{2}}} dx$$

$$y = \int_0^\pi \frac{1 + \tan^2 \frac{x}{2}}{a(1 + \tan^2 \frac{x}{2}) + b(1 - \tan^2 \frac{x}{2})} dx$$

$$y = \int_0^\pi \frac{\sec^2 \frac{x}{2}}{(a+b) + (a-b)\tan^2 \frac{x}{2}} dx$$

Let, $\tan \frac{x}{2} = t$

Differentiating both side with respect to x

$$\frac{dt}{dx} = \frac{1}{2} \sec^2 \frac{x}{2} \Rightarrow 2 dt = \sec^2 \frac{x}{2} dx$$

At $x = 0, t = 0$

At $x = \pi, t = \infty$

$$y = \int_0^\infty \frac{2}{(\sqrt{a+b})^2 + (t\sqrt{a-b})^2} dt$$

$$y = \frac{2}{\sqrt{a-b}} \left(\frac{1}{\sqrt{a+b}} \tan^{-1} \frac{t\sqrt{a-b}}{\sqrt{a+b}} \right)_0^\infty$$

$$y = \frac{2}{\sqrt{a^2 - b^2}} \left(\frac{\pi}{2} - 0 \right)$$

$$y = \frac{\pi}{\sqrt{a^2 - b^2}}$$

12. Question

$$\int_{\pi/6}^{\pi/3} \frac{1}{1 + \sqrt{\cot x}} dx$$
 is

A. $\pi/3$

B. $\pi/6$

C. $\pi/12$

D. $\pi/2$

Answer

$$y = \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{\sqrt{\sin x}}{\sqrt{\cos x} + \sqrt{\sin x}} dx \dots(1)$$

$$y = \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{\sqrt{\sin\left(\frac{\pi}{3} + \frac{\pi}{6} - x\right)}}{\sqrt{\cos\left(\frac{\pi}{3} + \frac{\pi}{6} - x\right)} + \sqrt{\sin\left(\frac{\pi}{3} + \frac{\pi}{6} - x\right)}} dx$$

$$y = \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{\sqrt{\cos x}}{\sqrt{\sin x} + \sqrt{\cos x}} dx \dots(2)$$

On adding eq(1) and eq(2)

$$2y = \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{\sqrt{\sin x} + \sqrt{\cos x}}{\sqrt{\sin x} + \sqrt{\cos x}} dx$$

$$2y = \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} dx$$

$$y = \frac{1}{2} (x) \Big|_{\frac{\pi}{6}}^{\frac{\pi}{3}}$$

$$y = \frac{1}{2} \left(\frac{\pi}{3} - \frac{\pi}{6} \right)$$

$$y = \frac{\pi}{12}$$

13. Question

Given that $\int_0^{\infty} \frac{x^2}{(x^2 + a^2)(x^2 + b^2)(x^2 + c^2)} dx = \frac{\pi}{2(a+b)(b+c)(c+a)}$, the value of $\int_0^{\infty} \frac{dx}{(x^2 + 4)(x^2 + 9)}$, is

A. $\frac{\pi}{60}$

B. $\frac{\pi}{20}$

C. $\frac{\pi}{40}$

D. $\frac{\pi}{80}$

Answer

In this question we use the method of partial fraction

$$\frac{1}{(x^2 + 4)(x^2 + 9)} = \frac{A}{(x^2 + 4)} + \frac{B}{(x^2 + 9)}$$

$$1 = A(x^2 + 9) + B(x^2 + 4)$$

$$A + B = 0, 9A + 4B = 1$$

$$A = 1/5, B = -1/5$$

$$y = \int_0^{\infty} \frac{\frac{1}{5}}{(x^2 + 4)} + \frac{\frac{-1}{5}}{(x^2 + 9)} dx$$

$$y = \left(\frac{1}{5 \times 2} \tan^{-1} \frac{x}{2} - \frac{1}{5 \times 3} \tan^{-1} \frac{x}{3} \right)_0^{\infty}$$

$$y = \left(\frac{\pi}{20} - \frac{\pi}{30} \right)$$

$$y = \frac{\pi}{60}$$

14. Question

$$\int_1^e \log x \, dx =$$

- A. 1
- B. $e - 1$
- C. $e + 1$
- D. 0

Answer

$$y = \int_1^e 1 \times \log x \, dx$$

By using integration by parts

Let, $\log x$ as 1st function and 1 as 2nd function

Use formula $\int I \times II \, dx = I \int II \, dx - \int \left(\frac{d}{dx} I \right) (\int II \, dx) \, dx$

$$y = \log x \int_1^e dx - \int_1^e \left(\frac{d}{dx} \log x \right) (\int_1^e dx) \, dx$$

$$y = (\log x)x - \int_1^e \left(\frac{1}{x} \right) (x) \, dx$$

$$y = x \log x - \int_1^e dx$$

$$y = (x \log x - x)_1^e$$

$$y = [(e - e) - (0 - 1)]$$

$$y = 1$$

15. Question

$$\int_1^{\sqrt{3}} \frac{1}{1+x^2} \, dx$$
 is equal to

$$A. \frac{\pi}{12}$$

$$B. \frac{\pi}{6}$$

C. $\frac{\pi}{4}$

D. $\frac{\pi}{3}$

Answer

$$y = \int_1^{\sqrt{3}} \frac{1}{1+x^2} dx$$

$$y = (\tan^{-1} x)_1^{\sqrt{3}}$$

$$y = \left(\frac{\pi}{3} - \frac{\pi}{4} \right)$$

$$y = \frac{\pi}{12}$$

16. Question

$$\int_0^3 \frac{3x+1}{x^2+9} dx =$$

A. $\frac{\pi}{12} + \log(2\sqrt{2})$

B. $\frac{\pi}{2} + \log(2\sqrt{2})$

C. $\frac{\pi}{6} + \log(2\sqrt{2})$

D. $\frac{\pi}{3} + \log(2\sqrt{2})$

Answer

$$y = \int_0^3 \frac{3x}{x^2+9} + \frac{1}{x^2+9} dx$$

$$A = \frac{3}{2} \int_0^3 \frac{2x}{x^2+9} dx$$

Let, $x^2 + 9 = t$

Differentiating both side with respect to x

$$\frac{dt}{dx} = 2x \Rightarrow dt = 2x dx$$

At $x = 0, t = 9$

At $x = 3, t = 18$

$$A = \frac{3}{2} \int_9^{18} \frac{1}{t} dt$$

$$= \frac{3}{2} (\log t)_9^{18}$$

$$= \frac{3}{2} (\log 18 - \log 9)$$

$$= \frac{3}{2} \log 2 = \log 2\sqrt{2}$$

$$B = \int_0^3 \frac{1}{x^2 + 9} dx$$

$$= \left(\frac{1}{3} \tan^{-1} \frac{x}{3} \right)_0^3$$

$$= \frac{1}{3} \left(\frac{\pi}{4} - 0 \right)$$

$$= \frac{\pi}{12}$$

So, the complete solution is $y = A + B$

$$y = \frac{\pi}{12} + \log 2\sqrt{2}$$

17. Question

The value of the integral $\int_0^\infty \frac{x}{(1+x)(1+x^2)} dx$ is

A. $\frac{\pi}{2}$

B. $\frac{\pi}{4}$

C. $\frac{\pi}{6}$

D. $\frac{\pi}{3}$

Answer

Let, $x = \tan t$

Differentiating both side with respect to t

$$\frac{dx}{dt} = \sec^2 t \Rightarrow dx = \sec^2 t dt$$

At $x = 0, t = 0$

At $x = \infty, t = \pi/2$

$$y = \int_0^{\pi/2} \frac{\tan t}{(1+\tan t)(1+\tan^2 t)} \sec^2 t dt$$

$$y = \int_0^{\pi/2} \frac{\tan t}{(1+\tan t)} dt$$

$$y = \int_0^{\pi/2} \frac{\sin t}{(\cos t + \sin t)} dt \dots (1)$$

By using the king rule

$$y = \int_0^{\pi/2} \frac{\sin(\frac{\pi}{2} - t)}{\left(\cos\left(\frac{\pi}{2} - t\right) + \sin\left(\frac{\pi}{2} - t\right) \right)} dt$$

$$y = \int_0^{\frac{\pi}{2}} \frac{\cos t}{(\cos t + \sin t)} dt \dots(2)$$

On adding eq(1) and eq(2)

$$2y = \int_0^{\frac{\pi}{2}} \frac{\sin t + \cos t}{(\cos t + \sin t)} dt$$

$$2y = \int_0^{\frac{\pi}{2}} dt$$

$$y = \frac{1}{2} (t) \Big|_0^{\frac{\pi}{2}}$$

$$y = \frac{\pi}{4}$$

18. Question

$$\int_{-\pi/2}^{\pi/2} \sin |x| dx \text{ is equal to}$$

- A. 1
- B. 2
- C. -1
- D. -2

Answer

In this question, we break the limit in two-part

$$y = \int_{-\frac{\pi}{2}}^0 \sin(-x) dx + \int_0^{\frac{\pi}{2}} \sin x dx$$

$$y = (\cos x) \Big|_{-\frac{\pi}{2}}^0 + (-\cos x) \Big|_0^{\frac{\pi}{2}}$$

$$y = (1 - 0) + [0 - (-1)]$$

$$y = 2$$

19. Question

$$\int_0^{\pi/2} \frac{1}{1 + \tan x} dx \text{ is equal to}$$

- A. $\frac{\pi}{4}$
- B. $\frac{\pi}{3}$
- C. $\frac{\pi}{2}$
- D. π

Answer

$$y = \int_0^{\frac{\pi}{2}} \frac{\cos x}{(\cos x + \sin x)} dx \dots(1)$$

By using the king rule

$$y = \int_0^{\frac{\pi}{2}} \frac{\cos\left(\frac{\pi}{2} - x\right)}{\left(\cos\left(\frac{\pi}{2} - x\right) + \sin\left(\frac{\pi}{2} - x\right)\right)} dx$$

$$y = \int_0^{\frac{\pi}{2}} \frac{\sin x}{(\cos x + \sin x)} dx \dots(2)$$

On adding eq(1) and eq(2)

$$2y = \int_0^{\frac{\pi}{2}} \frac{\sin x + \cos x}{(\cos x + \sin x)} dx$$

$$2y = \int_0^{\frac{\pi}{2}} dx$$

$$y = \frac{1}{2} (x) \Big|_0^{\frac{\pi}{2}}$$

$$y = \frac{\pi}{4}$$

20. Question

The value of $\int_0^{\pi/2} \cos x e^{\sin x} dx$ is

- A. 1
- B. $e - 1$
- C. 0
- D. -1

Answer

Let, $\sin x = t$

Differentiating both sides with respect to x

$$\frac{dt}{dx} = \cos x \Rightarrow dt = \cos x dx$$

At $x = 0, t = 0$

At $x = \pi/2, t = 1$

$$y = \int_0^1 e^t dt$$

$$y = (e^t) \Big|_0^1$$

$$y = e^1 - e^0$$

$$y = e - 1$$

21. Question

If $\int_0^a \frac{1}{1+4x^2} dx = \frac{\pi}{4}$ then a equals

$$A. \frac{\pi}{2}$$

B. $\frac{1}{2}$

C. $\frac{\pi}{4}$

D. 1

Answer

Given, $\int_0^a \frac{1}{1+4x^2} dx = \frac{\pi}{4}$

$$= \int_0^a \frac{1}{(1)^2 + (2x)^2} dx = \frac{\pi}{4}$$

$$= \left[\frac{1}{1} \tan^{-1} \left(\frac{2x}{1} \right) \right]_0^a + c = \frac{\pi}{4}$$

$$= [\tan^{-1}(2a) - \tan^{-1}(0)] = \frac{\pi}{4}$$

$$= [\tan^{-1}(2a) - 0] = \frac{\pi}{4}$$

$$= \tan^{-1}(2a) = \frac{\pi}{4}$$

$$= 2a = \tan \frac{\pi}{4}$$

$$= 2a = 1$$

$$= a = \frac{1}{2}$$

Option A: it's not option A, because clearly we got the value of 'a' as $\frac{1}{2}$ after solving.

Option C: it's not option C, because clearly we got the value of 'a' as $\frac{1}{2}$ after solving.

Option D: it's not option D, because clearly we got the value of 'a' as $\frac{1}{2}$ after solving.

22. Question

If $\int_0^1 f(x) dx = 1$, $\int_0^1 xf(x) dx = a$, $\int_0^1 x^2 f(x) dx = a^2$, then $\int_0^1 (a-x)^2 f(x) dx$ equals

A. $4a^2$

B. 0

C. $2a^2$

D. none of these

Answer

Given, $\int_0^1 f(x) dx = 1$, $\int_0^1 xf(x) dx = a$, $\int_0^1 x^2 f(x) dx = a^2$,

Now, $\int_0^1 (a-x)^2 f(x) dx$

$$\begin{aligned}
&= \int_0^1 (a^2 - 2ax + x^2)f(x)dx \\
&= \int_0^1 a^2 f(x) dx - \int_0^1 2axf(x) dx + \int_0^1 x^2 f(x)dx \\
&= a^2 \int_0^1 f(x)dx - 2a \int_0^1 x f(x)dx + \int_0^1 x^2 f(x)dx \\
&= a^2(1) - 2a(a) + a^2 \\
&= 2a^2 - 2a^2 \\
&= 0
\end{aligned}$$

Option A:- it's not option A , this is clearly justified on solving.

Option C: - it's not option C , this is clearly justified on solving.

Option D: - it's not option D, this is clearly justified on solving.

23. Question

The value of $\int_{-\pi}^{\pi} \sin^3 x \cos^2 x dx$ is

A. $\frac{\pi^4}{2}$

B. $\frac{\pi^4}{4}$

C. 0

D. none of these

Answer

let, $I_{3,2} = \int_{-\pi}^{\pi} \sin^3 x \cos^2 x dx$

$$\begin{aligned}
[\text{if, } I_{m,n} = \int \sin^m x \cos^n x dx \text{ then } I_{m,n} &= -\frac{\sin^{m-1} x \cos^{n+1} x}{m+n} + \frac{m-1}{m+n} I_{m-2,n}] \\
&= \frac{-\sin^2 x \cos^3 x}{5} + \frac{2}{5} I_{1,2} \\
&= \frac{-\sin^2 x \cos^3 x}{5} + \frac{2}{5} \left[\frac{-\sin^0 x \cos^3 x}{3} + 0 \right] \\
&= \left[\frac{-\sin^2 x \cos^3 x}{5} - \frac{2}{15} \cos^3 x \right]_{-\pi}^{\pi} \\
&= \left[\frac{-\sin^2 \pi \cos^2 \pi}{5} - \frac{2}{15} \cos^3 \pi \right] - \left[\frac{-\sin^2(-\pi) \cos^2(-\pi)}{5} - \frac{2}{15} \cos^3(-\pi) \right] \\
&= 0 - \frac{2}{15}(-1) - 0 + \frac{2}{15}(-1) \\
&= 0
\end{aligned}$$

Option A:- it's not option A , this is clearly justified on solving.



Option B: - it's not option B , this is clearly justified on solving.

Option D: - it's not option D, this is clearly justified on solving.

24. Question

$$\int_{\pi/6}^{\pi/3} \frac{1}{\sin 2x} dx \text{ is equal to}$$

- A. $\log_e 3$
- B. $\log_e \sqrt{3}$
- C. $\frac{1}{2} \log(-1)$
- D. $\log(-1)$

Answer

$$\text{Given, } \int_{\pi/6}^{\pi/3} \frac{1}{\sin 2x} dx$$

$$\begin{aligned}&= \int_{\pi/6}^{\pi/3} \csc 2x dx \\&= \left[\frac{\log |\csc 2x - \cot 2x|}{2} \right]_{\pi/6}^{\pi/3} \\&= \frac{1}{2} \left[\log \left| \csc \frac{2\pi}{3} - \cot \frac{2\pi}{3} \right| \right] - \frac{1}{2} \left[\log \left| \csc \frac{\pi}{3} - \cot \frac{\pi}{3} \right| \right] \\&= \frac{1}{2} \log \left[\frac{2}{\sqrt{3}} + \frac{1}{\sqrt{3}} \right] - \frac{1}{2} \log \left[\frac{2}{\sqrt{3}} - \frac{1}{\sqrt{3}} \right] = \\&= \frac{1}{2} \log \left[\frac{3}{\sqrt{3}} \right] - \frac{1}{2} \log \left[\frac{1}{\sqrt{3}} \right] \\&= \frac{1}{2} \log[\sqrt{3}] - \frac{1}{2} \log[\sqrt{3}]^{-1} \\&= \frac{1}{2} \log[\sqrt{3}] + \frac{1}{2} \log[\sqrt{3}] \\&= \log[\sqrt{3}]\end{aligned}$$

Option A:- it's not option A , this is clearly justified on solving.

Option C: - it's not option C , this is clearly justified on solving.

Option D: - it's not option D, this is clearly justified on solving.

25. Question

$$\int_{-1}^1 |1-x| dx \text{ is equal to}$$

- A. -2
- B. 2
- C. 0
- D. 4

Answer

Given, $\int_{-1}^1 |1-x| dx$

$$\text{Now, } |1-x| = \begin{cases} (1-x) & \text{if } x > 1 \\ (x-1) & \text{if } x < 1 \end{cases}$$

$$\begin{aligned} &= \int_{-1}^0 (x-1) dx + \int_0^1 (x-1) dx \\ &= \left[x - \frac{x^2}{2} \right]_{-1}^0 + \left[x - \frac{x^2}{2} \right]_0^1 \\ &= \left[(0-0) - \left((-1) - \frac{1}{2} \right) \right] + \left[\left(1 - \frac{1}{2} \right) - (0-0) \right] \\ &= 0 + 1 + \frac{1}{2} + 1 - \frac{1}{2} \\ &= 2 \end{aligned}$$

Option A:- it's not option A , this is clearly justified on solving.

Option C:- it's not option C , this is clearly justified on solving.

Option D:- it's not option D, this is clearly justified on solving.

26. Question

The derivative of $f(x) = \int_{x^2}^{x^3} \frac{1}{\log_e t} dt$, ($x > 0$), is

A. $\frac{1}{3 \ln x}$

B. $\frac{1}{3 \ln x} - \frac{1}{2 \ln x}$

C. $(\ln x)^{-1} \times (x-1)$

D. $\frac{3x^2}{\ln x}$

Answer

$$f'(x) = \frac{1}{\log_e x^3} \frac{d}{dx} x^3 - \frac{1}{\log_e x^2} \frac{d}{dx} x^2$$

$$f'(x) = \frac{3x^2}{3 \log_e x} - \frac{2x}{2 \log_e x}$$

$$f'(x) = \frac{x^2 - x}{\log_e x}$$

$$f'(x) = \frac{x(x-1)}{\log_e x}$$

$$f'(x) = (\ln x)^{-1} x (x-1)$$

27. Question

If $I_{10} = \int_0^{\pi/2} x^{10} \sin x dx$, then the value of $I_{10} + 90I_8$ is

A. $9\left(\frac{\pi}{2}\right)^9$

B. $10\left(\frac{\pi}{2}\right)^9$

C. $\left(\frac{\pi}{2}\right)^9$

D. $9\left(\frac{\pi}{2}\right)^8$

Answer

Use method of integration by parts

$$I_{10} = x^{10} \int \sin x dx - \int_0^{\pi/2} \frac{d}{dx} x^{10} \left(\int \sin x dx \right) dx$$

$$I_{10} = (x^{10}(-\cos x))_0^{\pi/2} - \int_0^{\pi/2} 10x^9(-\cos x) dx$$

$$I_{10} = \left(\frac{\pi}{2}\right)^{10} \times (-0) - 0 \times (-1) - \int_0^{\pi/2} 10x^9(-\cos x) dx$$

$$I_{10} = 0 + 10 \left(x^9 \int \cos x dx - \int_0^{\pi/2} \frac{d}{dx} x^9 \left(\int \cos x dx \right) dx \right)$$

$$I_{10} = 10 \left((x^9 \sin x)_0^{\pi/2} - \int_0^{\pi/2} 9x^8 \sin x dx \right)$$

$$I_{10} = 10 \left(\frac{\pi}{2} \right)^9 - 90I_8$$

$$I_{10} + 90I_8 = 10 \left(\frac{\pi}{2} \right)^9$$

28. Question

$$\int_0^1 \frac{x}{(1-x)^{54}} dx =$$

A. $\frac{15}{16}$

B. $\frac{3}{16}$

C. $-\frac{3}{16}$

D. $-\frac{16}{3}$

Answer

We know that $y = \int_a^b f(x)dx = \int_a^b f(a+b-x)dx$

$$y = \int_0^1 \frac{x}{(1-x)^{54}} dx = \int_0^1 \frac{1+0-x}{(1-(1+0-x))^{54}} dx$$

$$y = \int_0^1 \frac{1-x}{x^{54}} dx$$

$$y = \int_0^1 \frac{1}{x^{54}} - \frac{1}{x^{53}} dx$$

$$y = \left(\frac{1}{-53x^{53}} - \frac{1}{-52x^{52}} \right)_0^1$$

29. Question

$\lim_{n \rightarrow \infty} \left\{ \frac{1}{2n+1} + \frac{1}{2n+2} + \dots + \frac{1}{2n+n} \right\}$ is equal to

A. $\ln\left(\frac{1}{3}\right)$

B. $\ln\left(\frac{2}{3}\right)$

C. $\ln\left(\frac{3}{2}\right)$

D. $\ln\left(\frac{4}{2}\right)$

Answer

$$\text{Given, } \lim_{n \rightarrow \infty} \left\{ \frac{1}{2n+1} + \frac{1}{2n+2} + \dots + \frac{1}{2n+n} \right\}$$

$$= \lim_{n \rightarrow \infty} \left\{ \frac{1}{n\left(2+\frac{1}{n}\right)} + \frac{1}{n\left(2+\frac{2}{n}\right)} + \dots + \frac{1}{n\left(2+\frac{n}{n}\right)} \right\}$$

$$= \lim_{n \rightarrow \infty} \frac{1}{n} \left\{ \frac{1}{2+\frac{1}{n}} + \frac{1}{2+\frac{2}{n}} + \dots + \frac{1}{2+\frac{n}{n}} \right\}$$

$$= \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{r=1}^n \frac{1}{2+\frac{r}{n}}$$

Now for easy solvation, replace $\sum_{r=1}^n$ with \int_0^1 , $\frac{r}{n}$ with x and $\frac{1}{n}$ with dx

$$= \int_0^1 \frac{1}{2+x} dx$$

$$= [\log(2+x)]_0^1$$

$$= \log(2+1) - \log(2+0)$$

$$= \log 3 - \log 2$$

$$= \log\left(\frac{3}{2}\right)$$

$$= \ln\left(\frac{3}{2}\right)$$

Option A:- it's not option A , this is clearly justified on solving.

Option B:- it's not option B, this is clearly justified on solving.

Option D:- it's not option D, this is clearly justified on solving.

30. Question

The value of the integral $\int_{-2}^2 |1-x^2| dx$ is

A. 4

B. 2

C. -2

D. 0

Answer

$$\text{Given, } \int_{-2}^2 |1-x^2| dx$$

$$\text{Now, } |1-x^2| = \begin{cases} -(1-x^2) & \text{if } x < 0 \\ 1-x^2 & \text{if } x > 0 \end{cases}$$

$$= \int_{-2}^0 |1-x^2| dx + \int_0^2 |1-x^2| dx$$

$$= \int_{-2}^0 -(1-x^2) dx + \int_0^2 (1-x^2) dx$$

$$= \left[-\left(x - \frac{x^3}{3} \right) \right]_{-2}^0 + \left[x - \frac{x^3}{3} \right]_0^2$$

$$= \left[(0-0) - \left(\frac{(-2)^3}{3} - (-2) \right) \right] + \left[\left(2 - \frac{(2)^3}{3} \right) - (0-0) \right]$$

$$= \frac{8}{3} - 2 + 2 - \frac{8}{3}$$

$$= 0$$

Option A: - it's not option A , this is clearly justified on solving.

Option B: - it's not option B , this is clearly justified on solving.

Option C: - it's not option C, this is clearly justified on solving.

31. Question

$\int_0^{\pi/2} \frac{1}{1+\cot^3 x} dx$ is equal to

- A. 0
- B. 1
- C. $\pi/2$
- D. $\pi/4$

Answer

$$y = \int_0^{\frac{\pi}{2}} \frac{\sin^3 x}{\sin^3 x + \cos^3 x} dx \dots(1)$$

Use king's property $y = \int_a^b f(x)dx = \int_a^b f(a+b-x)dx$

$$y = \int_0^{\frac{\pi}{2}} \frac{\sin^3 \left(\frac{\pi}{2} - x\right)}{\sin^3 \left(\frac{\pi}{2} - x\right) + \cos^3 \left(\frac{\pi}{2} - x\right)} dx$$

$$y = \int_0^{\frac{\pi}{2}} \frac{\cos^3 x}{\sin^3 x + \cos^3 x} dx \dots(2)$$

On adding eq.(1) and (2)

$$2y = \int_0^{\frac{\pi}{2}} \frac{\sin^3 x}{\sin^3 x + \cos^3 x} dx + \int_0^{\frac{\pi}{2}} \frac{\cos^3 x}{\sin^3 x + \cos^3 x} dx$$

$$2y = \int_0^{\frac{\pi}{2}} \frac{\sin^3 x + \cos^3 x}{\sin^3 x + \cos^3 x} dx$$

$$y = \frac{1}{2} \int_0^{\frac{\pi}{2}} dx$$

$$y = \frac{1}{2} \left(\frac{\pi}{2} - 0 \right)$$

$$y = \frac{\pi}{4}$$

32. Question

$$\int_0^{\pi/2} \frac{\sin x}{\sin x + \cos x} dx \text{ equals to}$$

- A. π
- B. $\pi/2$
- C. $\pi/3$
- D. $\pi/4$

Answer

$$\text{Given, } \int_0^{\pi/2} \frac{\sin x}{\sin x + \cos x} dx$$

$$\text{Now, } \sin x = A(\sin x + \cos x) + B \frac{d}{dx}(\sin x + \cos x)$$

$$\sin x = A(\sin x + \cos x) + B(\cos x - \sin x)$$

Equating ' $\sin x$ ' coeff:- Equating ' $\cos x$ ' coeff:-

$$1=A-B \quad 0=A+B$$

$$A-B=1$$



$$A+B=0$$

$$\underline{2A=1}$$

$$A=\frac{1}{2}$$

$$B=-\frac{1}{2}$$

$$\sin x = \frac{1}{2}(\sin x + \cos x) - \frac{1}{2}(\cos x - \sin x)$$

$$= \int_0^{\pi/2} \frac{\frac{1}{2}(\sin x + \cos x) - \frac{1}{2}(\cos x - \sin x)}{\sin x + \cos x} dx$$

$$= \int_0^{\pi/2} \frac{1}{2} \left(\frac{\sin x + \cos x}{\sin x + \cos x} \right) dx - \int_0^{\pi/2} \frac{1}{2} \left(\frac{\cos x - \sin x}{\sin x + \cos x} \right) dx$$

$$= \frac{1}{2} \int_0^{\pi/2} 1 dx - \frac{1}{2} [\log(\sin x + \cos x)]_0^{\pi/2}$$

$$= \frac{1}{2} [\pi]_0^{\pi/2} - \frac{1}{2} [\log(\sin x + \cos x)]_0^{\pi/2}$$

$$= \frac{1}{2} \left[\frac{\pi}{2} - 0 \right] - \frac{1}{2} \left[\log(\sin \frac{\pi}{2} + \cos \frac{\pi}{2}) - \log(\sin 0 + \cos 0) \right]$$

$$= \frac{\pi}{4} - \frac{1}{2} [\log(1+0) - \log(0+1)]$$

$$= \frac{\pi}{4} - \frac{1}{2}[0-0]$$

$$= \frac{\pi}{4} - 0$$

$$= \frac{\pi}{4}$$

Option A: - it's not option A , this is clearly justified on solving.

Option B: - it's not option B , this is clearly justified on solving.

Option C: - it's not option C, this is clearly justified on solving.

33. Question

$$\int_0^1 \frac{d}{dx} \left\{ \sin^{-1} \left(\frac{2x}{1+x^2} \right) \right\} dx \text{ is equal to}$$

A. 0

B. π

C. $\pi/2$

D. $\pi/4$

Answer

$$\text{Given, } \int_0^1 \frac{d}{dx} \left\{ \sin^{-1} \left(\frac{2x}{1+x^2} \right) \right\} dx$$

$$\begin{aligned}
&= \int_0^1 \frac{1}{\sqrt{1 - \left(\frac{2x}{1+x^2}\right)^2}} \cdot \frac{d}{dx} \left(\frac{2x}{1+x^2} \right) dx \\
&= \int_0^1 \frac{1+x^2}{\sqrt{(1+x^2)^2 - 4x^2}} \cdot \frac{(1+x^2)2 - 2x(2x)}{(1+x^2)^2} dx \\
&= \int_0^1 \frac{1}{\sqrt{(1-x^2)^2}} \cdot \frac{2+2x^2-4x^2}{1+x^2} dx \\
&= \int_0^1 \frac{1}{1-x^2} \cdot \frac{2-2x^2}{1+x^2} dx \\
&= \int_0^1 \frac{1}{1-x^2} \cdot \frac{2(1-x^2)}{1+x^2} dx \\
&= \int_0^1 \frac{2}{1+x^2} dx \\
&= 2[\tan^{-1}(x)]_0^1 \\
&= 2[\tan^{-1}(1) - \tan^{-1}(0)] \\
&= 2\left[\frac{\pi}{4} - 0\right] \\
&= \frac{\pi}{2}
\end{aligned}$$

Option A: - it's not option A , this is clearly justified on solving.

Option B: - it's not option B , this is clearly justified on solving.

Option D: - it's not option D, this is clearly justified on solving.

34. Question

$$\int_0^{\pi/2} x \sin x \, dx$$
 is equal to

- A. $\pi/4$
- B. $\pi/2$
- C. π
- D. 1

Answer

Given, $\int_0^{\pi/2} x \sin x \, dx$

$$\begin{aligned}
&= \left[x \int_0^{\pi/2} \sin x \, dx \right]_0^{\pi/2} - \int_0^{\pi/2} \left[1 \int_0^{\pi/2} \sin x \, dx \right] dx
\end{aligned}$$

$$= [x(-\cos x)]_0^{\pi/2} - \int_0^{\pi/2} (-\cos x) dx$$

$$= \left[\frac{\pi}{2}(0) - 0 \right] + [\sin x]_0^{\pi/2}$$

$$= 0 + (1-0)$$

$$= 1$$

Option A: - it's not option A , this is clearly justified on solving.

Option B: - it's not option B , this is clearly justified on solving.

Option C: - it's not option C, this is clearly justified on solving.

35. Question

$$\int_0^{\pi/2} \sin 2x \log \tan x dx \text{ is equal to}$$

A. π

B. $\pi/2$

C. 0

D. 2π

Answer

$$\text{Given, } \int_0^{\pi/2} \sin 2x \log(\tan x) dx$$

$$= \left[\log(\tan x) \int \sin 2x dx \right]_0^{\pi/2} - \int_0^{\pi/2} \frac{d}{dx} [\log(\tan x)] \int \sin 2x dx$$

$$= \left[\log(\tan x) \cdot \frac{-\cos 2x}{2} \right]_0^{\pi/2} - \int_0^{\pi/2} \frac{1}{\tan x} (\sec^2 x) \cdot \frac{-\cos 2x}{2} dx$$

$$= (0-0) - \int_0^{\pi/2} \frac{\cos x}{\sin x} \cdot \frac{1}{\cos^2 x} \cdot \frac{-\cos 2x}{2} dx$$

$$= 0 - \int_0^{\pi/2} \frac{-\cos 2x}{2 \sin x \cos x} dx$$

$$= \int_0^{\pi/2} \frac{\cos 2x}{\sin 2x} dx$$

$$= \int_0^{\pi/2} \tan 2x dx$$

$$= \left[\frac{\log |\sec 2x|}{2} \right]_0^{\pi/2}$$

$$= \frac{1}{2} [(\log |\sec \pi|) - \log |\sec 0|]$$

$$= \frac{1}{2} (\log 1 - \log 1)$$

$$= \frac{1}{2} (0 - 0)$$

$$= 0$$

Option A: - it's not option A , this is clearly justified on solving.

Option B: - it's not option B , this is clearly justified on solving.

Option D: - it's not option D , this is clearly justified on solving

36. Question

The value of $\int_0^{\pi} \frac{1}{5+3\cos x} dx$ is

A. $\pi/4$

B. $\pi/8$

C. $\pi/2$

D. 0

Answer

Given, $\int_0^{\pi} \frac{1}{5+3\cos x} dx$

Now put, $\tan \frac{x}{2} = t$

$$dx = \frac{2}{1+t^2} dt \text{ and } \cos x = \frac{1-t^2}{1+t^2}$$

Limits will also be changed accordingly,

From (0 to π) To (0 to ∞)

$$= \int_0^{\infty} \frac{1}{5+3(\frac{1-t^2}{1+t^2})} \cdot \frac{2}{1+t^2} dt$$

$$= \int_0^{\infty} \frac{1+t^2}{5+5t^2+3-3t^2} \cdot \frac{2}{1+t^2} dt$$

$$= \int_0^{\infty} \frac{2}{2t^2+8} dt$$

$$= \int_0^{\infty} \frac{1}{4+t^2} dt$$

$$= \int_0^{\infty} \frac{1}{(t)^2 + (2)^2} dt$$

$$= \left[\frac{1}{2} \tan^{-1} \left(\frac{t}{2} \right) \right]_0^{\infty}$$

$$= \frac{1}{2} [\tan^{-1}(\infty) - \tan^{-1}(0)]$$

$$= \frac{1}{2} \left[\frac{\pi}{2} - 0 \right]$$

$$= \frac{\pi}{4}$$

Option B: - it's not option B , this is clearly justified on solving.

Option C: - it's not option C , this is clearly justified on solving.

Option D: - it's not option D , this is clearly justified on solving

37. Question

$$\int_0^{\infty} \log\left(x + \frac{1}{x}\right) \frac{1}{1+x^2} dx =$$

A. $\pi \ln 2$

B. $-\pi \ln 2$

C. 0

D. $-\frac{\pi}{2} \ln 2$

Answer

Given, $\int_0^{\infty} \log\left(x + \frac{1}{x}\right) \frac{1}{1+x^2} dx$

Put $x = \tan \theta$

$$dx = \sec^2 \theta d\theta$$

Limits also will be changed accordingly,

$$x = 0 \rightarrow \theta = 0$$

$$x = \infty \rightarrow \theta = \frac{\pi}{2}$$

$$= \int_0^{\pi/2} \log(\tan \theta + \cot \theta) \frac{1}{1+\tan^2 \theta} \sec^2 \theta d\theta$$

$$= \int_0^{\pi/2} \log(\tan \theta + \cot \theta) \frac{1}{\sec^2 \theta} \sec^2 \theta d\theta$$

$$= \int_0^{\pi/2} \log(\tan \theta + \cot \theta) d\theta$$

$$= \int_0^{\pi/2} \log\left(\frac{\sin \theta}{\cos \theta} + \frac{\cos \theta}{\sin \theta}\right) d\theta$$

$$= \int_0^{\pi/2} \log\left(\frac{\sin^2 \theta + \cos^2 \theta}{\sin \theta \cos \theta}\right) d\theta$$

$$= \int_0^{\pi/2} \log\left(\frac{1}{\sin \theta \cos \theta}\right) d\theta$$

$$= \int_0^{\pi/2} \log(\sin \theta \cos \theta)^{-1} d\theta$$

$$\begin{aligned}
 &= - \int_0^{\pi/2} \log(\sin \theta \cos \theta) d\theta \\
 &= - \left[\int_0^{\pi/2} \log(\sin \theta) d\theta + \int_0^{\pi/2} \log(\cos \theta) d\theta \right]
 \end{aligned}$$

(Some standard notations which we need to remember)

$$= - \left[-\frac{\pi}{2} \log 2 - \frac{\pi}{2} \log 2 \right]$$

$$= -[-\pi \log 2]$$

$$= \pi \log 2$$

Option B: - it's not option B, this is clearly justified on solving.

Option C: - it's not option C , this is clearly justified on solving.

Option D: - it's not option D , this is clearly justified on solving

38. Question

$\int_0^{2a} f(x) dx$ is equal to

A. $2 \int_0^a f(x) dx$

B. 0

C. $\int_0^a f(x) dx + \int_0^a f(2a-x) dx$

D. $\int_0^a f(x) dx + \int_0^{2a} f(2a-x) dx$

Answer

We know that $\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx$

$$y = \int_0^{2a} f(x) dx = \int_0^a f(x) dx + \int_a^{2a} f(x) dx$$

$$A = \int_0^a f(x) dx$$

$$B = \int_a^{2a} f(x) dx$$

Let, $t = 2a - x \Rightarrow x = 2a - t$

Differentiating both side with respect to x

$$\frac{dt}{dx} = -1 \Rightarrow dx = -dt$$

At $x = a$, $t = a$

At $x = 2a$, $t = 0$

$$B = - \int_a^0 f(2a - t) dt$$

Use $\int_a^b f(x) dx = - \int_b^a f(x) dx$ and $\int_a^b f(x) dx = \int_a^b f(t) dt$

$$B = \int_0^a f(2a - x) dx$$

The final is $y = A + B$

$$y = \int_0^a f(x) dx + \int_0^a f(2a - x) dx$$

39. Question

If $f(a + b - x) = f(x)$, then $\int_a^b x f(x) dx$ is equal to

A. $\frac{a+b}{2} \int_a^b f(b-x) dx$

B. $\frac{a+b}{2} \int_a^b f(b+x) dx$

C. $\frac{b-a}{2} \int_a^b f(x) dx$

D. $\frac{a+b}{2} \int_a^b f(x) dx$

Answer

Given, $f(a + b - x) = f(x)$

$$a + b - x = x$$

$$a + b = 2x$$

$$x = \frac{a+b}{2}$$

Now, $\int_a^b x f(x) dx$

$$= \int_a^b \frac{a+b}{2} f(x) dx$$

$$= \frac{a+b}{2} \int_a^b f(x) dx$$

Option A: - it's not option A , this is clearly justified on solving.

Option B: - it's not option B , this is clearly justified on solving.

Option C: - it's not option C , this is clearly justified on solving

40. Question

The value of $\int_0^1 \tan^{-1} \left(\frac{2x-1}{1+x-x^2} \right) dx$, is

- A. 1
- B. 0
- C. -1
- D. $\pi/4$

Answer

$$y = \int_0^1 \tan^{-1} \left(\frac{x+x-1}{1-x(x-1)} \right) dx$$

$$\text{Use } \tan^{-1} \left(\frac{a+b}{1-ab} \right) = \tan^{-1} a + \tan^{-1} b$$

$$y = \int_0^1 \tan^{-1} x + \tan^{-1}(x-1) dx$$

$$y = \int_0^1 \tan^{-1} x - \tan^{-1}(1-x) dx \dots(1)$$

$$\text{Use king's property } y = \int_a^b f(x) dx = \int_a^b f(a+b-x) dx$$

$$y = \int_0^1 \tan^{-1}(1-x) - \tan^{-1}(1-(1-x)) dx$$

$$y = \int_0^1 \tan^{-1}(1-x) - \tan^{-1} x dx \dots(2)$$

On adding eq(1) and (2)

$$2y = \int_0^1 \tan^{-1} x - \tan^{-1}(1-x) dx + \int_0^1 \tan^{-1}(1-x) - \tan^{-1} x dx$$

$$2y = 0$$

$$y = 0$$

41. Question

The value of $\int_0^{\pi/2} \log \left(\frac{4+3 \sin x}{4+3 \cos x} \right) dx$ is

- A. 2
- B. 3/4
- C. 0
- D. -2

Answer

$$y = \int_0^{\pi/2} \log \left(\frac{4+3 \sin x}{4+3 \cos x} \right) dx \dots(1)$$

$$\text{Use king's property } y = \int_a^b f(x) dx = \int_a^b f(a+b-x) dx$$

$$y = \int_0^{\pi/2} \log \left(\frac{4+3 \sin \left(\frac{\pi}{2}-x \right)}{4+3 \cos \left(\frac{\pi}{2}-x \right)} \right) dx$$

$$y = \int_0^{\frac{\pi}{2}} \log\left(\frac{4+3\cos x}{4+3\sin x}\right) dx \dots(2)$$

On adding eq.(1) and (2)

$$2y = \int_0^{\frac{\pi}{2}} \log\left(\frac{4+3\sin x}{4+3\cos x}\right) dx + \int_0^{\frac{\pi}{2}} \log\left(\frac{4+3\cos x}{4+3\sin x}\right) dx$$

$$2y = \int_0^{\frac{\pi}{2}} \log\left(\frac{4+3\sin x}{4+3\cos x}\right)\left(\frac{4+3\cos x}{4+3\sin x}\right) dx$$

$$2y = \int_0^{\frac{\pi}{2}} \log 1 dx$$

$$y = 0$$

42. Question

The value of $\int_{-\pi/2}^{\pi/2} (x^3 + x \cos x + \tan^5 x + 1) dx$, is

- A. 0
- B. 2
- C. π
- D. 1

Answer

Given, $\int_{-\pi/2}^{\pi/2} (x^3 + x \cos x + \tan^5 x + 1) dx$

$$\begin{aligned} &= \left[\frac{x^4}{4} \right]_{-\pi/2}^{\pi/2} + \left\{ [x(\sin x)]_{-\pi/2}^{\pi/2} - \int_{-\pi/2}^{\pi/2} 1 \cdot \sin x dx \right\} + \int_{-\pi/2}^{\pi/2} \tan^3 x \cdot \tan^2 x \\ &\quad + (x)_{-\pi/2}^{\pi/2} \\ &= \left[\frac{x^4}{4} \right]_{-\pi/2}^{\pi/2} + \left[\frac{\pi}{2} \left(\sin \frac{\pi}{2} \right) - \left(-\frac{\pi}{2} \left(\sin \frac{-\pi}{2} \right) \right) \right] \\ &\quad + \int_{-\pi/2}^{\pi/2} \tan^3 x (1 + \sec^2 x) dx + (x)_{-\pi/2}^{\pi/2} \\ &= \left[\frac{x^4}{4} \right]_{-\pi/2}^{\pi/2} + \frac{\pi}{2}(1) - \frac{\pi}{2}(1) + \int_{-\pi/2}^{\pi/2} \tan x (1 + \sec^2 x) dx + \int_{-\pi/2}^{\pi/2} \tan^3 x \cdot \sec^2 x dx + (x)_{-\pi/2}^{\pi/2} \\ &= \frac{1}{4} \left[\left(\frac{\pi}{2} \right)^4 - \left(-\frac{\pi}{2} \right)^4 \right] + 0 + \int_{-\pi/2}^{\pi/2} \tan x + \int_{-\pi/2}^{\pi/2} \tan x \cdot \sec^2 x dx + \left[\frac{\tan^4 x}{4} \right]_{-\pi/2}^{\pi/2} + \left[\frac{\pi}{2} - \left(-\frac{\pi}{2} \right) \right] \\ &= \frac{1}{4}(0) + 0 + [\log |\sec x|]_{-\pi/2}^{\pi/2} + \left[\frac{\tan^2 x}{2} \right]_{-\pi/2}^{\pi/2} + \left[\frac{\tan^4 x}{4} \right]_{-\pi/2}^{\pi/2} + \pi \\ &= 0 + 0 + 0 + 0 + \pi \\ &= \pi \end{aligned}$$

Option A: - it's not option A, this is clearly justified on solving.

Option B: - it's not option B, this is clearly justified on solving.

Option D: - it's not option D, this is clearly justified on solving