

Circle

Practice Set 3.1

Q. 1. In the adjoining figure the radius of a circle with centre C is 6 cm, line AB is a tangent at A. Answer the following questions.

- (1) What is the measure of $\angle CAB$? Why?
- (2) What is the distance of point C from line AB? Why?
- (3) $d(A,B) = 6$ cm, find $d(B,C)$.
- (4) What is the measure of $\angle ABC$? Why?




Fig. 3.19

Answer : (1) Here CA is the radius of the circle and A is the point of contact of the tangent AB.

$\Rightarrow \angle CAB = 90^\circ$ Using tangent-radius theorem which states that a tangent at any point of a circle is perpendicular to the radius at the point of contact.

(2) CA is the radius of the circle which is perpendicular to the tangent AB.

So, the perpendicular distance of line AB from C = CA = 6 cm

(3) In triangle ABC right-angled at A,

Given AB = 6 cm and CA = 6 cm

$$BC^2 = AB^2 + CA^2 \text{ \{Using Pythagoras theorem\}}$$

$$\Rightarrow BC^2 = 6^2 + 6^2$$

$$\Rightarrow BC^2 = 36 + 36$$

$$\Rightarrow BC = \sqrt{72}$$

$$\Rightarrow BC = 6\sqrt{2} \text{ cm}$$

(4) In triangle ABC right-angled at A,

$$AB = CA = 6 \text{ cm}$$

$$\Rightarrow \angle ABC = \angle ACB \text{ \{Angles opposite to equal sides are equal\}}$$

$$\Rightarrow \angle ABC + \angle ACB + \angle BAC = 180^\circ \text{ \{Angle sum property of the triangle\}}$$

$$\Rightarrow 2\angle ABC = 90^\circ \text{ \{ } \because \angle BAC = 90^\circ \text{ \}}$$

$$\Rightarrow \angle ABC = 45^\circ$$

Q. 2. In the adjoining figure, O is the centre of the circle. From point R, seg RM and seg RN are tangent segments touching the circle at M and N. If (OR) = 10 cm and radius of the circle = 5 cm, then

(1) What is the length of each tangent segment?

(2) What is the measure of $\angle MRO$?

(3) What is the measure of $\angle MRN$?




Fig. 3.20

Answer : (1) Here OM is the radius of the circle and M and N are the points of contact of MR and NR respectively.

$\Rightarrow \angle RMO = 90^\circ$ Using tangent-radius theorem which states that a tangent at any point of a circle is perpendicular to the radius at the point of contact.

In triangle ORM right-angled at M,

Given that OR = 10 cm and OM = 5 cm {Radius of the circle}

$$OR^2 = OM^2 + RM^2 \text{ \{Using Pythagoras theorem\}}$$

$$\Rightarrow MR^2 = 10^2 - 5^2$$

$$\Rightarrow MR^2 = 100 - 25$$

$$\Rightarrow MR = \sqrt{75}$$

$$\Rightarrow MR = 5\sqrt{3} \text{ cm}$$

Also, $RN = 5\sqrt{3} \text{ cm}$ $\{\because$ Tangents from the same external point are congruent to each other. $\}$

$$(2) \tan R = \frac{OM}{MR} = \frac{5}{5\sqrt{3}}$$

$$\Rightarrow \tan R = \frac{1}{\sqrt{3}} = \tan 30^\circ$$

$$\Rightarrow \angle MRO = 30^\circ$$

$$(3) \text{ Similarly, } \angle NRO = 30^\circ$$

$$\Rightarrow \angle MRN = \angle MRO + \angle NRO = 30^\circ + 30^\circ = 60^\circ$$

Q. 3. Seg RM and seg RN are tangent segments of a circle with centre O. Prove that seg OR bisects $\angle MRN$ as well as $\angle MON$.




Fig. 3.21

Answer : In triangle MOR and triangle NOR,

$MR = NR$ $\{\because$ Tangents from same external point are congruent to each other. $\}$

$OR = OR$ {Common}

$OM = ON$ {Radius of the circle}

$\Rightarrow \triangle MOR \cong \triangle NOR$ {By SSS}

$\Rightarrow \angle ROM = \angle RON$

And $\angle MRO = \angle NRO$ {C.P.C.T.}

Hence proved that seg OR bisects $\angle MRN$ as well as $\angle MON$.


Q. 4. What is the distance between two parallel tangents of a circle having radius 4.5 cm ? Justify your answer.

Answer : Let BC and DE be the parallel tangents to a circle centered at A with point of contact O and H respectively. On joining OH, we find OH is the diameter of the circle. $\angle BOA = 90^\circ = \angle DHA$ {Using tangent-radius theorem which states that a tangent at any point of a circle is perpendicular to the radius at the point of contact.}

Distance between BC and DE = OH

\therefore OH is perpendicular to BC and DE.

$OH = 2 \times 4.5 \text{ cm} = 9 \text{ cm}$



Practice Set 3.2

Q. 1. Two circles having radii 3.5 cm and 4.8 cm touch each other internally. Find the distance between their centres.

Answer : Given: Two circles are touching each other internally.

\therefore The distance between the centres of the circles touching internally is equal to the difference of their radii.

\Rightarrow Distance between their centres = $4.8 \text{ cm} - 3.5 \text{ cm} = 1.3 \text{ cm}$

Q. 2. Two circles of radii 5.5 cm and 4.2 cm touch each other externally. Find the distance between their centres.

Answer : Given: Two circles are touching each other externally


We know that if the circles touch each other externally, distance between their centres is equal to the sum of their radii.

$$\Rightarrow \text{Distance between their centres} = 5.5 \text{ cm} + 4.2 \text{ cm} = 9.7 \text{ cm}$$

Q. 3. If radii of two circles are 4 cm and 2.8 cm. Draw figure of these circles touching each other –


- (i) externally
- (ii) internally.

Answer :



Steps of construction:

1. Draw a circle with radius 4cm and centre A.
2. Draw another circle with radius 2.8 cm and centre B such that they touch each other externally.




Steps of construction:

1. Draw a circle with radius 4cm and centre A.
2. Draw another circle with radius 2.8 cm and centre B such that they touch each other internally.

Q. 4. In fig 3.27, the circles with centres P and Q touch each other at R. A line passing through R meets the circles at A and B respectively. Prove that -

- (1) $\text{seg AP} \parallel \text{seg BQ}$,
- (2) $\triangle APR \sim \triangle RQB$, and
- (3) Find $\angle RQB$ if $\angle PAR = 35^\circ$



Answer : (1) In $\triangle APR$,

$AP = RP$ {Radius of the circle with centre P}

$\angle PAR = \angle PRA \dots (1)$

In $\triangle RQB$,

$RQ = QB$ {Radius of the circle with centre Q}

$$\angle QRB = \angle QBR \dots (2)$$

$$\Rightarrow \angle PRA = \angle QRB \text{ \{Vertically Opposite Angle\} } \dots (3)$$

$$\Rightarrow \angle PAR = \angle QBR \text{ \{From (1), (2) and (3)\}}$$

\Rightarrow Alternate interior angles are equal.

$$\Rightarrow AP \parallel BQ$$

Hence, proved.

(2) In $\triangle APR$ and $\triangle RQB$,

$$\angle PAR = \angle QBR \text{ and } \angle PRA = \angle QRB \text{ \{From (1) and (2)\}}$$

$$\Rightarrow \triangle APR \sim \triangle RQB \text{ \{AA\}}$$

Hence, proved.

(4) Given: $\angle PAR = 35^\circ$

$$\Rightarrow \angle QBR = 35^\circ = \angle QRB \text{ \{Proved previously\}}$$


In $\triangle RQB$,

$$\Rightarrow \angle RQB + \angle QRB + \angle QBR = 180^\circ \text{ \{Angle sum property of the triangle\}}$$


$$\Rightarrow \angle RQB + 35^\circ + 35^\circ = 180^\circ$$

$$\Rightarrow \angle RQB = 180^\circ - 70^\circ = 110^\circ$$

Q. 5. In fig 3.28 the circles with centres A and B touch each other at E. Line is a common tangent which touches the circles at C and D respectively. Find the length of seg CD if the radii Fig. 3.28 of the circles are 4 cm, 6 cm.



Answer :



Given that two circles with centre A and B touch each other externally. We know that if the circles touch each other externally, distance between their centres is equal to the sum of their radii.

$$\Rightarrow AB = (4 + 6) \text{ cm} = 10 \text{ cm}$$

In $\triangle ABC$ right-angles at A,

$$BC^2 = CA^2 + AB^2 \text{ \{Using Pythagoras theorem\}}$$

$$\Rightarrow BC^2 = 4^2 + 10^2$$

$$\Rightarrow BC^2 = 16 + 100$$

$$\Rightarrow BC = \sqrt{116} \text{ cm}$$

In $\triangle DBC$,

$\angle BDC = 90^\circ$ because D is the point of contact of tangent CD to circle centred B

$$BC^2 = CD^2 + DB^2 \text{ \{Using Pythagoras theorem\}}$$

$$\Rightarrow CD^2 = 116 - 6^2$$


$$\Rightarrow CD^2 = 116 - 36$$

$$\Rightarrow CD = \sqrt{80} \text{ cm} = 4\sqrt{5}$$

Practice Set 3.3

Q. 1. In figure 3.37, points G, D, E, F are concyclic points of a circle with centre C.

$\angle ECF = 70^\circ$, $m(\text{arc DGF}) = 200^\circ$ find $m(\text{arc DE})$ and $m(\text{arc DEF})$.



Answer : Given $\angle ECF = 70^\circ$ and $m(\text{arc DGF}) = 200^\circ$

We know that measure of major arc = 360° - measure of minor arc

$$m(\text{arc DGF}) = 360^\circ - m(\text{arc DF})$$

$$\Rightarrow m(\text{arc DF}) = 360^\circ - 200^\circ = 160^\circ$$

$$\Rightarrow \angle DCF = 160^\circ$$

\therefore The measure of a minor arc is the measure of its central angle.

$$\therefore m(\text{arc DEF}) = 160^\circ$$

$$\text{So, } \angle DCE = \angle DCF - \angle ECF = 160^\circ - 70^\circ$$

$$\Rightarrow \angle DCE = 90^\circ$$

The measure of a minor arc is the measure of its central angle.

$$m(\text{arc DE}) = 90^\circ$$

Q. 2. In fig 3.38 $\triangle QRS$ is an equilateral triangle. Prove that,

(1) $\text{arc RS} \cong \text{arc QS} \cong \text{arc QR}$

(2) $m(\text{arc QRS}) = 240^\circ$.




Fig. 3.38

Answer : (1) Two arcs are congruent if their measures and radii are equal.

$\therefore \triangle QRS$ is an equilateral triangle

$$\therefore RS = QS = QR$$

$$\Rightarrow \text{arc } RS \cong \text{arc } QS \cong \text{arc } QR$$

(2) Let O be the centre of the circle.

$$m(\text{arc } QS) = \angle QOS$$

$$\angle QOS + \angle QOR + \angle SOR = 360^\circ$$

$$\Rightarrow 3\angle QOS = 360^\circ \{ \because \triangle QRS \text{ is an equilateral triangle} \}$$

$$\Rightarrow \angle QOS = 120^\circ$$

$$m(\text{arc } QS) = 120^\circ$$

$$m(\text{arc } QRS) = 360^\circ - 120^\circ \{ \because \text{Measure of a major arc} = 360^\circ - \text{measure of its corresponding minor arc} \}$$

$$\Rightarrow m(\text{arc } QRS) = 240^\circ$$

Q. 3. In fig 3.39 chord $AB \cong$ chord CD , Prove that, arc $AC \cong$ arc BD




Fig. 3.39

Answer : \because Chord $AB \cong$ chord CD

$\therefore m(\text{arc } AB) = m(\text{arc } CD)$ {Corresponding arcs of congruent chords of a circle (or congruent circles) are congruent}

Subtract $m(\text{arc } CB)$ from above,

$$m(\text{arc } AB) - m(\text{arc } CB) = m(\text{arc } CD) - m(\text{arc } CB)$$

$$\Rightarrow m(\text{arc } AC) = m(\text{arc } BD)$$

$$\Rightarrow \text{arc } AC \cong \text{arc } BD$$

Hence, proved.

Practice Set 3.4

Q. 1. In figure 3.56, in a circle with centre O, length of chord AB is equal to the radius of the circle. Find measure of each of the following.




Fig. 3.56

- (1) $\angle AOB$ (2) $\angle ACB$
 (3) arc AB (4) arc ACB.

Answer : (1) In $\triangle AOB$,

$AB = OA = OB = \text{radius of circle}$

$\Rightarrow \triangle AOB$ is an equilateral triangle

$\angle AOB + \angle ABO + \angle BAO = 180^\circ$ {Angle sum property}

$\Rightarrow 3\angle AOB = 180^\circ$ {All the angles are equal}

$\angle AOB = 60^\circ$

(2) $\angle AOB = 2 \times \angle ACB$ {The measure of an inscribed angle is half the measure of the arc intercepted by it.}

$\Rightarrow \angle ACB = 30^\circ$

(3) $\angle AOB = 60^\circ$

$\Rightarrow \text{arc}(AB) = 60^\circ$ {The measure of a minor arc is the measure of its central angle.}

(4) Using Measure of a major arc = $360^\circ - \text{measure of its corresponding minor arc}$

$\Rightarrow \text{arc}(ACB) = 360^\circ - \text{arc}(AB)$

$\Rightarrow \text{arc}(ACB) = 360^\circ - 60^\circ = 300^\circ$

Q. 2. In figure 3.57, $\square PQRS$ is cyclic. Side $PQ \cong \text{side } RQ$. $\angle PSR = 110^\circ$, Find-




Fig. 3.57

- (1) measure of $\angle PQR$
- (2) $m(\text{arc } PQR)$
- (3) $m(\text{arc } QR)$
- (4) measure of $\angle PRQ$

Answer : (1) Given PQRS is a cyclic quadrilateral.

\therefore Opposite angles of a cyclic quadrilateral are supplementary

$$\Rightarrow \angle PSR + \angle PQR = 180^\circ$$

$$\Rightarrow \angle PQR = 180^\circ - 110^\circ$$

$$\Rightarrow \angle PQR = 70^\circ$$

(2) $2 \times \angle PQR = m(\text{arc } PR)$ {The measure of an inscribed angle is half the measure of the arc intercepted by it.}

$$m(\text{arc } PR) = 140^\circ$$

$\Rightarrow m(\text{arc } PQR) = 360^\circ - 140^\circ = 220^\circ$ {Using Measure of a major arc = 360° - measure of its corresponding minor arc}

(3) Side $PQ \cong$ side RQ

$\therefore m(\text{arc } PQ) = m(\text{arc } RQ)$ {Corresponding arcs of congruent chords of a circle (or congruent circles) are congruent}

$$\Rightarrow m(\text{arc } PQR) = m(\text{arc } PQ) + m(\text{arc } RQ)$$

$$\Rightarrow m(\text{arc } PQR) = 2 \times m(\text{arc } PQ)$$

$$\Rightarrow m(\text{arc } PQ) = 110^\circ$$

(4) In $\triangle PQR$,

$$\angle PQR + \angle QRP + \angle RPQ = 180^\circ \text{ {Angle sum property}}$$

$$\Rightarrow \angle PRQ + \angle RPQ = 180^\circ - \angle PQR$$

$$\Rightarrow 2\angle PRQ = 180^\circ - 70^\circ \{\because \text{side } PQ \cong \text{side } RQ\}$$

$$\Rightarrow \angle PRQ = 55^\circ$$

Q. 3. $\square MRPN$ is cyclic, $\angle R = (5x - 13)^\circ$, $\angle N = (4x + 4)^\circ$. Find measures of \sqrt{R} and \sqrt{N} .

Answer : Given MRPN is a cyclic quadrilateral.

$$\Rightarrow \angle R + \angle N = 180^\circ \{\text{Using Opposite angles of a cyclic quadrilateral are supplementary}\}$$

$$\Rightarrow (5x - 13)^\circ + (4x + 4)^\circ = 180^\circ$$

$$\Rightarrow 9x - 9 = 180^\circ$$

$$\Rightarrow x - 1 = 20^\circ$$

$$\Rightarrow x = 21^\circ$$

$$\angle R = (5x - 13)^\circ = 5 \times 21 - 13 = 105 - 13 = 92^\circ$$

$$\angle N = (4x + 4)^\circ = 4 \times 21 + 4 = 84 + 4 = 88^\circ$$

Q. 4. In figure 3.58, seg RS is a diameter of the circle with centre O. Point T lies in the exterior of the circle. Prove that $\angle RTS$ is an acute angle.





Fig. 3.58

Answer :



Given RS is the diameter

$$\Rightarrow \angle ROS = 180^\circ$$

$$m(\text{arc RS}) = 180^\circ$$

Now, $\angle RTS$ is an external angle.

$$\angle RTS = \frac{1}{2}[m(\text{arc RS}) - m(\text{arc PQ})]$$


$$\Rightarrow \angle RTS = \frac{1}{2}[180^\circ - m(\text{arc PQ})]$$

$$\Rightarrow \angle RTS = 90^\circ - \frac{1}{2}m(\text{arc PQ})$$

Hence, $\angle RTS$ is an acute angle.

Q. 5. Prove that, any rectangle is a cyclic quadrilateral.

Answer :



In ABCD,

$\angle A = 90^\circ$ { \because angle of a rectangle is 90° . }

$\angle C = 90^\circ$ { opposite angles are equals }

$\Rightarrow \angle A + \angle C = 180^\circ$

If opposite angles are supplementary, the quadrilateral is cyclic.

\therefore ABCD is cyclic.

Q. 6. In figure 3.59, altitudes YZ and XT of

$\triangle WXY$ intersect at P. Prove that,




Fig. 3.59

(1) $\square WZPT$ is cyclic.

(2) Points X, Z, T, Y are concyclic.

Answer : (1) In WZPT,

$\angle WZP = \angle WTP = 90^\circ$ { YZ and XT are the altitudes }

If a pair of opposite angles of a quadrilateral is supplementary, then the quadrilateral is cyclic.

\Rightarrow WZPT is cyclic.

(2) \because X, Z, T, Y lie on same circle, \therefore they are concyclic.

Q. 7. In figure 3.60, $m(\text{arc NS}) = 125^\circ$, $m(\text{arc EF}) = 37^\circ$, find the measure $\angle NMS$.




Fig. 3.60

Answer : Given $m(\text{arc NS}) = 125^\circ$, $m(\text{arc EF}) = 37^\circ$

Also, $\angle NMS$ is an external angle, so

$$\angle NMS = \frac{1}{2} [m(\text{arc NS}) - m(\text{arc EF})]$$

$$\Rightarrow \angle NMS = \frac{1}{2} [125^\circ - 37^\circ]$$

$$\Rightarrow \angle NMS = \frac{1}{2} \times 88^\circ = 44^\circ$$

Q. 8. In figure 3.61, chords AC and DE intersect at B. If $\angle ABE = 108^\circ$, $m(\text{arc AE}) = 95^\circ$, find $m(\text{arc DC})$.




Fig. 3.61

Answer : Given $\angle ABE = 108^\circ$, $m(\text{arc AE}) = 95^\circ$

Using the property of the secant,

$$\angle ABE = \frac{1}{2} [m(\text{arc AE}) + m(\text{arc DC})]$$

$$\Rightarrow 108^\circ = \frac{1}{2}[95^\circ + m(\text{arc } DC)]$$

$$\Rightarrow m(\text{arc } DC) = 108^\circ \times 2 - 95^\circ$$

$$\Rightarrow m(\text{arc } DC) = 121^\circ$$

Practice Set 3.5

Q. 1. In figure 3.77, ray PQ touches the circle at point Q. PQ = 12, PR = 8, find PS and RS.




Fig. 3.77

Answer : Given PQ = 12, PR = 8

$$SP \times RP = PQ^2$$

This property is known as tangent secant segments theorem.

$$\Rightarrow PS \times 8 = 12^2$$

$$\Rightarrow PS = \frac{144}{8} = 18$$

$$RS = PS - RP = 18 - 8 = 10$$

Q. 2. In figure 3.78, chord MN and chord RS intersect at point D.

(1) If RD = 15, DS = 4, MD = 8 find DN

(2) If $RS = 18$, $MD = 9$, $DN = 8$ find DS




Fig. 3.78

Answer : (1) Given $RD = 15$, $DS = 4$, $MD = 8$

$$MD \times DN = RD \times DS$$

This property is known as theorem of chords intersecting inside the circle.

$$\Rightarrow 8 \times DN = 15 \times 4$$

$$\Rightarrow DN = \frac{15}{2} = 7.5$$

(2) Given $RS = 18$, $MD = 9$, $DN = 8$

Here, $RS = 18$

Let $RD = x$ and $DS = 18 - x$

$$MD \times DN = RD \times DS$$

This property is known as theorem of chords intersecting inside the circle.

$$\Rightarrow 8 \times 9 = x \times (18 - x)$$

$$\Rightarrow 18x - x^2 = 72$$

$$\Rightarrow x^2 - 18x + 72 = 0$$

$$\Rightarrow (x - 12)(x - 6) = 0$$

$$\Rightarrow x = 12 \text{ or } 6$$

$$\Rightarrow DS = 6 \text{ or } 12$$

Q. 3. In figure 3.79, O is the centre of the circle and B is a point of contact. seg $OE \perp$ seg AD, $AB = 12$, $AC = 8$, find




Fig. 3.79

- (1) AD
- (2) DC
- (3) DE.

Answer : (1) Given: $OE \perp AD$, $AB = 12$, $AC = 8$

$$\Rightarrow AD \times AC = AB^2$$

This property is known as tangent secant segments theorem.

$$\Rightarrow AD \times 8 = 12^2$$

$$\Rightarrow AD = \frac{144}{8} = 18$$

$$(2) DC = AD - AC = 18 - 8 = 10$$

(3) As we know that a perpendicular from centre divides the chord in two equal parts.
Here, $OE \perp AD$.


$$\Rightarrow DE = EC$$

$$\Rightarrow DE + EC = DC$$

$$\Rightarrow 2DE = DC$$

$$\Rightarrow DE = \frac{1}{2}DC = 5$$

Q. 4. In figure 3.80, if $PQ = 6$, $QR = 10$, $PS = 8$ find TS .



Answer : Given: $PQ = 6$, $QR = 10$, $PS = 8$

$$PT \times PS = PR \times PQ$$

This property is known as theorem of chords intersecting outside the circle.


$$\Rightarrow PR = PQ + RQ = 6 + 10 = 16$$

$$\Rightarrow PT \times 8 = 16 \times 6$$

$$\Rightarrow PT = 12$$

$$TS = PT - PS = 12 - 8 = 4$$

Q. 5. In figure 3.81, seg EF is a diameter and seg DF is a tangent segment. The radius of the circle is r . Prove that, $DE \times GE = 4r^2$



Answer : In $\triangle DEF$,

$\angle DFE = 90^\circ$ {Using tangent-radius theorem which states that a tangent at any point of a circle is perpendicular to the radius at the point of contact.}

Given: EF = diameter of the circle.

$$DE^2 = DF^2 + EF^2 \text{ \{Using Pythagoras theorem\}}$$

$$\Rightarrow DE^2 = DF^2 + (2r)^2$$

$$\Rightarrow DE^2 = DF^2 + 4r^2$$

$$\Rightarrow DF^2 = DE^2 - 4r^2$$

$$\text{Also, } DE \times DG = DF^2$$

This property is known as tangent secant segments theorem.

$$\Rightarrow DE \times DG = DE^2 - 4r^2$$

$$\Rightarrow DE^2 - DE \times DG = 4r^2$$

$$\Rightarrow DE(DE - DG) = 4r^2$$

$$\Rightarrow DE \times EG = 4r^2$$

Hence, proved.

Problem Set 3

Q. 1. A. Four alternative answers for each of the following questions are given. Choose the correct alternative.

Two circles of radii 5.5 cm and 3.3 cm respectively touch each other. What is the distance between their centers?

A. 4.4 cm

B. 8.8 cm

C. 2.2 cm

D. 8.8 or 2.2 cm

Answer : Given that both the circles touch each other but not specified externally or internally.

The distance between the centres of the circles touching internally is equal to the difference of their radii.

$$\Rightarrow \text{Distance between their centres} = 5.5 \text{ cm} - 3.3 \text{ cm} = 2.2 \text{ cm}$$

If the circles touch each other externally, distance between their centres is equal to the sum of their radii.




\Rightarrow Distance between their centres = $5.5 \text{ cm} + 3.3 \text{ cm} = 8.8 \text{ cm}$

Q. 1. B. Four alternative answers for each of the following questions are given. Choose the correct alternative.

Two circles intersect each other such that each circle passes through the centre of the other. If the distance between their centres is 12, what is the radius of each circle?

- A. 6 cm
- B. 12 cm
- C. 24 cm
- D. can't say

Answer : Given $OA = 12$



From the figure, OA is the radius of both the circles.

Given that distance between their centres is $OA = 12$


\therefore Radius of the circles = 12

Q. 1. C. Four alternative answers for each of the following questions are given. Choose the correct alternative.

A circle touches all sides of a parallelogram. So the parallelogram must be a,..... .

- A. rectangle
- B. rhombus
- C. square
- D. trapezium

Answer :



Let ABCD be a parallelogram which circumscribes the circle.

$AP = AS$ [Tangents drawn from an external point to a circle are equal in length]

$BP = BQ$ [Tangents drawn from an external point to a circle are equal in length]

$CR = CQ$ [Tangents drawn from an external point to a circle are equal in length]

$DR = DS$ [Tangents drawn from an external point to a circle are equal in length]

Consider, $(AP + BP) + (CR + DR) = (AS + DS) + (BQ + CQ)$

$AB + CD = AD + BC$

But $AB = CD$ and $BC = AD$ [Opposite sides of parallelogram ABCD]

$AB + CD = AD + BC$

Hence $2AB = 2BC$

Therefore, $AB = BC$

Similarly, we get $AB = DA$ and $DA = CD$

Thus, ABCD is a rhombus.


Q. 1. D. Four alternative answers for each of the following questions are given. Choose the correct alternative.

Length of a tangent segment drawn from a point which is at a distance 12.5 cm from the centre of a circle is 12 cm, find the diameter of the circle.

A. 25 cm

- B. 24 cm
- C. 7 cm
- D. 14 cm

Answer :



Given: $BO = 12.5$ cm and $AB = 12$ cm

In $\triangle AOB$,

$\angle OAB = 90^\circ$ {Using tangent-radius theorem which states that a tangent at any point of a circle is perpendicular to the radius at the point of contact.}

$BO^2 = AB^2 + OA^2$ {Using Pythagoras theorem}

$$\Rightarrow (12.5)^2 = 12^2 + OA^2$$

$$\Rightarrow OA^2 = 156.25 - 144$$

$$\Rightarrow OA = \sqrt{12.25}$$

$$\Rightarrow OA = 3.5 \text{ cm}$$

Radius = 3.5 cm

$$\Rightarrow \text{Diameter} = 7 \text{ cm}$$


Q. 1. E. Four alternative answers for each of the following questions are given. Choose the correct alternative.

If two circles are touching externally, how many common tangents of them can be drawn?

- A. One
- B. Two

- C. Three
- D. Four

Answer :



If two circles are touching each other externally, they have 3 tangents in common. The above figure proves this statement.


There are three common tangents for the given two circles.

Q. 1. F. Four alternative answers for each of the following questions are given. Choose the correct alternative.

$\angle ACB$ is inscribed in arc ACB of a circle with centre O . If $\angle ACB = 65^\circ$, find $m(\text{arc } ACB)$.

- A. 65°
- B. 130°
- C. 295°
- D. 230°

Answer :



Given $\angle ACB = 65^\circ$

$\Rightarrow \angle AOB = 2 \times 65^\circ = 130^\circ$ $\{\because$ The measure of an inscribed angle is half the measure of the arc intercepted by it. $\}$

$$m(\text{AB}) = 130^\circ$$

So, $m(\text{arc ACB}) = 360^\circ - m(\text{AB})$ $\{\because$ Measure of a major arc = 360° - measure of its corresponding minor arc $\}$

$$\Rightarrow m(\text{arc ACB}) = 360^\circ - 130^\circ = 230^\circ$$

Q. 1. G. Four alternative answers for each of the following questions are given. Choose the correct alternative.

Chords AB and CD of a circle intersect inside the circle at point E. If $AE = 5.6$, $EB = 10$, $CE = 8$, find ED.


A. 7

B. 8

C. 11.2

D. 9

Answer :



Given: $AE = 5.6$, $EB = 10$, $CE = 8$

We know that $AE \times EB = CE \times ED$

This property is known as theorem of chords intersecting inside the circle.

$$\Rightarrow 5.6 \times 10 = 8 \times ED$$

$$\Rightarrow ED = 7$$

Q. 1. H. Four alternative answers for each of the following questions are given. Choose the correct alternative.

In a cyclic □ ABCD, twice the measure of $\angle A$ is thrice the measure of $\angle C$. Find the measure of $\angle C$?

- A. 36
- B. 72
- C. 90
- D. 108

Answer : Given that $2\angle A = 3\angle C$

We know that in a cyclic quadrilateral opposite angles are supplementary to each other.

$$\Rightarrow \angle A + \angle C = 180^\circ$$

$$\Rightarrow \frac{3}{2}\angle C + \angle C = 180^\circ$$

$$\Rightarrow \frac{5}{2}\angle C = 180^\circ$$

$$\Rightarrow \angle C = 72^\circ$$

Q. 1. I. Four alternative answers for each of the following questions are given. Choose the correct alternative.

Points A, B, C are on a circle, such that $m(\text{arc AB}) = m(\text{arc BC}) = 120^\circ$. No point, except point B, is common to the arcs. Which is the type of $\triangle ABC$?

- A. Equilateral triangle
- B. Scalene triangle
- C. Right angled triangle
- D. Isosceles triangle

Answer : Angle subtended by the arcs at centre = 120°

\Rightarrow Angle subtended by the arc at the remaining part of the circle = 60° {The measure of an inscribed angle is half the measure of the arc intercepted by it.}

\therefore Interior angles of the triangle ABC = 60°

\therefore It is an equilateral triangle.

Q. 1. J. Four alternative answers for each of the following questions are given. Choose the correct alternative.




Seg XZ is a diameter of a circle. Point Y lies in its interior. How many of the following statements are true?


- (i) It is not possible that $\angle XYZ$ is an acute angle.
- (ii) $\angle XYZ$ can't be a right angle.
- (iii) $\angle XYZ$ is an obtuse angle.
- (iv) Can't make a definite statement for measure of $\angle XYZ$.

- A. Only one
- B. Only two
- C. Only three
- D. All


Answer : If Y would have lied on circumference $\angle XYZ = 90^\circ$ since XZ is the diameter.



If Y lied outside the circle, $\angle XYZ =$ acute angle



$\therefore \angle XYZ$ is an obtuse angle.



Statements (i), (ii) and (iii) are true.

Q. 2. Line l touches a circle with centre O at point P . If radius of the circle is 9 cm, answer the following.

- (1) What is $d(O, P)$ = ? Why ?
- (2) If $d(O, Q) = 8$ cm, where does the point Q lie?
- (3) If $d(PQ) = 15$ cm, How many locations of point R are line online l ? At what distance will each of them be from point P ?




Fig. 3.82

Answer : (1) The perpendicular distance of O from P = radius of the circle = 9 cm.

(2) Q lies in the interior of the circle because P lying on the circumference of the circle is at a distance of 9 cm.

(3) Position of R is not specified.

Q. 3. In figure 3.83, M is the centre of the circle and seg KL is a tangent segment.

If $MK = 12$, $KL = 6\sqrt{3}$ then find -

(1) Radius of the circle.

(2) Measures of $\angle K$ and $\angle M$.




Fig. 3.83

Answer : (1) Here LM is the radius of the circle

$\Rightarrow \angle KML = 90^\circ$ Using tangent-radius theorem which states that a tangent at any point of a circle is perpendicular to the radius at the point of contact.

In triangle MLK right-angled at L,

Given $MK = 12$, $KL = 6\sqrt{3}$,

$MK^2 = LM^2 + KL^2$ {Using Pythagoras theorem}

$$\Rightarrow LM^2 = 12^2 - 6\sqrt{3}^2$$

$$\Rightarrow LM^2 = 144 - 108$$

$$\Rightarrow LM = \sqrt{36}$$

$$\Rightarrow LM = 6 \text{ cm}$$

$$(2) \tan K = \frac{ML}{LK}$$

$$\Rightarrow \tan K = \frac{6}{6\sqrt{3}} = \frac{1}{\sqrt{3}} = \tan 30^\circ$$

$$\Rightarrow \angle K = 30^\circ$$

$$\angle M + \angle K + \angle L = 180^\circ \text{ \{Angle sum property of the triangle\}}$$

$$\text{So, } \angle M = 180^\circ - \angle K - \angle L$$

$$\Rightarrow \angle M = 180^\circ - 30^\circ - 90^\circ = 60^\circ$$

Q. 4. In figure 3.84, O is the centre of the circle. Seg AB, seg AC are tangent segments. Radius of the circle is r and $l(AB) = r$, Prove that, $\square ABOC$ is a square.




Fig. 3.84

Answer : Given: $AB = r =$ radius of the circle

Here, $AB = AC = r$ {tangents from the same external point are equal}

And $OB = OC = r =$ radius of the circle.

$\Rightarrow \angle OBA = \angle OCA = 90^\circ$ Using tangent-radius theorem which states that a tangent at any point of a circle is perpendicular to the radius at the point of contact.

\therefore Sides of $ABOC$ are equal and opposite angles are 90° each

Hence, $ABOC$ is a square.

Q. 5. In figure 3.85, $\square ABOC$ is a parallelogram. It circumscribes the circle with centre T . Point E, F, G, H are touching points. If $AE = 4.5, EB = 5.5$, find AD .




Fig. 3.85

Answer : Given: $AE = 4.5, EB = 5.5$

Here, $AE = AH = 4.5$ {tangents from same external point are equal}

$EB = BF = 5.5$ {tangents from same external point are equal}

\therefore Opposite sides of a parallelogram are equal

$\therefore AE = DG$ and $EB = GC$

Also, $DH = DG = 4.5$ {tangents from same external point are equal}

And $FC = GC = 5.5$ {tangents from same external point are equal}

$$\Rightarrow AD = AH + HD = 10$$

Q. 6. In figure 3.86, circle with centre M touches the circle with centre N at point T. Radius RM touches the smaller circle at S. Radii of circles are 9 cm and 2.5 cm. Find the answers to the following questions hence find the ratio MS:SR.

- (1) Find the length of segment MT
- (2) Find the length of seg MN
- (3) Find the measure of $\angle NSM$.




Fig. 3.86

Answer : (1) MT = radius of the big circle = 9 cm

$$(2) \quad MN = MT - TN = 9 - 2.5 = 6.5 \text{ cm}$$

(3) SM is the tangent to the circle with radius 2.5 cm with S being point of contact.

$\angle NSM = 90^\circ$ Using tangent-radius theorem which states that a tangent at any point of a circle is perpendicular to the radius at the point of contact.

In $\triangle MSN$,

$$\angle MSN = 90^\circ \{ \because MS \text{ is the tangent to the small circle with point of contact } S \}$$

$$\Rightarrow MN^2 = MS^2 + NS^2$$

$$MS^2 = MN^2 - NS^2$$

$$\Rightarrow MS^2 = 6.5^2 - 2.5^2$$

$$\Rightarrow MS^2 = 36$$

$$\Rightarrow MS = 6 \text{ cm}$$

Now, $SR = MR - MS = 9 - 6 = 3$ cm

$\Rightarrow MS: SR = 6:3 = 2:1$

Q. 7. In the adjoining figure circles with centres X and Y touch each other at point Z. A secant passing through Z intersects the circles at points A and B respectively. Prove that, radius $XA \parallel$ radius YB . Fill in the blanks and complete the proof.




Fig. 3.87

Construction: Draw segments XZ and YZ..... .

Proof: By theorem of touching circles, points X, Z, Y are ..concylic..... .

$\therefore \angle XZA \cong \angle YZB$ vertically opposite angles

Let $\angle XZA = \angle BZY = a$ (I)

Now, seg $XA \cong$ seg XZ (...radius of the same circle.....)

$\therefore \angle XAZ = \angle XZA$ = a (isosceles triangle theorem) (II)

similarly, seg $YB \cong$ seg YZ (...radius of the same circle.....)

$\therefore \angle BZY = \angle ZBY$ = a (.isosceles triangle theorem.....) (III)

\therefore from (I), (II), (III),

$\angle XAZ = \angle ZBY$

\therefore radius $XA \parallel$ radius YB (...since alternate interior angles are equal.....)

Answer : Construction: Draw segments XZ and YZ.

Proof: By theorem of touching circles, points X, Z, Y are concyclic.

$\angle XZA = \angle YZB$ {vertically opposite angles}

Let $\angle XZA = \angle BZY = a$ (I)

Now, $\text{seg } XA \cong \text{seg } XZ$ (radius of the same circle)

$\therefore \angle XAZ = \angle XZA = a$ (isosceles triangle theorem) (II)

Similarly, $\text{seg } YB \cong \text{seg } YZ$ (radius of the same circle)

$\therefore \angle BZY = \angle ZBY = a$ (isosceles triangle theorem) (III)

\therefore from (I), (II), (III),

$\angle XAZ = \angle ZBY$

\therefore Radius $XA \parallel$ radius YB (since alternate interior angles are equal)

Q. 8. In figure 3.88, circles with centres X and Y touch internally at point Z. Seg BZ is a chord of bigger circle and intersects smaller circle at point A. Prove that, $\text{seg } AX \parallel \text{seg } BY$.




Fig. 3.88

Answer : XA and YB are the radii of the respective circles.

AZ and BZ are the chords of the circles.

In triangle XAZ ,

$AX = XZ$ {Radii of the same circle}

$\Rightarrow \angle XAZ = \angle XZA$ {angles opposite to equal sides are equal}

In triangle YBZ ,

$YB = YZ$ {Radii of the same circle}

$\Rightarrow \angle YBZ = \angle YZB$ {angles opposite to equal sides are equal}

$\Rightarrow \angle XAZ = \angle XZA = \angle YBZ = \angle YZB$



∴ Corresponding angles are equal

⇒ $XA \parallel YB$

Q. 9. In figure 3.89, line l touches the circle with centre O at point P . Q is the mid point of radius OP . RS is a chord through Q such that chords $RS \parallel$ line l . If $RS = 12$ find the radius of the circle.




Fig. 3.89

Answer : The radius of the circle will bisect the chord RS . Therefore, $RQ = QS = \frac{1}{2} \times 12 = 6$

Let the radius of circle be r ,

Now, in ΔOQS , we have,

$$RQ = 6$$

$$OR = r$$

$$OQ = \frac{1}{2} r$$

Applying Pythagoras theorem, we get,

$$r^2 = \left(\frac{r}{2}\right)^2 + (6)^2$$

$$r^2 - \frac{r^2}{4} = 36$$

$$\frac{3r^2}{4} = 36$$

$$3r^2 = 4 \times 36$$

$$r^2 = 4 \times 12 = 48$$

$$r = \sqrt{48} \text{ units}$$

Q. 10. In figure 3.90, seg AB is a diameter of a circle with centre C. Line PQ is a tangent, which touches the circle at point T. seg AP \perp line PQ and seg BQ \perp line PQ. Prove that, seg CP \cong seg CQ.





Fig. 3.90

Answer : To Prove: seg CP \cong seg CQ

Construction: Join CP, CQ and CT

Figure:



Since PQ is a tangent to the circle, $\angle CTP = \angle CTQ = 90$

Since $\angle APT = \angle CTP = 90$

AP \parallel CT.

Similarly,

CT \parallel BQ.

So, we can say that,

AP \parallel CT \parallel BQ

AB is a line cutting all three parallel lines.

$AC = CB$ (Radius of the circle, and AB is diameter, C is center)

Since C is center point of line AB cutting parallel lines.

We can say these parallel lines are equal distance.

Therefore, $PT = TQ$.

Now in $\triangle CTP$ and $\triangle CTQ$,


CT is a common side, $PT = TQ$ and $\angle CTP = \angle CTQ = 90^\circ$

$CP = CQ$. (Pythagoras theorem or congruent triangle theorem)

Hence, Proved.

Q. 11. Draw circles with centres A , B and C each of radius 3 cm, such that each circle touches the other two circles.


Answer :



Draw a circle with radius 3 and centred at A . Similarly draw other two circles with centre B and C and same radius touching each other externally.

Q. 12. Prove that any three points on a circle cannot be collinear.

Answer :



We draw a circle of any radius and take any three points A, B and C on the circle.

We join A to B and B to C.

We draw perpendicular bisectors of AB and BC.

We know that perpendicular from the center bisects the chord.

Hence the center lies on both of the perpendicular bisectors.

The point where they intersect is the center of the circle.

The perpendiculars of the line segments drawn by joining collinear points is always parallel whereas in circle any three point's perpendicular bisector will always intersect at the center.

Hence, any three points on the circle cannot be collinear.

Q. 13. In figure 3.91, line PR touches the circle at point Q. Answer the following questions with the help of the figure.

- (1) What is the sum of $\angle TAQ$ and $\angle TSQ$?
- (2) Find the angles which are congruent to $\angle AQP$.
- (3) Which angles are congruent to $\angle QTS$?
- (4) $\angle TAS = 65^\circ$, find the measure of $\angle TQS$ and arc TS.

(5) If $\angle AQP = 42^\circ$ and $\angle SQR = 58^\circ$ find measure of $\angle ATS$.




Fig. 3.91

Answer : (1) As TAQS is a cyclic quadrilateral,

$\angle TAQ + \angle TSQ = 180^\circ$ (Sum of opposite angles of a cyclic quadrilateral is 180°)

(2) $\angle ASQ$ and $\angle ATQ$

(3) $\angle QAS$ and $\angle SQR$

(4) $\angle TAS = 65^\circ$

$\angle TQS = \angle TAS = 65^\circ$ (angle by same arc TS in the same sector)

$m(\text{arc TS}) = \angle TQS + \angle TAS$

$\Rightarrow m(\text{arc TS}) = 65 + 65 = 130^\circ$

(5) $\angle AQP + \angle AQS + \angle SQR = 180^\circ$

$\Rightarrow 42 + \angle AQS + 58 = 180$

$\Rightarrow \angle AQS + 100 = 180$

$\Rightarrow \angle AQS = 80$

$\angle AQS + \angle ATS = 180^\circ$ (opposite angles of a cyclic quadrilateral)

$\Rightarrow 80 + \angle ATS = 180$

$\Rightarrow \angle ATS = 100^\circ$

Q. 14. In figure 3.92, O is the centre of a circle, chord $PQ \cong$ chord RS If $\angle POR = 70^\circ$ and $(\text{arc RS}) = 80^\circ$, find -

(1) $m(\text{arc PR})$

- (2) $m(\text{arc QS})$
 (3) $m(\text{arc QSR})$




Fig. 3.92

Answer : (1) $m(\text{arc PR}) = \angle POR = 70^\circ$

$$(2) \angle POQ + \angle QOS + \angle ROS + \angle POR = 360^\circ$$

As $PQ = RS$, $\angle POQ = \angle ROS = 80^\circ$

$$\Rightarrow \angle POQ + \angle QOS + \angle ROS + \angle POR = 360^\circ$$

$$\Rightarrow 80 + \angle QOS + 80 + 70 = 360$$

$$\Rightarrow 230 + \angle QOS = 360$$

$$\Rightarrow \angle QOS = 130^\circ$$

$$m(\text{arc QS}) = \angle QOS = 130^\circ$$

$$(3) m(\text{arc QSR}) = \angle QOS + \angle ROS = 130 + 80 = 210^\circ$$

Q. 15. In figure 3.93, $m(\text{arc WY}) = 44^\circ$, $m(\text{arc ZX}) = 68^\circ$, then

- (1) Find the measure of $\angle ZTX$.
 (2) If $WT = 4.8$, $TX = 8.0$, $YT = 6.4$, find TZ .

(3) If $WX = 25$, $YT = 8$, $YZ = 26$, find WT .




Fig. 3.93

Answer : (1) Given: $m(\text{arc } WY) = 44^\circ$, $m(\text{arc } ZX) = 68^\circ$

We know that

$$\angle ZTX = \frac{1}{2} [m(\text{arc } ZX) + m(\text{arc } WX)]$$

$$\Rightarrow \angle ZTX = \frac{1}{2} (44^\circ + 68^\circ) = 56^\circ$$

(2) Given: $WT = 4.8$, $TX = 8.0$, $YT = 6.4$

We know that $WT \times TX = YT \times TZ$ {Using secant-tangent theorem}

$$\Rightarrow 6.4 \times TZ = 4.8 \times 8$$

$$\Rightarrow TZ = 6$$

(3) Given: $WX = 25$, $YT = 8$, $YZ = 26$

Let $WT = x$ and $TX = 25 - x$

$$WT \times TX = YT \times TZ$$

$$\Rightarrow x(25 - x) = 8 \times 26$$

$$\Rightarrow (x - 16)(x - 9) = 0$$

$$\Rightarrow WT = 16 \text{ or } 9$$

Q. 16. In figure 3.94,

(1) $m(\text{arc } CE) = 54^\circ$,

$m(\text{arc } BD) = 23^\circ$, find measure of $\angle CAE$.

(2) If $AB = 4.2$, $BC = 5.4$,
 $AE = 12.0$, find AD

(3) If $AB = 3.6$, $AC = 9.0$,
 $AD = 5.4$, find AE




Fig. 3.94

Answer : (1) Given: $m(\text{arc } CE) = 54^\circ$,

$m(\text{arc } BD) = 23^\circ$

$\angle CAE$ is an external angle.

$$\angle CAE = \frac{1}{2} [m(\text{arc } CE) - m(\text{arc } BD)]$$

$$\angle CAE = \frac{1}{2} [54^\circ - 23^\circ] = 15.5^\circ$$

(2) Given: $AB = 4.2$, $BC = 5.4$, $AE = 12.0$

Here, $AB \times AC = AD \times EA$

$$\Rightarrow AD \times 12 = 4.2 \times 5.4$$

$$\Rightarrow AD = 3.36$$

(3) Given $AB = 3.6$, $AC = 9.0$,

$AD = 5.4$

Here, $AB \times AC = AD \times EA$

$$\Rightarrow AE \times 5.4 = 3.6 \times 9$$

$$\Rightarrow AE = 6$$

Q. 17. In figure 3.95, chord $EF \parallel$ chord GH . Prove that, chord $EG \cong$ chord FH .

Fill in the blanks and write the proof.

Proof : Draw seg GF .

$\angle EFG = \angle FGH$...Alternate interior angles..... ☐ (I)

$\angle EFG = \dots\dots\dots 90^\circ$ {inscribed angle theorem}(II)

$\angle FGH = \dots\dots\dots 90^\circ$ {inscribed angle theorem} (III)

$\therefore m(\text{arc } EG) = \dots\dots\dots 90^\circ$ from (I), (II), (III).

chord $EG \cong$ chord FH ☐




Fig. 3.95

Answer : Proof : Draw seg GF .

$\angle EFG = \angle FGH$ {Alternate interior angles} (I)

$\angle EFG = 90^\circ$ {inscribed angle theorem}(II)

$\angle FGH = 90^\circ$ {inscribed angle theorem} (III)

$\therefore m(\text{arc } EG) = 90^\circ$ from (I), (II), (III).

Chord $EG \cong$ chord FH {Corresponding chords of congruent arcs of a circle (or congruent circles) are congruent}

Q. 18. In figure 3.96 P is the point of contact.

(1) If $m(\text{arc } PR) = 140^\circ$, $\angle POR = 36^\circ$, find $m(\text{arc } PQ)$

(2) If $OP = 7.2$, $OQ = 3.2$, find OR and QR

(3) If $OP = 7.2$, $OR = 16.2$, find QR .




Fig. 3.96

Answer : (1) Given: $m(\text{arc PR}) = 140^\circ$, $\angle POR = 36^\circ$

$\angle ROP$ is an external angle.

$$\angle ROP = \frac{1}{2}[m(\text{arc RP}) - m(\text{arc PQ})]$$

$$\Rightarrow m(\text{arc PQ}) = 140^\circ - 2 \times 36^\circ$$

$$\Rightarrow m(\text{arc PQ}) = 68^\circ$$

(2) Given: $OP = 7.2$, $OQ = 3.2$

Here, $RO \times OQ = OP^2$

$$\Rightarrow RO \times 3.2 = 7.2 \times 7.2$$

$$\Rightarrow RO = 16.2$$

$$QR = RO - OQ = 16.2 - 3.2 = 13$$

(3) Given: $OP = 7.2$, $OR = 16.2$

Here, $RO \times OQ = OP^2$

$$\Rightarrow 16.2 \times OQ = 7.2 \times 7.2$$

$$\Rightarrow OQ = 3.2$$

$$QR = RO - OQ = 16.2 - 3.2 = 13$$

Q. 19. In figure 3.97, circles with centres C and D touch internally at point E . D lies on the inner circle. Chord EB of the outer circle intersects inner circle at point A . Prove that, $\text{seg } EA \cong \text{seg } AB$.





Fig. 3.97

Answer :



We see that the line joining D to E passes through C.

In the smaller circle,

A lies in the semicircle,

$$\therefore \angle EAD = 90^\circ$$

\Rightarrow DA is perpendicular on the chord EB of the bigger circle.

We know that perpendicular from the center bisects the chord.

Therefore, EA = AB.

Q. 20. In figure 3.98, seg AB is a diameter of a circle with centre O . The bisector of $\angle ACB$ intersects the circle at point D. Prove that, seg AD \cong seg BD.




Fig. 3.98

Complete the following proof by filling in the blanks.

Proof: Draw seg OD.

$\angle ACB = \square$ 90° angle inscribed in semicircle

$\angle DCB = \square$ 45° CD is the bisector of $\angle C$

$m(\text{arc DB}) = \square$ 45° .. inscribed angle theorem

$\angle DOB = \square$ 90° definition of measure of an arc (I)

seg OA \cong seg OBradii of the circle... \square (II)

\therefore line OD is \square of seg ABbisector.... From (I) and (II)

\therefore seg AD \cong seg BD

Answer : Proof: Draw seg OD.

$\angle ACB = 90^\circ$ {angle inscribed in semicircle}

$\angle DCB = 45^\circ$ {CD is the bisector of $\angle C$ }

$m(\text{arc DB}) = 45^\circ$ {inscribed angle theorem}

$\angle DOB = 90^\circ$ {definition of measure of an arc} (I)

seg OA \cong seg OB {radii of the circle}(II)

\therefore line OD is bisector of seg AB From (I) and (II)

\therefore seg AD \cong seg BD

Q. 21. In figure 3.99, seg MN is a chord of a circle with centre O. $MN = 25$, L is a point on chord MN such that $ML = 9$ and $d(O, L) = 5$.

Find the radius of the circle.





Fig. 3.99

Answer : The figure is shown below:



Draw perpendicular on MN from the center O.

Mark the point as A. Join O to N.

As we know that perpendicular on a chord bisects the chord.

$$AM = MN/2$$

$$\Rightarrow AM = 25/2 = 12.5$$

Given that $LM = 9$

$$\Rightarrow LM + LA = AM$$

$$\Rightarrow 9 + LA = 12.5$$

$$\Rightarrow LA = 3.5$$

In ΔOAL ,

$$\Rightarrow OL^2 = OA^2 + AL^2$$

$$\Rightarrow 5^2 = OA^2 + (3.5)^2$$

$$\Rightarrow OA^2 = 25 - 12.25$$

$$\Rightarrow OA^2 = 12.75$$

In ΔOAN ,

$$\Rightarrow ON^2 = OA^2 + AN^2$$

$$\Rightarrow ON^2 = 12.75 + (12.5)^2$$

$$\Rightarrow ON^2 = 12.75 + 156.25$$

$$\Rightarrow ON^2 = 169$$

$$\Rightarrow ON = 13$$

Therefore, the radius of the circle is 13.

Q. 22. In figure 3.100, two circles intersect each other at points S and R. Their common tangent PQ touches the circle at points P, Q.

Prove that, $\angle PRQ + \angle PSQ = 180^\circ$




Fig. 3.100

Answer : We join R to S,




Fig. 3.100

As PQ is the tangent at P, we have

$$\angle RPQ = \angle PSR \dots\dots\dots(1)$$

As PQ is tangent at Q, we have

$$\angle RQP = \angle RSQ \dots\dots\dots(2)$$

In $\triangle RPQ$, we have

$$\Rightarrow \angle RPQ + \angle RQP + \angle PRQ = 180^\circ \text{ (Sum of all angles of a triangle)}$$

$$\Rightarrow \angle PSR + \angle RSQ + \angle PRQ = 180^\circ \text{ (From (1) and (2))}$$

$$\Rightarrow \angle PSQ + \angle PRQ = 180^\circ \text{ (}\angle PSR + \angle RSQ = \angle PSQ\text{)}$$

Hence Proved.

Q. 23. In figure 3.101, two circles intersect at points M and N. Secants drawn through M and N intersect the circles at points R, S and P, Q respectively.

Prove that :seg SQ || seg RP.





Fig. 3.101

Answer :



We join MN.

As PRMN is a cyclic quadrilateral,

$$\angle R + \angle PNM = 180^\circ \dots\dots\dots(1) \text{ (opposite angles of a cyclic quadrilateral)}$$

Also, QSMN is a cyclic quadrilateral,

$$\angle S + \angle QNM = 180^\circ \dots\dots\dots(2) \text{ (opposite angles of a cyclic quadrilateral)}$$

Adding (1) and (2)

$$\angle R + \angle S + \angle PNM + \angle QNM = 360^\circ$$

$$\Rightarrow \angle R + \angle S + 180 = 360 \text{ (PQ is a straight line)}$$

$$\Rightarrow \angle R + \angle S = 180^\circ$$

Similarly we have,

As PRMN is a cyclic quadrilateral,

$$\angle P + \angle RMN = 180^\circ \dots\dots\dots(3) \text{ (opposite angles of a cyclic quadrilateral)}$$

Also, QSMN is a cyclic quadrilateral,

$$\angle Q + \angle SMN = 180^\circ \dots\dots\dots(4) \text{ (opposite angles of a cyclic quadrilateral)}$$

Adding (3) and (4)

$$\angle P + \angle Q + \angle RMN + \angle SMN = 360^\circ$$

$$\Rightarrow \angle P + \angle Q + 180 = 360 \text{ (RS is a straight line)}$$

$$\Rightarrow \angle P + \angle Q = 180^\circ$$

Therefore, $PR \parallel SQ$.

Q. 24. In figure 3.102, two circles intersect each other at points A and E. Their common secant through E intersects the circles at points B and D. The tangents of the circles at points B and D intersect each other at point C.

Prove that $\square ABCD$ is cyclic.





Fig. 3.102

Answer : We join A to B and A to D and A to E



As BC is a tangent at B, we have

$$\angle CBD = \angle BAE \dots\dots\dots(1)$$

As CD is a tangent at D, we have

$$\angle CDB = \angle DAE \dots\dots\dots(2)$$

In $\triangle BCD$, we have

$$\Rightarrow \angle CBD + \angle CDB + \angle BCD = 180^\circ \text{ (Sum of all angles of a triangle)}$$

$$\Rightarrow \angle BAE + \angle DAE + \angle BCD = 180^\circ \text{ (From (1) and (2))}$$

$$\Rightarrow \angle BAD + \angle BCD = 180^\circ \text{ (}\angle BAE + \angle DAE = \angle BAD\text{)}$$

In quadrilateral ABCD,

We have $\angle A + \angle C = 180^\circ$ (Proved above)

$$\Rightarrow \angle A + \angle B + \angle C + \angle D = 360^\circ$$

$$\Rightarrow \angle B + \angle D + 180 = 360$$

$$\Rightarrow \angle B + \angle D = 180$$

Therefore, opposite angles of the quadrilateral sum to 180. Hence ABCD is a cyclic quadrilateral.

Q. 25. In figure 3.103, seg AD \perp side BC, seg BE \perp side AC, seg CF \perp side AB. Point O is the orthocentre. Prove that, point O is the incentre of $\triangle DEF$.





Fig. 3.103

Answer :



Join D to E, D to F and E to F.

In $\triangle ABE$,

$$\Rightarrow \angle ABE + \angle BAE + \angle BEA = 180^\circ \text{ (Sum of all angles of a triangle)}$$

$$\Rightarrow \angle ABE + \angle BAE + 90 = 180$$

$$\Rightarrow \angle ABE + \angle BAE = 90$$

$$\Rightarrow \angle ABO + \angle BAC = 90$$

$$\Rightarrow \angle ABO = 90 - \angle BAC \dots\dots\dots(1)$$

In quadrilateral BFOD, we have

$$\text{We have } \angle F = 90, \angle D = 90$$

$$\Rightarrow \angle B + \angle F + \angle O + \angle D = 360^\circ$$

$$\Rightarrow \angle B + \angle O + 180 = 360$$

$$\Rightarrow \angle B + \angle O = 180$$

Therefore, BFOD is a cyclic quadrilateral.

$$\angle FBO = \angle FDO \text{ (angle by the same arc)}$$

$$\Rightarrow \angle ABO = \angle FDO$$

From (1),

$$\Rightarrow \angle FDO = 90 - \angle BAC \dots\dots\dots(2)$$

In $\triangle AFC$,

$$\Rightarrow \angle CAF + \angle FCA + \angle AFC = 180^\circ \text{ (Sum of all angles of a triangle)}$$

$$\Rightarrow \angle CAF + \angle FCA + 90 = 180$$

$$\Rightarrow \angle CAF + \angle FCA = 90$$

$$\Rightarrow \angle BAC + \angle OCE = 90$$

$$\Rightarrow \angle OCE = 90 - \angle BAC \dots\dots\dots(3)$$

In quadrilateral CEOD, we have

We have $\angle E = 90$, $\angle D = 90$

$$\Rightarrow \angle C + \angle E + \angle O + \angle D = 360^\circ$$

$$\Rightarrow \angle C + \angle O + 180 = 360$$

$$\Rightarrow \angle C + \angle O = 180$$

Therefore, CEOD is a cyclic quadrilateral.

$\angle ODE = \angle OCE$ (angle by the same arc)

From (3),

$$\Rightarrow \angle ODE = 90 - \angle BAC \dots\dots\dots(4)$$

From (2) and (4) we conclude,

$$\angle FDO = \angle ODE$$

OD bisects $\angle D$.

Similarly, we can prove that OE bisects $\angle E$ and OF bisects $\angle F$.

Hence O is the incentre of $\triangle DEF$.