

Indices

PRACTICE SET 26 [PAGE 44]

Practice Set 26 | Q 1 | Page 44

Complete the table below.

Sr. No	Indices (Numbers in index form)	Base	Index	Multiplication form	Value
(i)	3^4	3	4	$3 \times 3 \times 3 \times 3$	81
(ii)	16^3				
(iii)		(-8)	2		
(iv)				$3/7 \times 3/7 \times 3/7 \times 3/7$	81/2401
(v)	$(-13)^4$				

Solution:

Sr. No	Indices (Numbers in index form)	Base	Index	Multiplication form	Value
(i)	3^4	3	4	$3 \times 3 \times 3 \times 3$	81
(ii)	16^3	16	3	$16 \times 16 \times 16$	4096
(iii)	$(-8)^2$	(-8)	2	$(-8) \times (-8)$	64
(iv)	$\left(\frac{3}{7}\right)^4$	$\frac{3}{7}$	4	$\frac{3}{7} \times \frac{3}{7} \times \frac{3}{7} \times \frac{3}{7}$	$\frac{81}{2401}$
(v)	$(-13)^4$	- 13	4	$(-13) \times (-13) \times (-13) \times (-13)$	28561

Practice Set 26 | Q 2.1 | Page 44

Find the value of 2^{10} .



Solution: $2^{10} = 2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2$
 $= 1024$

Practice Set 26 | Q 2.2 | Page 44

Find the value of 5^3 .

Solution: $5^3 = 5 \times 5 \times 5 = 125$

Practice Set 26 | Q 2.3 | Page 44

Find the value of $(-7)^4$

Solution: $(-7)^4 = (-7) \times (-7) \times (-7) \times (-7) = 2401$

Practice Set 26 | Q 2.4 | Page 44

Find the value of $(-6)^3$

Solution: $(-6)^3 = (-6) \times (-6) \times (-6) = -216$

Practice Set 26 | Q 2.5 | Page 44

Find the value of 9^3

Solution: $9^3 = 9 \times 9 \times 9 = 729$

Practice Set 26 | Q 2.6 | Page 44

Find the value of 8^1

Solution: $8^1 = 8$

Practice Set 26 | Q 2.7 | Page 44

Find the value of $\left(\frac{4}{5}\right)^3$

Solution:

$$\begin{aligned}\left(\frac{4}{5}\right)^3 &= \frac{4}{5} \times \frac{4}{5} \times \frac{4}{5} \\ &= \frac{64}{125}\end{aligned}$$

Practice Set 26 | Q 2.8 | Page 44

Find the value of $\left(-\frac{1}{2}\right)^4$

Solution:

$$\left(-\frac{1}{2}\right)^4 = -\frac{1}{2} \times -\frac{1}{2} \times -\frac{1}{2} \times -\frac{1}{2}$$

$$= \frac{1}{16}$$

PRACTICE SET 27 [PAGE 45]

Practice Set 27 | Q 1.1 | Page 45

Simplify: $7^4 \times 7^2$

Solution: It is known that, $a^m \times a^n = a^{m+n}$, where m and n are integers and a is a non-zero rational number.

$$7^4 \times 7^2$$

$$= 7^{4+2}$$

$$= 7^6$$

Practice Set 27 | Q 1.2 | Page 45

Simplify: $(-11)^5 \times (-11)^2$

Solution: It is known that, $a^m \times a^n = a^{m+n}$, where m and n are integers and a is a non-zero rational number.

$$(-11)^5 \times (-11)^2$$

$$= (-11)^{5+2}$$

$$= (-11)^7$$

Practice Set 27 | Q 1.3 | Page 45

Simplify: $\left(\frac{6}{7}\right)^3 \times \left(\frac{6}{7}\right)^5$

Solution: It is known that, $a^m \times a^n = a^{m+n}$, where m and n are integers and a is a non-zero rational number.

$$\begin{aligned} & \left(\frac{6}{7}\right)^3 \times \left(\frac{6}{7}\right)^5 \\ &= \left(\frac{6}{7}\right)^{3+5} \\ &= \left(\frac{6}{7}\right)^8 \end{aligned}$$

Practice Set 27 | Q 1.4 | Page 45

Simplify: $\left(-\frac{3}{2}\right)^5 \times \left(-\frac{3}{2}\right)^3$

Solution: It is known that, $a^m \times a^n = a^{m+n}$, where m and n are integers and a is a non-zero rational number.

$$\begin{aligned} & \left(-\frac{3}{2}\right)^5 \times \left(-\frac{3}{2}\right)^3 \\ &= \left(-\frac{3}{2}\right)^{5+3} \\ &= \left(-\frac{3}{2}\right)^8 \end{aligned}$$

Practice Set 27 | Q 1.5 | Page 45

Simplify: $a^{16} \times a^7$

Solution: It is known that, $a^m \times a^n = a^{m+n}$, where m and n are integers and a is a non-zero rational number.

$$a^{16} \times a^7$$

$$= a^{16+7}$$

$$= a^{23}$$

Practice Set 27 | Q 1.6 | Page 45

Simplify: $\left(\frac{P}{5}\right)^3 \times \left(\frac{P}{5}\right)^7$

Solution: It is known that, $a^m \times a^n = a^{m+n}$, where m and n are integers and a is a non-zero rational number.

$$\left(\frac{P}{5}\right)^3 \times \left(\frac{P}{5}\right)^7$$

$$= \left(\frac{P}{5}\right)^{3+7}$$

$$= \left(\frac{P}{5}\right)^{10}$$

PRACTICE SET 28 [PAGE 46]

Practice Set 28 | Q 1.1 | Page 46

Simplify: $a^6 \div a^4$

Solution: It is known that, $a^m \div a^n = a^{m-n}$, where m and n are integers and a is non-zero rational number.

$$a^6 \div a^4$$

$$= a^{6-4}$$

$$= a^2$$

Practice Set 28 | Q 1.2 | Page 46

Simplify: $m^5 \div m^8$

Solution: It is known that, $a^m \div a^n = a^{m-n}$, where m and n are integers and a is non-zero rational number.

$$m^5 \div m^8$$

$$= m^{5-8}$$

$$= m^{-3}$$

Practice Set 28 | Q 1.3 | Page 46

Simplify: $p^3 \div p^{13}$



Solution: It is known that, $a^m \div a^n = a^{m-n}$, where m and n are integers and a is non-zero rational number.

$$\begin{aligned} p^3 \div p^{13} \\ &= p^{3-13} \\ &= p^{-10} \end{aligned}$$

Practice Set 28 | Q 1.4 | Page 46

Simplify: $x^{10} \div x^{10}$

Solution: It is known that, $a^m \div a^n = a^{m-n}$, where m and n are integers and a is non-zero rational number.

$$\begin{aligned} x^{10} \div x^{10} \\ &= x^{10-10} \\ &= x^0 \\ &= 1 \quad (\because a^0 = 1) \end{aligned}$$

Practice Set 28 | Q 2.1 | Page 46

Find the value.

$$(-7)^{12} \div (-7)^{12}$$

Solution: It is known that, $a^m \div a^n = a^{m-n}$, where m and n are integers and a is a non-zero rational number.

$$\begin{aligned} (-7)^{12} \div (-7)^{12} \\ &= (-7)^{12-12} \\ &= (-7)^0 \\ &= 1 \quad (\because a^0 = 1) \end{aligned}$$

Practice Set 28 | Q 2.2 | Page 46

Find the value.

$$7^5 \div 7^3$$

Solution: It is known that, $a^m \div a^n = a^{m-n}$, where m and n are integers and a is a non-zero rational number.

$$7^5 \div 7^3$$

$$= 7^{5-3}$$

$$= 7^2$$

$$= 7 \times 7$$

$$= 49$$

Practice Set 28 | Q 2.3 | Page 46

Find the value.

$$\left(\frac{4}{5}\right)^3 \div \left(\frac{4}{5}\right)^2$$

Solution: It is known that, $a^m \div a^n = a^{m-n}$, where m and n are integers and a is a non-zero rational number.

$$\left(\frac{4}{5}\right)^3 \div \left(\frac{4}{5}\right)^2$$

$$= \left(\frac{4}{5}\right)^{3-2}$$

$$= \left(\frac{4}{5}\right)^1$$

$$= \frac{4}{5}$$

Practice Set 28 | Q 2.4 | Page 46

Find the value.

$$4^7 \div 4^5$$

Solution: It is known that, $a^m \div a^n = a^{m-n}$, where m and n are integers and a is a non-zero rational number.

$$4^7 \div 4^5$$

$$= 4^{7-5}$$

$$= 4^2$$

$$= 4 \times 4$$

$$= 16$$



PRACTICE SET 29 [PAGE 48]

Practice Set 29 | Q 1.01 | Page 48

Simplify: $\left[\left(\frac{15}{12}\right)^3\right]^4$

Solution: It is known that, $(a^m)^n = a^{mn}$, where m and n are integers and a is a non-zero rational number.

$$\begin{aligned} & \left[\left(\frac{15}{12}\right)^3\right]^4 \\ &= \left(\frac{15}{12}\right)^{3 \times 4} \\ &= \left(\frac{15}{12}\right)^{12} \end{aligned}$$

Practice Set 29 | Q 1.02 | Page 48

Simplify: $(3^4)^{-2}$

Solution: It is known that, $(a^m)^n = a^{mn}$, where m and n are integers and a is a non-zero rational number.

$$\begin{aligned} & (3^4)^{-2} \\ &= 3^4 \times (-2) \\ &= 3^{-8} \end{aligned}$$

Practice Set 29 | Q 1.03 | Page 48

Simplify: $\left(\left(\frac{1}{7}\right)^{-3}\right)^4$

Solution: It is known that, $(a^m)^n = a^{mn}$, where m and n are integers and a is a non-zero rational number.

$$\begin{aligned} & \left(\left(\frac{1}{7} \right)^{-3} \right)^4 \\ &= \left(\frac{1}{7} \right)^{-3 \times 4} \\ &= \left(\frac{1}{7} \right)^{-12} \end{aligned}$$

Practice Set 29 | Q 1.04 | Page 48

Simplify: $\left(\left(\frac{2}{5} \right)^{-2} \right)^{-3}$

Solution: It is known that, $(a^m)^n = a^{mn}$, where m and n are integers and a is a non-zero rational number.

$$\begin{aligned} & \left(\left(\frac{2}{5} \right)^{-2} \right)^{-3} \\ &= \left(\frac{2}{5} \right)^{(-2) \times (-3)} \\ &= \left(\frac{2}{5} \right)^6 \end{aligned}$$

Practice Set 29 | Q 1.05 | Page 48

Simplify: $(6^5)^4$

Solution: It is known that, $(a^m)^n = a^{mn}$, where m and n are integers and a is a non-zero rational number.

$$\begin{aligned}
 &(6^5)^4 \\
 &= 6^{5 \times 4} \\
 &= 6^{20}
 \end{aligned}$$

Practice Set 29 | Q 1.06 | Page 48

Simplify: $\left[\left(\frac{6}{7}\right)^5\right]^2$

Solution: It is known that, $(a^m)^n = a^{mn}$, where m and n are integers and a is a non-zero rational number.

$$\begin{aligned}
 &\left[\left(\frac{6}{7}\right)^5\right]^2 \\
 &= \left(\frac{6}{7}\right)^{5 \times 2} \\
 &= \left(\frac{6}{7}\right)^{10}
 \end{aligned}$$

Practice Set 29 | Q 1.07 | Page 48

Simplify: $\left[\left(\frac{2}{3}\right)^{-4}\right]^5$

Solution: It is known that, $(a^m)^n = a^{mn}$, where m and n are integers and a is a non-zero rational number.

$$\begin{aligned} & \left[\left(\frac{2}{3} \right)^{-4} \right]^5 \\ &= \left(\frac{2}{3} \right)^{-4 \times 5} \\ &= \left(\frac{2}{3} \right)^{-20} \end{aligned}$$

Practice Set 29 | Q 1.08 | Page 48

Simplify: $\left[\left(\frac{5}{8} \right)^3 \right]^{-2}$

Solution: It is known that, $(a^m)^n = a^{mn}$, where m and n are integers and a is a non-zero rational number.

$$\begin{aligned} & \left[\left(\frac{5}{8} \right)^3 \right]^{-2} \\ &= \left(\frac{5}{8} \right)^{3 \times (-2)} \\ &= \left(\frac{5}{8} \right)^{-6} \end{aligned}$$

Practice Set 29 | Q 1.09 | Page 48

Simplify: $\left[\left(\frac{3}{4} \right)^6 \right]^1$

Solution: It is known that, $(a^m)^n = a^{mn}$, where m and n are integers and a is a non-zero rational number.

$$\begin{aligned} & \left[\left(\frac{3}{4} \right)^6 \right]^1 \\ &= \left(\frac{3}{4} \right)^{6 \times 1} \\ &= \left(\frac{3}{4} \right)^6 \end{aligned}$$

Practice Set 29 | Q 1.1 | Page 48

Simplify: $\left[\left(\frac{2}{5} \right)^{-3} \right]^2$

Solution: It is known that, $(a^m)^n = a^{mn}$, where m and n are integers and a is a non-zero rational number.

$$\begin{aligned} & \left[\left(\frac{2}{5} \right)^{-3} \right]^2 \\ &= \left(\frac{2}{5} \right)^{-3 \times 2} \\ &= \left(\frac{2}{5} \right)^{-6} \end{aligned}$$

Practice Set 29 | Q 2.1 | Page 48

Write the following numbers using positive indices.

$$\left(\frac{2}{7} \right)^{-2}$$

Solution:

It is known that, $a^{-m} = \frac{1}{a^m}$ where m is an integer and a is a non-zero rational number.

$$\begin{aligned} & \left(\frac{2}{7}\right)^{-2} \\ &= \frac{1}{\left(\frac{2}{7}\right)^2} \\ &= \left(\frac{7}{2}\right)^2 \end{aligned}$$

Practice Set 29 | Q 2.2 | Page 48

Write the following numbers using positive indices.

$$\left(\frac{11}{3}\right)^{-5}$$

Solution:

It is known that, $a^{-m} = \frac{1}{a^m}$ where m is an integer and a is a non-zero rational number.

$$\begin{aligned} & \left(\frac{11}{3}\right)^{-5} \\ &= \frac{1}{\left(\frac{11}{3}\right)^5} \\ &= \left(\frac{3}{11}\right)^5 \end{aligned}$$

Practice Set 29 | Q 2.3 | Page 48

Write the following numbers using positive indices.

$$\left(\frac{1}{6}\right)^{-3}$$

Solution:

It is known that, $a^{-m} = \frac{1}{a^m}$ where m is an integer and a is a non-zero rational number.

$$\begin{aligned} &\left(\frac{1}{6}\right)^{-3} \\ &= \frac{1}{\left(\frac{1}{6}\right)^3} \\ &= \left(\frac{6}{1}\right)^3 \\ &= 6^3 \end{aligned}$$

Practice Set 29 | Q 2.4 | Page 48

Write the following numbers using positive indices.

$$(y)^{-4}$$

Solution:

It is known that, $a^{-m} = \frac{1}{a^m}$ where m is an integer and a is a non-zero rational number.

$$\begin{aligned} &(y)^{-4} \\ &= \frac{1}{y^4} \\ &= \left(\frac{1}{y}\right)^4 \end{aligned}$$

PRACTICE SET 30 [PAGE 50]

Practice Set 30 | Q 1 | Page 50

Find the square root of 625.

Solution: The prime factorization of 625 is,

$$625 = \underline{5 \times 5} \times \underline{5 \times 5}$$

To find the square root, we will take one number from each pair and multiply.

$$\sqrt{625} = 5 \times 5 = 25$$

$$\sqrt{625} = 25$$

Practice Set 30 | Q 2 | Page 50

Find the square root of 1225.

Solution: The prime factorization of 1225 is,

$$1225 = \underline{5 \times 5} \times \underline{7 \times 7}$$

To find the square root, we will take one number from each pair and multiply.

$$\sqrt{1225} = 5 \times 7 = 35$$

$$\sqrt{1225} = 35$$

Practice Set 30 | Q 3 | Page 50

Find the square root of 289.

Solution: The prime factorization of 289 is,

$$289 = \underline{17 \times 17}$$

To find the square root, we will take one number from each pair and multiply.

$$\sqrt{289} = 17 \times 17 = 17$$

$$\sqrt{289} = 17$$

Practice Set 30 | Q 4 | Page 50

Find the square root of 4096.

Solution: The prime factorization of 4096 is,

$$4096 = \underline{2 \times 2} \times \underline{2 \times 2} \times \underline{2 \times 2} \times \underline{2 \times 2} \times \underline{2 \times 2} \times \underline{2 \times 2}$$

To find the square root, we will take one number from each pair and multiply.

$$\sqrt{4096} = 2 \times 2 \times 2 \times 2 \times 2 \times 2 = 64$$

$$\sqrt{4096} = 64$$

Practice Set 30 | Q 5 | Page 50

Find the square root of 1089.

Solution: The prime factorization of 1089 is,

$$1089 = \underline{3 \times 3} \times \underline{11 \times 11}$$

To find the square root, we will take one number from each pair and multiply.

$$\sqrt{1089} = 3 \times 11 = 33$$

$$\sqrt{1089} = 33$$

