

Class: XII Session: 2020-21
Subject: Mathematics
Sample Question Paper (Theory)

Time Allowed: 3 Hours

Maximum Marks: 80

General Instructions:

1. This question paper contains two **parts A and B**. Each part is compulsory. Part A carries **24** marks and Part B carries **56** marks
2. **Part-A** has Objective Type Questions and **Part -B** has Descriptive Type Questions
3. Both Part A and Part B have choices.

Part – A:

1. It consists of two sections- **I and II**.
2. Section **I** comprises of 16 very short answer type questions.
3. Section **II** contains **2** case studies. Each case study comprises of 5 case-based MCQs. An examinee is to attempt **any 4 out of 5 MCQs**.

Part – B:



1. It consists of three sections- **III, IV and V**.
2. Section **III** comprises of 10 questions of **2 marks** each.
3. Section **IV** comprises of 7 questions of **3 marks** each.
4. Section **V** comprises of 3 questions of **5 marks** each.
5. Internal choice is provided in **3** questions of Section –III, **2** questions of Section-IV and **3** questions of Section-V. You have to attempt only one of the alternatives in all such questions.

Sr. No.	Part – A	Marks
	Section I All questions are compulsory. In case of internal choices attempt any one.	
1	Check whether the function $f: R \rightarrow R$ defined as $f(x) = x^3$ is one-one or not. OR	1



	How many reflexive relations are possible in a set A whose $n(A) = 3$.	1
2	A relation R in $S = \{1,2,3\}$ is defined as $R = \{(1,1), (1,2), (2,2), (3,3)\}$. Which element(s) of relation R be removed to make R an equivalence relation?	1
3	A relation R in the set of real numbers $\mathbf{R}$ defined as $R = \{(a,b): \sqrt{a} = b\}$ is a function or not. Justify OR An equivalence relation R in A divides it into equivalence classes A_1, A_2, A_3 . What is the value of $A_1 \cup A_2 \cup A_3$ and $A_1 \cap A_2 \cap A_3$	1 1
4	If A and B are matrices of order $3 \times n$ and $m \times 5$ respectively, then find the order of matrix $5A - 3B$, given that it is defined.	1
5	Find the value of A^2 , where A is a 2×2 matrix whose elements are given by $a_{ij} = \begin{cases} 1 & \text{if } i \neq j \\ 0 & \text{if } i = j \end{cases}$ OR Given that A is a square matrix of order 3×3 and $ A = -4$. Find $ \text{adj } A $	1 1
6	Let $A = [a_{ij}]$ be a square matrix of order 3×3 and $ A = -7$. Find the value of $a_{11} A_{21} + a_{12} A_{22} + a_{13} A_{23}$ where A_{ij} is the cofactor of element a_{ij}	1
7	Find $\int e^x (1 - \cot x + \operatorname{cosec}^2 x) dx$ OR Evaluate $\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} x^2 \sin x dx$	1 1
8	Find the area bounded by $y = x^2$, the x-axis and the lines $x = -1$ and $x = 1$.	1
9	How many arbitrary constants are there in the particular solution of the differential equation $\frac{dy}{dx} = -4xy^2$; $y(0) = 1$ OR For what value of n is the following a homogeneous differential equation: $\frac{dy}{dx} = \frac{x^3 - y^n}{x^2 y + xy^2}$	1 1
10	Find a unit vector in the direction opposite to $-\frac{3}{4} \hat{j}$	1
11	Find the area of the triangle whose two sides are represented by the vectors $2\hat{i}$ and $-3\hat{j}$.	1



12	Find the angle between the unit vectors $\hat{a}$ and $\hat{b}$, given that $ \hat{a} + \hat{b} = 1$	1
13	Find the direction cosines of the normal to YZ plane?	1
14	Find the coordinates of the point where the line $\frac{x+3}{3} = \frac{y-1}{-1} = \frac{z-5}{-5}$ cuts the XY plane.	1
15	The probabilities of A and B solving a problem independently are $\frac{1}{3}$ and $\frac{1}{4}$ respectively. If both of them try to solve the problem independently, what is the probability that the problem is solved?	1
16	The probability that it will rain on any particular day is 50%. Find the probability that it rains only on first 4 days of the week.	1
	Section II Both the Case study based questions are compulsory. Attempt any 4 sub parts from each question (17-21) and (22-26). Each question carries 1 mark	
17	An architect designs a building for a multi-national company. The floor consists of a rectangular region with semicircular ends having a perimeter of 200m as shown below: Design of Floor  Building  Based on the above information answer the following:	
	(i) If x and y represents the length and breadth of the rectangular region, then the relation between the variables is a) $x + \pi y = 100$ b) $2x + \pi y = 200$ c) $\pi x + y = 50$ d) $x + y = 100$	

	(ii) The area of the rectangular region A expressed as a function of x is a) $\frac{2}{\pi} (100x - x^2)$ b) $\frac{1}{\pi} (100x - x^2)$ c) $\frac{x}{\pi} (100 - x)$ d) $\pi y^2 + \frac{2}{\pi} (100x - x^2)$	1
	(iii) The maximum value of area A is a) $\frac{\pi}{3200} m^2$ b) $\frac{3200}{\pi} m^2$ c) $\frac{5000}{\pi} m^2$ d) $\frac{1000}{\pi} m^2$	1
	(iv) The CEO of the multi-national company is interested in maximizing the area of the whole floor including the semi-circular ends. For this to happen the value of x should be a) 0 m b) 30 m c) 50 m d) 80 m	1
	(v) The extra area generated if the area of the whole floor is maximized is : a) $\frac{3000}{\pi} m^2$ b) $\frac{5000}{\pi} m^2$ c) $\frac{7000}{\pi} m^2$ d) No change Both areas are equal	1




18	In an office three employees Vinay, Sonia and Iqbal process incoming copies of a certain form. Vinay process 50% of the forms. Sonia processes 20% and Iqbal the remaining 30% of the forms. Vinay has an error rate of 0.06, Sonia has an error rate of 0.04 and Iqbal has an error rate of 0.03  Based on the above information answer the following:	
	(i) The conditional probability that an error is committed in processing given that Sonia processed the form is : a) 0.0210 b) 0.04 c) 0.47 d) 0.06	1
	(ii) The probability that Sonia processed the form and committed an error is : a) 0.005 b) 0.006 c) 0.008 d) 0.68	1
	(iii) The total probability of committing an error in processing the form is a) 0 b) 0.047 c) 0.234	1

	d) 1	
	(iv) The manager of the company wants to do a quality check. During inspection he selects a form at random from the days output of processed forms. If the form selected at random has an error, the probability that the form is NOT processed by Vinay is : a) 1 b) 30/47 c) 20/47 d) 17/47	1
	(v) Let A be the event of committing an error in processing the form and let E_1, E_2 and E_3 be the events that Vinay, Sonia and Iqbal processed the form. The value of $\sum_{i=1}^3 P(E_i A)$ is a) 0 b) 0.03 c) 0.06 d) 1	1
	Part – B	
	Section III	
19	Express $\tan^{-1}\left(\frac{\cos x}{1 - \sin x}\right)$, $\frac{-3\pi}{2} < x < \frac{\pi}{2}$ in the simplest form.	2
20	If A is a square matrix of order 3 such that $A^2 = 2A$, then find the value of A . OR If $A = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix}$, show that $A^2 - 5A + 7I = 0$. Hence find A^{-1}.	2 2
21	Find the value(s) of k so that the following function is continuous at $x = 0$	2



	$f(x) = \begin{cases} \frac{1-\cos kx}{x \sin x} & \text{if } x \neq 0 \\ \frac{1}{2} & \text{if } x = 0 \end{cases}$	
22	Find the equation of the normal to the curve $y = x + \frac{1}{x}$, $x > 0$ perpendicular to the line $3x - 4y = 7$.	2
23	Find $\int \frac{1}{\cos^2 x (1 - \tan x)^2} dx$ OR Evaluate $\int_0^1 x(1-x)^n dx$	2 2
24	Find the area of the region bounded by the parabola $y^2 = 8x$ and the line $x = 2$.	2
25	Solve the following differential equation: $\frac{dy}{dx} = x^3 \operatorname{cosec} y$, given that $y(0) = 0$.	2
26	Find the area of the parallelogram whose one side and a diagonal are represented by coinitial vectors $\hat{i} - \hat{j} + \hat{k}$ and $4\hat{i} + 5\hat{k}$ respectively	2
27	Find the vector equation of the plane that passes through the point $(1,0,0)$ and contains the line $\vec{r} = \lambda \hat{j}$.	2
28	A refrigerator box contains 2 milk chocolates and 4 dark chocolates. Two chocolates are drawn at random. Find the probability distribution of the number of milk chocolates. What is the most likely outcome? OR Given that E and F are events such that $P(E) = 0.8$, $P(F) = 0.7$, $P(E \cap F) = 0.6$. Find $P(\bar{E} \bar{F})$	2 2
	Section IV All questions are compulsory. In case of internal choices attempt any one.	
29	Check whether the relation R in the set Z of integers defined as $R = \{(a, b) : a + b \text{ is "divisible by 2"}\}$ is reflexive, symmetric or transitive. Write the equivalence class containing 0 i.e. $[0]$.	3
30	If $y = e^{x \sin^2 x} + (\sin x)^x$, find $\frac{dy}{dx}$.	3
31	Prove that the greatest integer function defined by $f(x) = [x]$, $0 < x < 2$ is not differentiable at $x = 1$	3



	$2x + 3y + 4z = 17$ $y + 2z = 7$	
37	Find the shortest distance between the lines $\vec{r} = 3\hat{i} + 2\hat{j} - 4\hat{k} + \lambda(\hat{i} + 2\hat{j} + 2\hat{k})$ and $\vec{r} = 5\hat{i} - 2\hat{j} + \mu(3\hat{i} + 2\hat{j} + 6\hat{k})$ If the lines intersect find their point of intersection OR Find the foot of the perpendicular drawn from the point $(-1, 3, -6)$ to the plane $2x + y - 2z + 5 = 0$. Also find the equation and length of the perpendicular.	5
38	Solve the following linear programming problem (L.P.P) graphically. Maximize $Z = x + 2y$ subject to constraints ; $x + 2y \geq 100$ $2x - y \leq 0$ $2x + y \leq 200$ $x, y \geq 0$ OR The corner points of the feasible region determined by the system of linear constraints are as shown below:  Answer each of the following: (i) Let $Z = 3x - 4y$ be the objective function. Find the maximum and minimum value of Z and also the corresponding points at which the maximum and minimum value occurs.	5

	(ii) Let $Z = px + qy$, where $p, q > 0$ be the objective function. Find the condition on p and q so that the maximum value of Z occurs at $B(4,10)$ and $C(6,8)$. Also mention the number of optimal solutions in this case.	
--	--	--



Class: XII Session: 2020-21

Subject: Mathematics

Marking Scheme (Theory)

Sr.No.	Objective type Question Section I	Marks
1	Let $f(x_1) = f(x_2)$ for some $x_1, x_2 \in R$ $\Rightarrow (x_1)^3 = (x_2)^3$ $\Rightarrow x_1 = x_2$, Hence $f(x)$ is one – one OR 2^6 reflexive relations	1 1
2	(1,2)	1
3	Since $\sqrt{a}$ is not defined for $a \in (-\infty, 0)$ $\therefore \sqrt{a} = b$ is not a function. OR $A_1 \cup A_2 \cup A_3 = A$ and $A_1 \cap A_2 \cap A_3 = \phi$	1 1
4	3×5	1
5	$A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \Rightarrow A^2 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ OR $ \text{adj } A = (-4)^{3-1} = 16$	1
6	0	1
7	$e^x(1 - \cot x) + C$ OR $\therefore f(x)$ is an odd function $\therefore \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} x^2 \sin x \, dx = 0$	1 1
8	$A = 2 \int_0^1 x^2 \, dx = \frac{2}{3} [x^3]_0^1$ $= \frac{2}{3} \text{sq unit}$	1



9	0 OR 3	1 1
10	$\hat{j}$	1
11	$\frac{1}{2} 2\hat{i} \times (-3\hat{j}) = \frac{1}{2} -6\hat{k} = 3 \text{ sq units}$	1
12	$ \hat{a} + \hat{b} ^2 = 1$ $\Rightarrow \hat{a}^2 + \hat{b}^2 + 2\hat{a} \cdot \hat{b} = 1$ $\Rightarrow 2\hat{a} \cdot \hat{b} = 1 - 1 - 1$ $\Rightarrow \hat{a} \cdot \hat{b} = \frac{-1}{2} \Rightarrow \hat{a} \hat{b} \cos \theta = \frac{-1}{2} \Rightarrow \theta = \pi - \frac{\pi}{3}$ $\Rightarrow \theta = \frac{2\pi}{3}$	1
13	1,0,0	1
14	(0,0,0)	1
15	$1 - \frac{2}{3} \times \frac{3}{4} = \frac{1}{2}$	1
16	$\left(\frac{1}{2}\right)^4 \left(\frac{1}{2}\right)^3 = \left(\frac{1}{2}\right)^7$	1
Section II		
17(i)	(b)	1
17(ii)	(a)	1
17(iii)	(c)	1
17(iv)	(a)	1
17(v)	(d)	1
18(i)	(b)	1
18(ii)	(c)	1
18(iii)	(b)	1
18(iv)	(d)	1
18(v)	(d)	1
Section III		
19	$\tan^{-1}\left(\frac{\cos x}{1-\sin x}\right) = \tan^{-1}\left[\frac{\sin\left(\frac{\pi}{2}-x\right)}{1-\cos\left(\frac{\pi}{2}-x\right)}\right]$ $\tan^{-1}\left[\frac{2\sin\left(\frac{\pi}{4}-\frac{x}{2}\right)\cos\left(\frac{\pi}{4}-\frac{x}{2}\right)}{2\sin^2\left(\frac{\pi}{4}-\frac{x}{2}\right)}\right]$	$\frac{1}{2}$



	$\tan^{-1} \left[\cot \left(\frac{\pi}{4} - \frac{x}{2} \right) \right] = \tan^{-1} \left[\tan \frac{\pi}{2} - \left(\frac{\pi}{4} - \frac{x}{2} \right) \right]$ $\tan^{-1} \left[\tan \left(\frac{\pi}{4} + \frac{x}{2} \right) \right] = \frac{\pi}{4} + \frac{x}{2}$	1 $\frac{1}{2}$
20	$A^2 = 2A$ $\Rightarrow AA = 2A $ $\Rightarrow A A = 8 A \quad (\because AB = A B \text{ and } 2A = 2^3 A)$ $\Rightarrow A (A - 8) = 0$ $\Rightarrow A = 0 \text{ or } 8$ OR $A^2 = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix} = \begin{bmatrix} 8 & 5 \\ -5 & 3 \end{bmatrix}$ $5A = \begin{bmatrix} 15 & 5 \\ -5 & 10 \end{bmatrix}, 7I = \begin{bmatrix} 7 & 0 \\ 0 & 7 \end{bmatrix}$ $\Rightarrow A^2 - 5A + 7I = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} = 0$ $\Rightarrow A^{-1}(A^2 - 5A + 7I) = A^{-1}0$ $\Rightarrow A - 5I + 7A^{-1} = 0$ $\Rightarrow 7A^{-1} = 5I - A$ $\Rightarrow A^{-1} = \frac{1}{7} \left(\begin{bmatrix} 5 & 0 \\ 0 & 5 \end{bmatrix} - \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix} \right)$ $\Rightarrow A^{-1} = \frac{1}{7} \begin{bmatrix} 2 & -1 \\ 1 & 3 \end{bmatrix}$	$\frac{1}{2}$ $\frac{1}{2}$ 1 1
21	$\lim_{x \rightarrow 0} \frac{1 - \cos kx}{x \sin x} = \lim_{x \rightarrow 0} \frac{2 \sin^2 \left(\frac{kx}{2} \right)}{x \sin x}$ $= \lim_{x \rightarrow 0} \frac{2 \sin^2 \left(\frac{kx}{2} \right)}{\frac{x^2}{\frac{x \sin x}{x^2}}}$ $= \frac{\lim_{x \rightarrow 0} \frac{2 \sin^2 \left(\frac{kx}{2} \right)}{\left(\frac{kx}{2} \right)^2} \times \left(\frac{k}{2} \right)^2}{\lim_{x \rightarrow 0} \frac{\sin x}{x}} = \frac{2 \times 1 \times \frac{k^2}{4}}{1}$	$1 \frac{1}{2}$

	$= 4\sqrt{2} \left[\frac{2}{3} x^{\frac{3}{2}} \right]_0^2$ $= \frac{8}{3} \sqrt{2} \left[2^{\frac{3}{2}} - 0 \right] = \frac{8\sqrt{2}}{3} \times 2\sqrt{2}$ $= \frac{32}{3} \text{ sq units}$	$\frac{1}{2}$ $\frac{1}{2}$		
25	$\frac{dy}{dx} = x^3 \operatorname{cosec} y \quad ; \quad y(0) = 0$ $\int \frac{dy}{\operatorname{cosec} y} = \int x^3 dx$ $\int \sin y \, dy = \int x^3 dx$ $-\cos y = \frac{x^4}{4} + c$ $-1 = c \quad (\because y = 0, \text{ when } x = 0)$ $\cos y = 1 - \frac{x^4}{4}$	$\frac{1}{2}$ 1 $\frac{1}{2}$		
26	Let $\vec{a} = \hat{i} - \hat{j} + \hat{k}$ $\vec{d} = 4\hat{i} + 5\hat{k}$ $\therefore \vec{a} + \vec{b} = \vec{d} \therefore \vec{b} = \vec{d} - \vec{a} = 3\hat{i} + \hat{j} + 4\hat{k}$ $\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & -1 & 1 \\ 3 & 1 & 4 \end{vmatrix} = -5\hat{i} - 1\hat{j} + 4\hat{k}$ Area of parallelogram $= \vec{a} \times \vec{b} = \sqrt{25 + 1 + 16} = \sqrt{42} \text{ sq units}$	$\frac{1}{2}$ 1 $\frac{1}{2}$		
27	Let the normal vector to the plane be $\vec{n}$ Equation of the plane passing through (1,0,0), i.e., $\hat{i}$ is $(\vec{r} - \hat{i}) \cdot \vec{n} = 0 \dots\dots\dots(1)$ $\therefore$ plane (1) contains the line $\vec{r} = \vec{o} + \lambda \hat{j}$ $\therefore \hat{i} \cdot \vec{n} = 0$ and $\hat{j} \cdot \vec{n} = 0 \Rightarrow \vec{n} = \hat{k}$ Hence equation of the plane is $(\vec{r} - \hat{i}) \cdot \hat{k} = 0$ i.e., $\vec{r} \cdot \hat{k} = 0$	1 1		
28	Let x denote the number of milk chocolates drawn <table border="1" style="margin-left: auto; margin-right: auto;"><tr><td>X</td><td>P(x)</td></tr></table>	X	P(x)	
X	P(x)			



	<table><tr><td>0</td><td>$\frac{4}{6} \times \frac{3}{5} = \frac{12}{30}$</td></tr><tr><td>1</td><td>$\left(\frac{2}{6} \times \frac{4}{5}\right) \times 2 = \frac{16}{30}$</td></tr><tr><td>2</td><td>$\frac{2}{6} \times \frac{1}{5} = \frac{2}{30}$</td></tr></table>	0	$\frac{4}{6} \times \frac{3}{5} = \frac{12}{30}$	1	$\left(\frac{2}{6} \times \frac{4}{5}\right) \times 2 = \frac{16}{30}$	2	$\frac{2}{6} \times \frac{1}{5} = \frac{2}{30}$	$1\frac{1}{2}$
0	$\frac{4}{6} \times \frac{3}{5} = \frac{12}{30}$							
1	$\left(\frac{2}{6} \times \frac{4}{5}\right) \times 2 = \frac{16}{30}$							
2	$\frac{2}{6} \times \frac{1}{5} = \frac{2}{30}$							
	Most likely outcome is getting one chocolate of each type OR $P(\bar{E} \bar{F}) = P\left(\frac{\bar{E} \cap \bar{F}}{P(\bar{F})}\right) = \frac{P(\bar{E} \cup \bar{F})}{P(\bar{F})} = \frac{1 - P(E \cup F)}{1 - P(F)} \text{-----(1)}$ Now $P(E \cup F) = P(E) + P(F) - P(E \cap F)$ $= 0.8 + 0.7 - 0.6 = 0.9$ Substituting value of $P(E \cup F)$ in (1) $P(\bar{E} \bar{F}) = \frac{1 - 0.9}{1 - 0.7} = \frac{0.1}{0.3} = \frac{1}{3}$	$\frac{1}{2}$ $\frac{1}{2}$						
	Section IV							
29	(i) Reflexive : Since, $a+a=2a$ which is even $\therefore (a,a) \in R \forall a \in Z$ Hence R is reflexive (ii) Symmetric: If $(a,b) \in R$, then $a+b = 2\lambda \Rightarrow b+a = 2\lambda$ $\Rightarrow (b,a) \in R$, Hence R is symmetric (iii) Transitive: If $(a,b) \in R$ and $(b,c) \in R$ then $a+b = 2\lambda$---(1) and $b+c = 2\mu$ ---- (2) Adding (1) and (2) we get $a+2b+c=2(\lambda + \mu)$ $\Rightarrow a+c=2(\lambda + \mu - b)$ $\Rightarrow a+c=2k$, where $\lambda + \mu - b = k \Rightarrow (a,c) \in R$ Hence R is transitive $[0] = \{...-4, -2, 0, 2, 4...\}$	$\frac{1}{2}$ $\frac{1}{2}$						
30	Let $u = e^{x \sin^2 x}$ and $v = (\sin x)^x$	$\frac{1}{2}$						





	so that $y = u + v \Rightarrow \frac{dy}{dx} = \frac{du}{dx} + \frac{dv}{dx}$-----(1) Now, $u = e^{x \sin^2 x}$, Differentiating both sides w.r.t. x, we get $\Rightarrow \frac{du}{dx} = e^{x \sin^2 x} [x(\sin 2x) + \sin^2 x] \quad \text{----- (2)}$ Also , $v = (\sin x)^x$ $\Rightarrow \log v = x \log (\sin x)$ Differentiating both sides w.r.t. x, we get $\frac{1}{v} \frac{dv}{dx} = x \cot x + \log (\sin x)$ $\frac{dv}{dx} = (\sin x)^x [x \cot x + \log(\sin x)] \quad \text{----- (3)}$ Substituting from – (2), – (3) in – (1) we get $\frac{dy}{dx} = e^{x \sin^2 x} [x \sin 2x + \sin^2 x] + (\sin x)^x [x \cot x + \log(\sin x)]$	1 1 $\frac{1}{2}$
31	RHD = $\lim_{h \rightarrow 0} \frac{f(1+h) - f(1)}{h} = \lim_{h \rightarrow 0} \frac{[1+h] - [1]}{h}$ $= \lim_{h \rightarrow 0} \frac{(1-1)}{h} = 0$ LHD = $\lim_{h \rightarrow 0} \frac{f(1-h) - f(1)}{-h} = \lim_{h \rightarrow 0} \frac{[1-h] - [1]}{-h} = \lim_{h \rightarrow 0} \frac{0-1}{-h}$ $= \lim_{h \rightarrow 0} \frac{1}{h} = \infty$ Since, RHD $\neq$ LHD Therefore $f(x)$ is not differentiable at $x = 1$ OR $y = b \tan \theta \Rightarrow \frac{dy}{d\theta} = b \sec^2 \theta \dots (1)$ $x = a \sec \theta \Rightarrow \frac{dx}{d\theta} = a \sec \theta \tan \theta \dots (2)$	1 1 1




	$\frac{dy}{dx} = \frac{\frac{dy}{d\theta}}{\frac{dx}{d\theta}} = \frac{b \sec^2 \theta}{a \sec \theta \tan \theta} = \frac{b}{a} \operatorname{cosec} \theta$ <i>Differentiating both sides w.r.t. x, we get</i> $\begin{aligned} \frac{d^2y}{dx^2} &= \frac{-b}{a} \operatorname{cosec} \theta \cot \theta \times \frac{d\theta}{dx} \\ &= \frac{-b}{a} \operatorname{cosec} \theta \cot \theta \times \frac{1}{a \sec \theta \tan \theta} \quad [\text{using (2)}] \\ &= \frac{-b}{a.a} \cot^3 \theta \end{aligned}$ $\left. \frac{d^2y}{dx^2} \right _{\theta=\frac{\pi}{6}} = \frac{-b}{a} \left[\cot \frac{\pi}{6} \right]^3 = \frac{-b}{a} (\sqrt{3})^3 = -\frac{3\sqrt{3}b}{a.a}$	$1\frac{1}{2}$ $\frac{1}{2}$
32	$f(x) = \tan x - 4x$ $f'(x) = \sec^2 x - 4$ a) For $f(x)$ to be strictly increasing $f'(x) > 0$ $\Rightarrow \sec^2 x - 4 > 0$ $\Rightarrow \sec^2 x > 4$ $\Rightarrow \cos^2 x < \frac{1}{4} \Rightarrow \cos^2 x < \left(\frac{1}{2}\right)^2$ $\Rightarrow -\frac{1}{2} < \cos x < \frac{1}{2} \Rightarrow \frac{\pi}{3} < x < \frac{\pi}{2}$ b) For $f(x)$ to be strictly decreasing $f'(x) < 0$ $\Rightarrow \sec^2 x - 4 < 0$ $\Rightarrow \sec^2 x < 4$ $\Rightarrow \cos^2 x > \frac{1}{4}$ $\Rightarrow \cos^2 x > \left(\frac{1}{2}\right)^2$ $\Rightarrow \cos x > \frac{1}{2} \left[\because x \in \left(0, \frac{\pi}{2}\right) \right]$ $\Rightarrow 0 < x < \frac{\pi}{3}$	$\frac{1}{2}$ $1\frac{1}{2}$ 1



	Required Area = $\frac{4}{3} \int_0^6 \sqrt{6^2 - x^2} dx$  $= \frac{4}{3} \left[\frac{x}{2} \sqrt{6^2 - x^2} + 18 \sin^{-1} \left(\frac{x}{6} \right) \right]_0^6$ $= \frac{4}{3} \left[18 \times \frac{\pi}{2} - 0 \right] = 12\pi \text{ sq units}$	$\frac{1}{2}$ $\frac{1}{2}$ 1 1
35	The given differential equation can be written as $\frac{dy}{dx} = \frac{y + 2x^2}{x} \Rightarrow \frac{dy}{dx} - \frac{1}{x}y = 2x$ Here $P = -\frac{1}{x}$, $Q = 2x$ $IF = e^{\int P dx} = e^{-\int \frac{1}{x} dx} = e^{-\log x} = \frac{1}{x}$ The solutions is : $y \times \frac{1}{x} = \int \left(2x \times \frac{1}{x} \right) dx$ $\Rightarrow \frac{y}{x} = 2x + c$ $\Rightarrow y = 2x^2 + cx$	$\frac{1}{2}$ 1 1 $\frac{1}{2}$
36	$A = 1(-1 - 2) - 2(-2 - 0) = -3 + 4 = 1$ A is nonsingular, therefore A^{-1} exists $Adj A = \begin{bmatrix} -3 & -2 & -4 \\ 2 & 1 & 2 \\ 2 & 1 & 3 \end{bmatrix}$ $\Rightarrow A^{-1} = \frac{1}{ A } (Adj A) = \begin{bmatrix} -3 & -2 & -4 \\ 2 & 1 & 2 \\ 2 & 1 & 3 \end{bmatrix}$	$\frac{1}{2}$ $1\frac{1}{2}$

$a_2 = 5i - 2j$ $b_2 = 3\hat{i} + 2\hat{j} + 6\hat{k}$ $\vec{a}_2 - \vec{a}_1 = 2\hat{i} - 4\hat{j} + 4\hat{k}$ $\vec{b}_1 \times \vec{b}_2 = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 2 & 2 \\ 3 & 2 & 6 \end{vmatrix} = \hat{i}(12 - 4) - \hat{j}(6 - 6) + \hat{k}(2 - 6)$ $\vec{b}_1 \times \vec{b}_2 = 8\hat{i} + 0\hat{j} - 4\hat{k} = 8\hat{i} - 4\hat{k}$ $\therefore (\vec{b}_1 \times \vec{b}_2) \cdot (\vec{a}_2 - \vec{a}_1) = 16 - 16 = 0$ $\therefore$ The lines are intersecting and the shortest distance between the lines is 0. Now for point of intersection $3\hat{i} + 2\hat{j} - 4\hat{k} + \lambda(\hat{i} + 2\hat{j} + 2\hat{k}) = 5\hat{i} - 2\hat{j} + \mu(3\hat{i} + 2\hat{j} + 6\hat{k})$ $\Rightarrow 3 + \lambda = 5 + 3\mu \quad \text{--- -- (1)}$ $2 + 2\lambda = -2 + 2\mu \quad \text{--- -- (2)}$ $-4 + 2\lambda = 6\mu \quad \text{--- -- (3)}$ Solving (1) and (2) we get, $\mu = -2$ and $\lambda = -4$ Substituting in equation of line we get $\vec{r} = 5i - 2j + (-2)(3i + 2j + 6k) = -i - 6j - 12k$ Point of intersection is $(-1, -6, -12)$ OR Let P be the given point and Q be the foot of the perpendicular. Equation of PQ $\frac{x+1}{2} = \frac{y-3}{1} = \frac{z+6}{-2} = \lambda$  Let coordinates of Q be $(2\lambda - 1, \lambda + 3, -2\lambda - 6)$ Since Q lies in the plane $2x + y - 2z + 5 = 0$ $\therefore 2(2\lambda - 1) + (\lambda + 3) - 2(-2\lambda - 6) + 5 = 0$ $\Rightarrow 4\lambda - 2 + \lambda + 3 + 4\lambda + 12 + 5 = 0$	1 1 1 1 1 1 1 1
---	---

	$\Rightarrow 9\lambda + 18 = 0 \qquad \Rightarrow \lambda = -2$$\therefore \text{ coordinates of } Q \text{ are } (-5, \quad 1, -2)$ Length of the perpendicular = $\sqrt{(-5 + 1)^2 + (1 - 3)^2 + (-2 + 6)^2}$ $= 6 \text{ units}$	1 1 1										
38	Max $Z = 3x + y$ Subject to $x + 2y \geq 100$ $2x - y \leq 0$ $2x + y \leq 200$ $x \geq 0, \quad y \geq 0$ ----- ----- ----- (1) (2) (3)  <table><tr><th>Corner Points</th><th>$Z = 3x + y$</th></tr><tr><td>A (0, 50)</td><td>50</td></tr><tr><td>B (0, 200)</td><td>200</td></tr><tr><td>C (50, 100)</td><td>250</td></tr><tr><td>D (20, 40)</td><td>100</td></tr></table> $\text{Max } z = 250 \text{ at } x = 50, \quad y = 100$	Corner Points	$Z = 3x + y$	A (0, 50)	50	B (0, 200)	200	C (50, 100)	250	D (20, 40)	100	3 1 1
Corner Points	$Z = 3x + y$											
A (0, 50)	50											
B (0, 200)	200											
C (50, 100)	250											
D (20, 40)	100											



OR

(i)

Corner points	$Z = 3x - 4y$
O(0,0)	0
A(0,8)	-32
B(4,10)	-28
C(6,8)	-14
D(6,5)	-2
E(4,0)	12

$Max\ Z = 12\ at\ E(4,0)$

$Min\ Z = -32\ at\ A(0,8)$

(ii) Since maximum value of Z occurs at B(4,10) and C(6, 8)

$\therefore 4p + 10q = 6p + 8q$

$\Rightarrow 2q = 2p$

$\Rightarrow p = q$

Number of optimal solution are infinite

$1\frac{1}{2}$

<

