

Class XI Session 2025-26

Subject - Chemistry

Sample Question Paper - 3

Time Allowed: 3 hours

Maximum Marks: 70

General Instructions:

1. There are 33 questions in this question paper with internal choice.
2. SECTION A consists of 16 multiple-choice questions carrying 1 mark each.
3. SECTION B consists of 5 very short answer questions carrying 2 marks each.
4. SECTION C consists of 7 short answer questions carrying 3 marks each.
5. SECTION D consists of 2 case-based questions carrying 4 marks each.
6. SECTION E consists of 3 long answer questions carrying 5 marks each.
7. All questions are compulsory.
8. The use of log tables and calculators is not allowed

Section A

1. Which one of the following is isoelectronic with Ne? [1]
a) N^{3-} b) All of these
c) Al^{3+} d) Mg^{2+}
2. Maximum entropy will be in which of the following? [1]
a) water vapour b) Ice
c) liquid water d) snow
3. The number of moles of solute present in 1 kg of solvent is called: [1]
a) normality b) mole fraction
c) molarity d) molality
4. Which conformation of ethane has the lowest potential energy? [1]
a) Staggered b) Eclipsed
c) All will have equal PE d) Skewed
5. Position isomerism is not shown by _____. [1]
a) alkanes b) alkenes
c) aldehydes d) alcohols
6. The wavelength of a ball of mass 0.1 kg moving with a velocity of 10 ms^{-1} will be: [1]
a) $6.626 \times 10^{-35} \text{ m}$ b) $6.626 \times 10^{-34} \text{ m}$



Reason (R): Addition of HBr on alkene proceed by carbocation intermediate.

- a) Both A and R are true and R is the correct explanation of A. b) Both A and R are true but R is not the correct explanation of A.
c) A is true but R is false. d) A is false but R is true.

15. **Assertion (A):** Atomic mass has no unit but is expressed in amu. [1]

Reason (R): It is the average mass of an atom taking care of the relative abundance of all its possible isotopes.

- a) Both A and R are true and R is the correct explanation of A. b) Both A and R are true but R is not the correct explanation of A.
c) A is true but R is false. d) A is false but R is true.

16. **Assertion (A):** The nuclear isomers are the atoms with the same atomic number and same mass number, but with different radioactive properties. [1]

Reason (R): The nucleus in the excited state will evidently have a different half-life as compared to that in the ground state.

- a) Both A and R are true and R is the correct explanation of A. b) Both A and R are true but R is not the correct explanation of A.
c) A is true but R is false. d) A is false but R is true.

Section B

17. Which element do you think would have been named by: [2]

- i. Lawrence Berkeley Laboratory
ii. Seaborg's group?

18. Oxygen is prepared by the catalytic decomposition of potassium chlorate (KClO_3). Decomposition of potassium chlorate gives potassium chloride (KCl) and oxygen (O_2). If 2.4 moles of oxygen is needed for an experiment, how many grams of potassium chlorate must be decomposed? [2]

19. Write the structure and IUPAC names of different structural isomers of alkenes corresponding to C_5H_{10} . [2]

OR

How will you convert benzene into






- i. p - nitrobromobenzene
ii. m - nitrobromobenzene


20. Lead chloride has a solubility product of 1.7×10^{-5} at 298 K. Calculate its solubility at this temperature. [2]

21. Which of the following will not show deflection from the path on passing through an electric field? [2]
Proton, cathode rays, electron, neutron.

Section C

22. Identify the type of redox reaction taking place in the following. [3]





23. **Answer:** [3]
- (a) Give the mathematical expression of enthalpy. [1]
- (b) Neither q nor W is a state function but $q + W$ is a state function. Explain why? [1]
- (c) The standard heat of formation of $\text{Fe}_2\text{O}_3(\text{s})$ is 824.2 kJ mol^{-1} . Calculate heat change for the reaction. [1]
- $$4\text{Fe}(\text{s}) + 3\text{O}_2(\text{g}) \rightarrow 2\text{Fe}_2\text{O}_3(\text{s})$$

24. The reactant which is entirely consumed in reaction is known as limiting reagent. In the reaction $2\text{A} + 4\text{B} \rightarrow 3\text{C} + 4\text{D}$, when 5 moles of A react with 6 moles of B, then [3]

- i. Which is the limiting reagent?
- ii. Calculate the amount of C formed?

25. The electronic configuration of some elements are given below: [3]

- a. $1s^2, 2s^2, 2p^6, 3s^2$
- b. $1s^2, 2s^2, 2p^6$
- c. $1s^2, 2s^2, 2p^2$
- d. $1s^2, 2s^2, 2p^6, 3s^1$
- e. $1s^2, 2s^2, 2p^5$

Answer the following questions:

- i. Name the elements.
- ii. Which of these have the lowest Ionization enthalpy?
- iii. Which is a halogen?
- iv. Which is an alkali metal?
- v. Which is an inert gas?
26. Dual behaviour of matter proposed by de Broglie led to the discovery of electron microscope often used for the highly magnified images of biological molecules and other types of material. If the velocity of the electron in this microscope is $1.6 \times 10^6 \text{ ms}^{-1}$, Calculate de Broglie wavelength associated with this electron. [3]
27. Describe the change in hybridisation (if any) of the Al atom in the following reaction. [3]
- $$\text{AlCl}_3 + \text{Cl}^- \rightarrow \text{AlCl}_4^-$$
28. 100 mL of a liquid is contained in an insulated container at a pressure of 1 bar. The pressure is steeply increased to 100 bar. The volume of the liquid is decreased by 1 mL at this constant pressure. Find ΔH and ΔU . [3]

Section D

29. **Read the text carefully and answer the questions:** [4]

The existing large number of organic compounds and their ever-increasing numbers has made it necessary to classify them on the basis of their structures. Organic compounds are broadly classified as open-chain compounds which are also called aliphatic compounds. Aliphatic compounds further classified as homocyclic and heterocyclic compounds. Aromatic compounds are special types of compounds. Alicyclic compounds, aromatic compounds may also have heteroatom in the ring. Such compounds are called heterocyclic aromatic compounds. Organic compounds can also be classified on the basis of functional groups, into families or homologous series. The members of a homologous series can be represented by general molecular formula and the successive members differ from each other in a molecular formula by a $-\text{CH}_2$ unit.

- (a) The successive members of a homologous series differ by which mass of amu?



OR

Is tetrahydrofuran is aromatic compounds?

- (b) Does Pyridine, pyrrole, thiophene are all heteroaromatic compounds
- (c) Difference between heterocyclic and homocyclic compound.

30. **Read the text carefully and answer the questions:**

[4]

When anions and cations approach each other, the valence shell of anions are pulled towards the cation nucleus and thus, the shape of the anion is deformed. The phenomenon of deformation of anion by a cation is known as polarization and the ability of the cation to polarize the anion is called as polarizing power of cation. Due to polarization, sharing of electrons occurs between two ions to some extent and the bond shows some covalent character.

The magnitude of polarization depends upon a number of factors.

- (a) Out of AlCl_3 and AlI_3 which halides show maximum polarization?
- (b) Out of AlCl_3 and CaCl_2 which one is more covalent in nature?
- (c) The non-aqueous solvent like ether is added to the mixture of LiCl , NaCl and KCl . Which will be extracted into the ether?

OR

Out of CaF_2 and CaI_2 which one has a minimum melting point?

Section E

31. **Answer:**

[5]


- (a) i. Why is nitric acid added to sodium extract before adding silver nitrate for testing halogens? [2.5]
- ii. How does [2.5]
 - i. an electron withdrawing group (EWG) and
 - ii. an electron donating group (EDG) influence the acid strength of carboxylic end?

OR

- i. Suggest a method to purify [2.5]
 - i. a liquid which decomposes at its boiling point.
 - ii. kerosene oil containing water.
 - iii. camphor containing traces of common salt.
- ii. Differentiate between the principle of estimation of nitrogen in an organic compound [2.5]
 - i. Dumas method
 - ii. Kjeldahl's method.

32. **Attempt any five of the following:**

[5]

- (a) Can a catalyst change the position of equilibrium in a reaction? [1]
- (b) To which category of compounds does cyclohexane belong? [1]
- (c) Write an IUPAC name:  [1]
- (d) What is hydrogenation? [1]
- (e) What are conformations? [1]
- (f) The intermediate carbocation formed in the reactions of HI , HBr and HCl with propene is the same [1]
and the bond energy of HCl , HBr and HI is $430.5 \text{ kJ mol}^{-1}$, $363.7 \text{ kJ mol}^{-1}$ and $296.8 \text{ kJ mol}^{-1}$ respectively. What will be the order of reactivity of these halogen acids?



(g) State Le chatelier's principle.

[1]

33. Calculate the degree of ionization of 0.05 M acetic acid if its pK_a value is 4.74. How is the degree of dissociation affected when its solution also contains
- 0.01 M
 - 0.1 M HCl?

[5]

OR

Write a relation between ΔG and Q and define the meaning of each term and answer the following:

- Why a reaction proceeds forward when $Q < K$ and no net reaction occurs when $Q = K$.
- Explain the effect of an increase in pressure in terms of reaction quotient Q for the reaction:
$$\text{CO(g)} + 3\text{H}_2\text{(g)} \rightleftharpoons \text{CH}_4\text{(g)} + \text{H}_2\text{O(g)}$$



Solution

Section A

1. **(b)** All of these
Explanation:
Isoelectronic species have same number of electrons.
Ne (Z=10) has 10 electrons.
N (Z=7) has 7 electrons and with addition of 3 more electrons it becomes N^{3-} anion which has 10 electrons.
Mg (Z=12) has 12 electron and with removal of 2 electrons it becomes Mg^{2+} cation which has 10 electrons.
Al (Z=13) has 13 electrons and with removal of 3 electrons it becomes Al^{3+} cation which has 10 electrons.
Since all the species have same number of electrons that is 10 , so they are isoelectronic.
2. **(a)** water vapour
Explanation:
water vapour
3. **(d)** molality
Explanation:
Mathematically, molality is expressed as, $\text{Molality}(m) = \frac{\text{No. of moles of solute}}{\text{Mass of solvent in kg}}$
4. **(a)** Staggered
Explanation:
In staggered conformation of ethane, any two H-atoms on adjacent C-atoms are as far apart as possible. As a result, the repulsions between the electron clouds of σ -bonds of two non bonded H-atoms are minimum and hence stability is maximum.
5. **(c)** aldehydes
Explanation:
aldehydes
6. **(b)** $6.626 \times 10^{-34} \text{ m}$
Explanation:
Given : mass = 0.1 kg; velocity = 10 m/s.
We know,
 $\lambda = \frac{h}{mv}$ where h is Planck's constant.
 $\lambda = \frac{6.626 \times 10^{-34}}{10 \times 0.1} = 6.626 \times 10^{-34} \text{ m}$
7. **(d)** $-74.8 \text{ kJ mol}^{-1}$
Explanation:
 $\text{CH}_4 + 2\text{O}_2 \rightarrow \text{CO}_2 + 2\text{H}_2\text{O} \quad \Delta H_1 = -890.3 \text{ kJ/mol} \dots(1)$
 $\text{C(s)} + \text{O}_2 \rightarrow \text{CO}_2 \quad \Delta H_2 = -393.5 \text{ kJ/mol} \dots(2)$
 $\text{H}_2 + 0.5\text{O}_2 \rightarrow \text{H}_2\text{O} \quad \Delta H_3 = -285.8 \text{ kJ/mol} \dots(3)$
 $\text{C(s)} + 2\text{H}_2 \rightarrow \text{CH}_4 \quad \Delta H = \Delta H_2 + 2(\Delta H_3) - \Delta H_1$
 $\Delta H = -393.5 + 2(-285.8) - (-890.3)$
 $= -74.8 \text{ kJ/mol}$



8. **(d) 2**
Explanation:
 Explanation: number of angular node= l
 for 4d orbital, $l=2$
9. **(d) isotopes**
Explanation:
 isotopes
10. **(a) increases with increase in molecular mass.**
Explanation:
 As molecular mass increases, the magnitude of Van der Waals forces of attraction increases and hence boiling point increases accordingly.
11. **(a) a, b**
Explanation:
 a) $K_4[Fe(CN)_6] + H_2SO_4 + H_2O \rightarrow K_2SO_4 + CO + FeSO_4 + (NH_4)_2SO_4$
 b) $CuSO_4 + NH_3 \rightarrow [Cu(NH_3)_4]SO_4$
 Since, oxidation number of each element does not change in these reactions, so these are not redox reactions.
12. **(d) the enthalpy change that accompanies melting of one mole of a solid substance in the standard state.**
Explanation:
 Molar enthalpy of fusion (always increases i.e. $\Delta H = \text{positive}$) is the amount of energy needed to change completely one mole of a substance from the solid phase to the liquid phase in the standard state.
13. **(a) Both A and R are true and R is the correct explanation of A.**
Explanation:
 The liquids are having sufficient difference in their boiling points. Liquids having different boiling points vaporize at different temperatures. The vapours are cooled and the liquids so formed are collected separately.
14. **(b) Both A and R are true but R is not the correct explanation of A.**
Explanation:
 Both A and R are true but R is not the correct explanation of A.
15. **(a) Both A and R are true and R is the correct explanation of A.**
Explanation:

$$\text{Atomic mass} = \frac{\text{Average mass of an atom}}{1/12 \times \text{Mass of an atom of } ^{12}\text{C}}$$

$$\text{Average mass of an atom} = \frac{R \cdot A(1) \times M.No + R \cdot A(2) \times M.No}{R \cdot A(1) + R \cdot A(2)}$$
 Here R.A = Relative abundance, M.No = Mass number and 1 and 2 refers the two possible isotopes. As atomic mass is a ratio so its has no unit.
16. **(a) Both A and R are true and R is the correct explanation of A.**
Explanation:
 The nuclear isomerism in the nuclei of the same mass number and same atomic number arises due to different radioactive properties. The reason for nuclear isomerism is the different energy states of two isomeric nuclei. One may be in the ground state and the other in an excited state. The nucleus in the excited state will have a different half-life.


Section B

17. i. Lawrencium (Lr) with atomic number (Z) 103.



ii. Seaborgium (Sg) with atomic number (Z) 106.

18. The balanced equation is




2 mol of KClO₃ produced = 3 moles of O₂

3 moles of O₂ is produced by decomposition of 245g of KClO₃

2.4 moles of O₂ will be produced by the decomposition of KClO₃ = $\frac{245 \times 2.4}{3} = 196.0 \text{ g}$.

19.


- Pent-1-ene



- Pent-2-ene




- 2-Methylbut-2-ene




- 3-Methylbut-2-ene



- 2-Methylbut-ene




OR



20. According to the question, the solubility product of lead chloride at 298 K is 1.7×10^{-5} .

Reaction:



Let the solubility of PbCl₂ be S mol/L.

Then the solution will contain S moles of Pb²⁺ ions and 2S moles of Cl⁻ ions respectively per litre.

$$\therefore K_{sp} = [\text{Pb}^{2+}] [\text{Cl}^{-}]^2$$

$$= S \times (2S)^2$$

$$= 4S^3$$

$$\Rightarrow 4S^3 = 1.7 \times 10^{-5}$$

$$\Rightarrow S^3 = \frac{1.7 \times 10^{-5}}{4} = 0.425 \times 10^{-5}$$

$$\text{Therefore, } S = 1.620 \times 10^{-2} \text{ mol L}^{-1}.$$

21. Out of proton, cathode rays, electron, and neutron. The neutral particles are only neutron. So, it will not show deflection from the path on passing through an electric field.

Section C

22. i. Combination reaction

ii. Displacement reaction

iii. Decomposition reaction

iv. Metal displacement reaction

v. Non-metal displacement reaction

vi. Disproportionation reaction

23. Answer:


(i) Mathematical expression of enthalpy is $H = U + pV$,
where U is internal energy.

(ii) q and W are not state functions. But as we know that,
 $q + W = \Delta U$, which is a state function.
Hence, $q + W$ is a state function.

(iii) According to the question, the standard heat of formation of $\text{Fe}_2\text{O}_3(\text{s})$ is $824.2 \text{ kJ mol}^{-1}$.

The standard heat of formation of Fe and O_2 is zero because they are in their basic standard states.

Reaction:



$$\begin{aligned}\text{We know that, } \Delta H^\circ &= \sum \Delta H_f^\circ(\text{products}) - \sum \Delta H_f^\circ(\text{reactants}) \\ &= [2 \times \Delta H_f^\circ \text{Fe}_2\text{O}_3(\text{s})] - [4\Delta H_f^\circ \text{Fe}(\text{s}) + 3\Delta H_f^\circ \text{O}_2(\text{g})] \\ &= 2(-824.2) - [4 \times 0 + 3 \times 0] \\ &= -1648.4 \text{ kJ}.\end{aligned}$$

24. The given equation is : $2\text{A} + 4\text{B} \rightarrow 3\text{C} + 4\text{D}$

i. It is clear from the above equation that: 2 moles of 'A' requires 4 moles of 'B' for the reaction i.e. ratio of moles of A to B is 2: 4 or 1:2. Hence, for 5 moles of 'A', the moles of 'B' required = 5 mole of A $\times \frac{4 \text{ mol of B}}{2 \text{ mol of A}} = 10 \text{ mol of B}$. But we have only 6 moles of 'B', hence, 'B' is the limiting reagent.

ii. Since 4 moles of 'B' gives 3 moles of 'C'. Hence, 6 moles of 'B' will produce $\frac{3}{4} \times 6 = 4.5$ mole of C.

25. i. (a) Magnesium, (b) Neon (c) Carbon (d) Sodium (e) Flourine

ii. $1s^2, 2s^2, 2p^6, 3s^1$ (Sodium)


iii. $1s^2, 2s^2, 2p^5$ (Flourine)

iv. $1s^2, 2s^2, 2p^6, 3s^1$ (Sodium)


v. $1s^2, 2s^2, 2p^6$ (Neon)

$$\begin{aligned}26. \lambda &= \frac{h}{mv} = \frac{(6.626 \times 10^{-34} \text{ Js})}{(9.1 \times 10^{-31} \text{ kg}) \times (1.6 \times 10^6 \text{ ms}^{-1})} \\ &= 0.455 \times 10^{-34+25} \text{ m} \\ &= 0.455 \text{ nm} \\ &= 455 \text{ pm}\end{aligned}$$

27. The valence orbital picture of aluminum in the ground state can be shown as:



The orbital picture of aluminum in the excited state can be shown as:



Hence, it undergoes sp^2 hybridization to give a trigonal planar arrangement (in AlCl_3). To form AlCl_4^- , the empty $3p_z$ orbital also gets involved and the hybridization changes from sp^2 to sp^3 . As a result, the shape becomes tetrahedral.

28. According to the question, $p_1 = 1 \text{ bar}$, $p_2 = 100 \text{ bar}$, $V_1 = 100 \text{ mL}$, $V_2 = 99 \text{ mL}$.

We know that, $\Delta U = q + W$

For the adiabatic process, $q = 0$

So, $\Delta U = W$

Now, $W = -p\Delta V = -100(99 - 100) = 100 \text{ bar ml}$

We know that, $\Delta H = \Delta U + \Delta pV$

$$= 100 + p_2 V_2 - p_1 V_1$$

$$= 100 + (100 \times 99) - (1 \times 100)$$

$$= 100 + 9900 - 100$$

$$= 9900 \text{ bar mL}$$



Section D

29. Read the text carefully and answer the questions:

The existing large number of organic compounds and their ever-increasing numbers has made it necessary to classify them on the basis of their structures. Organic compounds are broadly classified as open-chain compounds which are also called aliphatic compounds. Aliphatic compounds further classified as homocyclic and heterocyclic compounds. Aromatic compounds are special types of compounds. Alicyclic compounds, aromatic compounds may also have heteroatom in the ring. Such compounds are called heterocyclic aromatic compounds. Organic compounds can also be classified on the basis of functional groups, into families or homologous series. The members of a homologous series can be represented by general molecular formula and the successive members differ from each other in a molecular formula by a $-\text{CH}_2$ unit.

- (i) The successive members of a homologous series are differ by a $-\text{CH}_2$ group. The molecular mass of a $-\text{CH}_2$ group is 14 amu. Hence, each successive homologue of a homologous series differ by a mass of 14 amu.

OR

Tetrahydrofuran is non-aromatic, due to absence of conjugation in π electrons, and it does not follow Huckel's rule.

- (ii) Heterocyclic compounds are a major class of organic compounds characterized by the fact that some or all of the atoms in their molecules are joined in rings containing at least one atom of an element other than carbon and follow Huckels rule, the most common heterocycles are those having five or six-membered rings and containing hetero members of Nitrogen, oxygen, sulphur. Pyridine, pyrrole, thiophene are all heteroaromatic compounds
- (iii) A cyclic compound in which the ring includes at least one atom of an element different from the rest is called heterocyclic compound. A homocyclic compound is a cyclic compound in which all the ring atoms are the same.

30. Read the text carefully and answer the questions:

When anions and cations approach each other, the valence shell of anions are pulled towards the cation nucleus and thus, the shape of the anion is deformed. The phenomenon of deformation of anion by a cation is known as polarization and the ability of the cation to polarize the anion is called as polarizing power of cation. Due to polarization, sharing of electrons occurs between two ions to some extent and the bond shows some covalent character.

The magnitude of polarization depends upon a number of factors.

- (i) AlI_3 halides show maximum polarization. The most covalent halide is AlI_3 .

Since lesser, the electronegativity difference, the more covalent is the aluminum halide.

- (ii) AlCl_3 is more covalent in nature.

- (iii) LiCl will be extracted into the ether.

OR

CaI_2 has a minimum melting point.


Section E

31. Answer:


- (i) i. Nitric acid is added to sodium extract so as to decompose



- ii. The influence of the inductive effect on acidity is best understood in terms of the conjugate base, RCOO^- and can be summarised as follows



Electron withdrawing group destabilises RCOO^- because there exists a repulsion between electrons from EDG and negative charge of O. Hence, EDG weakens the acid.



Electron withdrawing group stabilises RCOO^- by taking negative charge from O. Hence, EWG strengthens the acid.

OR

- i. i. Distillation under reduced pressure.

- ii. Since the two liquids are immiscible, the technique of solvent extraction with a separating funnel is used.

Kerosene being lighter than water forms the upper layer while water forms the lower layer.



The lower water layer is run off when kerosene oil is obtained. It is dried over anhydrous CaCl_2 or MgSO_4 and then distilled to give pure kerosene oil.

iii. Sublimation Camphor sublimes while common salt remains as residue in the China dish.


ii. (i) **Dumas method:** The organic compound is heated strongly with excess of CuO (Cupric oxide) in an atmosphere of CO_2 when free nitrogen, CO_2 and H_2O are obtained.

(ii) **Kjeldahl's method:** A known mass of the organic compound is heated strongly with conc. H_2SO_4 and a little potassium sulphate and a little mercury (a catalyst). As a result of reaction the nitrogen present in the organic compound is converted to ammonium sulphate.

32. Attempt any five of the following:


(i) A catalyst speeds up the forward and back reaction to the same extent. Because adding a catalyst doesn't affect the relative rates of the two reactions, a catalyst cannot change the position of equilibrium in a chemical reaction

(ii) Cyclohexane belongs to saturated alicyclic hydrocarbons.



(iv) **Hydrogenation:** Addition of hydrogen to alkenes and alkynes in the presence of finely divided catalysts like Pt, Pd or Ni to form alkanes is known as hydrogenation.

Example:



(v) Conformations are spatial arrangements which are obtained by rotation around sigma bonds.

Explanation : In chemistry, **conformational isomerism** is a form of stereoisomers in which the isomers can be interconverted just by rotations about formally single bonds (refer to figure on single bond rotation). Such isomers are generally referred to as **conformational isomers** or **conformers**.

(vi) The bond dissociation enthalpy decreases in the order $\text{HCl} > \text{HBr} > \text{HI}$, the order of reactivity of these halogen acids is in the reverse order i.e., $\text{HI} > \text{HBr} > \text{HCl}$.

(vii) **Le chatelier's principle:** If a system at equilibrium is subjected to change in the temperature, pressure or concentration of the reactants or the products that govern the equilibrium, then the equilibrium shifts in that direction in which this change is reduced or nullified.


33. $\text{pK}_a = -\log K_a = 4.74$

$$\text{or } \log K_a = -4.74 = \bar{5}.26 \therefore K_a = 1.82 \times 10^{-5}$$

$$\alpha = \sqrt{K_a/C} = \sqrt{(1.82 \times 10^{-5})/(5 \times 10^{-2})} = 1.908 \times 10^{-2}$$

In presence of HCl , due to high concentration of H^+ ion, dissociation equilibrium will shift backward, i.e. dissociation of acetic acid will decrease.

a. In presence of 0.01 M HCl , if x is the amount dissociated, then



(0.01 M H^+ ions are obtained from 0.01 M HCl)

$$\therefore K_a = \frac{x(0.01)}{0.05} \text{ or } \frac{x}{0.05} = \frac{K_a}{0.01} = \frac{1.82 \times 10^{-5}}{10^{-2}} = 1.82 \times 10^{-3}$$

$$\text{or } \alpha = 1.82 \times 10^{-3} \left(\because \alpha = \frac{\text{Amount dissociated}}{\text{Amount taken}} \right)$$

The degree of ionization is $\alpha = 1.82 \times 10^{-3}$

b. In the presence of 0.1 M HCl , if y is the amount of acetic acid dissociated, then at equilibrium

$$[\text{CH}_3\text{COOH}] = 0.05 - y \approx 0.05 \text{ M}$$

$$[\text{CH}_3\text{COO}^-] = y, [\text{H}^+] = 0.1 \text{ M} + y \approx 0.1 \text{ M}$$

$$K_a = \frac{y(0.1)}{0.05} \text{ or } \frac{y}{0.05} = \frac{K_2}{0.1} = \frac{1.82 \times 10^{-5}}{10^{-1}} = 1.82 \times 10^{-4} \text{ i.e.}$$

$$\alpha = 1.82 \times 10^{-4}$$

The degree of ionization is $\alpha = 1.82 \times 10^{-4}$

OR

a. We know that, $\Delta G = \Delta G^\circ + RT \ln Q$

Where,

ΔG° = Change in free energy as the reaction proceeds

ΔG = Standard free energy change

Q = Reaction quotient

T = Absolute temperature

Also, $\Delta G^\circ = -RT \ln K$


$$\Rightarrow \Delta G = -RT \ln K + RT \ln Q$$

$$\therefore \Delta G = RT \ln \frac{Q}{K}$$

If $Q < K$, ΔG will be negative. So, the reaction proceeds in the forward direction.

If $Q = K$, $\Delta G = 0$, reaction will be at equilibrium.

b. Reaction:



On increasing the pressure equilibrium will shift in forward direction, it means $Q < K$.

