

Chapter 8. Differential Equation

Formation of Differential Equations

1 Mark Questions

1. Write the degree of the differential equation

$$\left(\frac{dy}{dx}\right)^4 + 3y \frac{d^2y}{dx^2} = 0.$$

Delhi 2013C



The degree of the differential equation is the degree of the highest order derivative, when differential coefficients are made free from radicals and fractions sign.

Given differential equation is

$$\left(\frac{dy}{dx}\right)^4 + 3y \frac{d^2y}{dx^2} = 0$$

Here, highest order derivative is d^2y/dx^2 , whose degree is one. So, degree of differential equation is 1. (1)

2. Write the degree of the differential equation

$$x^3 \left(\frac{d^2y}{dx^2}\right)^2 + x \left(\frac{dy}{dx}\right)^4 = 0.$$

Delhi 2013

Given differential equation is

$$x^3 \left(\frac{d^2y}{dx^2}\right)^2 + x \left(\frac{dy}{dx}\right)^4 = 0$$

Here, all differential coefficients are free from radical sign.

\therefore Degree = 2 (1)

3. Write the degree of the differential equation

$$\left(\frac{dy}{dx}\right)^4 + 3x \frac{d^2y}{dx^2} = 0.$$

Delhi 2013

Given differential equation is

$$\left(\frac{dy}{dx}\right)^4 + 3x\left(\frac{d^2y}{dx^2}\right) = 0$$

Here, all differential coefficients are free from radical sign.

\therefore Degree = 1 (1)

4. Write the differential equation representing the family of curves $y = mx$, where m is an arbitrary constant. All India 2013

Given, family of curves is $y = mx$... (i)

where, m is an arbitrary constant.

Now, differentiating Eq. (i) w.r.t. x , we get

$$\frac{dy}{dx} = m$$

On putting $m = \frac{dy}{dx}$ in Eq. (i), we get

$$y = x \frac{dy}{dx}$$

which is the required differential equation. (1)


4 Marks Questions

5. Find the differential equation of the family of circles in the first quadrant which touch the coordinate axes. All India 2010C



The equation of family of circles in first quadrant, which touch the coordinate axes, is $(x - a)^2 + (y - a)^2 = a^2$, where a is radius of circle. Differentiate it one time and eliminate the arbitrary constant a .

Let a be the radius of family of circles in the first quadrant, which touch the coordinate axes.



Then, coordinates of centre of circle = (a, a) .

(1)

We know that, equation of circle which has centre (h, k) and radius r is given by

$$(x - h)^2 + (y - k)^2 = r^2$$

Here, $(h, k) = (a, a)$ and $r = a$

\therefore Equation of family of such circles is

$$(x - a)^2 + (y - a)^2 = a^2 \quad \dots(i) \quad (1)$$

On differentiating both sides w.r.t. x , we get

$$2(x - a) + 2(y - a) \frac{dy}{dx} = 0$$

$$\Rightarrow x - a + (y - a) \cdot y' = 0 \quad \left[\because \frac{dy}{dx} = y' \right]$$

$$\Rightarrow x + yy' = a + ay'$$

$$\Rightarrow a = \frac{x + yy'}{1 + y'} \quad (1)$$

On putting above value of a in Eq. (i), we get

$$\begin{aligned} \left[x - \frac{x + yy'}{y' + 1} \right]^2 + \left[y - \frac{x + yy'}{y' + 1} \right]^2 &= \left(\frac{x + yy'}{y' + 1} \right)^2 \\ \Rightarrow \left[\frac{xy' + x - x - yy'}{y' + 1} \right]^2 + \left[\frac{yy' + y - x - yy'}{y' + 1} \right]^2 &= \left(\frac{x + yy'}{y' + 1} \right)^2 \end{aligned}$$

On multiplying both sides by $(y' + 1)^2$, we get

$$\begin{aligned} (xy' - yy')^2 + (y - x)^2 &= (x + yy')^2 \\ \Rightarrow (x - y)^2 (y')^2 + (x - y)^2 &= (x + yy')^2 \\ &[\because (x - y)^2 = (y - x)^2] \\ \Rightarrow (x - y)^2 [(y')^2 + 1] &= (x + yy')^2 \end{aligned}$$

which is the required differential equation. **(1)**

- 6.** Find the differential equation of family of circles touching Y-axis at the origin.

HOTS; Delhi 2010; All India 2009, 2008C



The equation of family of circles touching Y-axis at origin is given by $(x - a)^2 + y^2 = a^2$, where a is radius of circle.

Differentiate this equation once because one arbitrary constant is present in the equation and eliminate a .


Given, family of circles touch Y-axis at the origin.

Let radius of family of circles be a .

\therefore Centre of circle = $(a, 0)$ (1)

Now, equation of family of circles with centre $(a, 0)$ and radius a is

$$(x - a)^2 + y^2 = a^2 \quad (1)$$



[putting $(h, k) = (a, 0)$ and $r = a$
in $(x - h)^2 + (y - k)^2 = r^2$]

$$\Rightarrow x^2 + a^2 - 2ax + y^2 = a^2$$

$$\Rightarrow x^2 - 2ax + y^2 = 0 \quad \dots(i)$$

On differentiating both sides w.r.t. x , we get

$$2x - 2a + 2y \frac{dy}{dx} = 0$$

$$\Rightarrow x^2 - 2ax + y^2 = 0 \quad \dots(i)$$

On differentiating both sides w.r.t. x , we get

$$2x - 2a + 2y \frac{dy}{dx} = 0$$

$$\Rightarrow a = x + y \frac{dy}{dx} \quad (1)$$

On putting above value of a in Eq. (i), we get

$$x^2 + y^2 - 2 \left(x + y \frac{dy}{dx} \right) x = 0$$

$$\Rightarrow 2xy \frac{dy}{dx} + x^2 - y^2 = 0$$

$$\Rightarrow 2xyy' + x^2 - y^2 = 0$$


$$\text{or } 2xy \frac{dy}{dx} + x^2 - y^2 = 0$$

which is the required differential equation. (1)

- 7.** Find the differential equation of family of circles touching X-axis at the origin.

HOTS; Delhi 2010C; All India 2009C

Let a be the radius of family of circles which touch X -axis at origin.



\therefore Centre of circle = $(0, a)$

Now, equation of family of such circles is

$$x^2 + (y - a)^2 = a^2 \quad (1)$$

$$\begin{aligned} &[\text{putting } (h, k) = (0, a) \text{ and } r = a \\ &\text{in } (x - h)^2 + (y - k)^2 = r^2] \end{aligned}$$

$$\Rightarrow x^2 + y^2 - 2ay = 0 \quad \dots(i)$$

On differentiating both sides w.r.t. x , we get

$$2x + 2y \frac{dy}{dx} - 2a \frac{dy}{dx} = 0$$

$$\Rightarrow x + (y - a) \frac{dy}{dx} = 0 \quad (1)$$

$$\Rightarrow x + yy' - ay' = 0, \left[\text{where, } y' = \frac{dy}{dx} \right]$$

$$\Rightarrow x + yy' = ay'$$

$$\Rightarrow a = \frac{x + yy'}{y'} \quad (1)$$

On putting above value of a in Eq. (i), we get

$$x^2 + y^2 = 2y \left(\frac{x + yy'}{y'} \right)$$

$$\Rightarrow (x^2 + y^2) \cdot y' = 2xy + 2y^2 \cdot y'$$

$$\Rightarrow x^2 y' + y^2 y' - 2xy - 2y^2 y' = 0$$

$$\Rightarrow x^2 y' - 2xy - y^2 y' = 0$$

$$\Rightarrow y'(x^2 - y^2) = 2xy$$

$$\Rightarrow y' = \frac{2xy}{x^2 - y^2} \quad \text{or} \quad \frac{dy}{dx} = \frac{2xy}{x^2 - y^2}$$

which is the required differential equation. (1)


- 8.** Form the differential equation representing family of ellipses having foci on X-axis and centre at the origin. HOTS; Delhi 2009C



The equation of family of ellipses having foci on X-axis and centre at origin is $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1, a > b$.

Differentiate this equation two times and eliminate two arbitrary constants a and b to get the required result.

We know that, the equation of family of ellipse having foci on X-axis and centre at origin is given by



$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 \quad \dots(i)$$

where, $a > b$ (1)

On differentiating both sides of Eq. (i) w.r.t. x , we get

$$\begin{aligned} \frac{2x}{a^2} + \frac{2yy'}{b^2} &= 0 \quad \left[\text{put } \frac{dy}{dx} = y' \right] \\ \Rightarrow \frac{x}{a^2} &= \frac{-yy'}{b^2} \\ \Rightarrow \frac{yy'}{x} &= \frac{-b^2}{a^2} \quad \dots(ii) \quad (1) \end{aligned}$$

Again, differentiating both sides of Eq. (ii) w.r.t. x , we get

$$\begin{aligned} \frac{\left[x \cdot \frac{d}{dx}(yy') - yy' \cdot \frac{d}{dx}(x) \right]}{x^2} &= 0 \\ \left[\text{using quotient rule of differentiation} \right. \\ \left. \text{in LHS and } \frac{d}{dx} \left(\frac{-b^2}{a^2} \right) = 0 \right] \\ \Rightarrow x \left[y \cdot \frac{d}{dx}(y') + y' \cdot \frac{d}{dx}(y) \right] - yy' \cdot 1 &= 0 \quad (1) \end{aligned}$$

$$\Rightarrow xyy'' + x(y')^2 - yy' = 0$$

$$\text{or } xy \frac{d^2y}{dx^2} + x \left(\frac{dy}{dx} \right)^2 - y \frac{dy}{dx} = 0$$

which is the required differential equation. (1)

- 9.** Form the differential equation representing family of curves given by $(x - a)^2 + 2y^2 = a^2$ where, a is an arbitrary constant. All India 2009

Given equation of family of curves is

$$(x - a)^2 + 2y^2 = a^2 \quad \dots(i)$$

On differentiating both sides w.r.t. x in Eq. (i), we get

$$2(x - a) + 4yy' = 0 \quad \left[\because \frac{d}{dx}(y^2) = 2yy' \right] \quad (1)$$

$$\Rightarrow x - a + 2yy' = 0$$

$$\Rightarrow a = x + 2yy' \quad (1)$$

On putting above value of a in Eq. (i), we get

$$(x - x - 2yy')^2 + 2y^2 = (x + 2yy')^2$$

$$\Rightarrow 4y^2(y')^2 + 2y^2 = x^2 + 4y^2(y')^2 + 4xyy'$$

$$\Rightarrow 2y^2 = x^2 + 4xyy' \quad (1)$$

Hence, the required differential equation is

$$x^2 + 4xyy' = 2y^2$$

$$\text{or } x^2 + 4xy \frac{dy}{dx} = 2y^2 \quad (1)$$

Solution of Different Types of Differential Equations

4 Marks Questions

1. Find the particular solution of the differential equation $\frac{dy}{dx} = 1 + x + y + xy$, given that $y = 0$ when $x = 1$. All India 2014

Given differential equation is

$$\begin{aligned}\frac{dy}{dx} &= 1 + x + y + xy \\ \Rightarrow \frac{dy}{dx} &= 1(1 + x) + y(1 + x) \\ \Rightarrow \frac{dy}{dx} &= (1 + x)(1 + y) \quad \dots(i) \quad (1)\end{aligned}$$

On separating variables, we get

$$\frac{1}{(1 + y)} dy = (1 + x) dx \quad \dots(ii)$$

On integrating both sides of Eq. (ii), we get

$$\int \frac{1}{1 + y} dy = \int (1 + x) dx$$

$$\Rightarrow \log |1+y| = x + \frac{x^2}{2} + C \quad \dots(\text{iii}) \quad (1)$$

Also, given that $y = 0$, when $x = 1$.

On substituting $x = 1, y = 0$ in Eq. (iii), we get

$$\log |1+0| = 1 + \frac{1}{2} + C \Rightarrow C = -\frac{3}{2} \quad [\because \log 1 = 0] \quad (1)$$

Now, on substituting the value of C in Eq. (iii), we get

$$\log |1+y| = x + \frac{x^2}{2} - \frac{3}{2}$$

which is the required particular solution of given differential equation. (1)

2. Find the particular solution of the differential

$$\text{equation } x \frac{dy}{dx} - y + x \operatorname{cosec} \left(\frac{y}{x} \right) = 0 \text{ or}$$

$$\frac{dy}{dx} - \frac{y}{x} + \operatorname{cosec} \left(\frac{y}{x} \right) = 0, \text{ given that } y = 0, \text{ when}$$

$$x = 1.$$

All India 2014C, 2011; Delhi 2009

Given differential equation is

$$x \frac{dy}{dx} - y + x \operatorname{cosec} \left(\frac{y}{x} \right) = 0$$

$$\Rightarrow \frac{dy}{dx} - \frac{y}{x} + \operatorname{cosec} \left(\frac{y}{x} \right) = 0$$

Above equation can be written as

$$\frac{dy}{dx} = \frac{y}{x} - \operatorname{cosec} \left(\frac{y}{x} \right) \quad \dots(i)$$

which is a homogeneous differential equation.

On putting $y = vx$,

$$\Rightarrow \frac{dy}{dx} = v + x \frac{dv}{dx} \text{ in Eq. (i), we get}$$

$$v + x \frac{dv}{dx} = \frac{vx}{x} - \operatorname{cosec} \left(\frac{vx}{x} \right)$$

$$\Rightarrow v + x \frac{dv}{dx} = v - \operatorname{cosec} v$$

$$\Rightarrow x \frac{dv}{dx} = -\operatorname{cosec} v \Rightarrow \frac{dv}{\operatorname{cosec} v} = \frac{-dx}{x} \quad (1)$$

On integrating both sides, we get

$$\int \frac{dv}{\operatorname{cosec} v} = \int -\frac{dx}{x}$$

$$\Rightarrow \int \sin v \, dv = \int -\frac{dx}{x} \quad \left[\because \frac{1}{\operatorname{cosec} v} = \sin v \right]$$

$$\Rightarrow -\cos v = -\log |x| + C$$

$$\left[\begin{array}{l} \because \int \sin x \, dx = -\cos x + C \\ \text{and } \int \frac{1}{x} \, dx = \log |x| + C \end{array} \right]$$

On putting $v = \frac{y}{x}$, we get

$$-\cos \frac{y}{x} = -\log |x| + C$$

$$\Rightarrow \cos \frac{y}{x} = +(\log |x| - C)$$

$$\Rightarrow \frac{y}{x} = \cos^{-1}(\log |x| - C)$$

$$\Rightarrow y = x \cos^{-1}(\log |x| - C) \quad \dots(ii) \quad (1\frac{1}{2})$$

Also, given that $x = 1$ and $y = 0$.

On putting above values in Eq. (ii), we get

$$0 = 1 \cos^{-1}(\log |1| - C)$$

$$\Rightarrow \cos 0^\circ = 0 - C$$

$$\Rightarrow 1 = 0 - C$$

$$\Rightarrow C = -1$$

$$\therefore y = x \cos^{-1}(\log |x| + 1) \quad (1\frac{1}{2})$$

which is required solution.

3. Solve the differential equation

$$x \log x \frac{dy}{dx} + y = \frac{2}{x} \log x. \text{ Foreign 2014; Delhi 2009}$$

Given differential equation is

$$(x \log x) \cdot \frac{dy}{dx} + y = \frac{2}{x} \log x$$

On dividing both sides by $x \log x$, we get

$$\frac{dy}{dx} + \frac{y}{x \log x} = \frac{2 \log x}{x^2 \log x} = \frac{2}{x^2} \quad \dots(i)$$

which is a linear differential equation of first order and is of the form

$$\frac{dy}{dx} + Py = Q \quad \dots(ii)$$

On comparing Eqs. (i) and (ii), we get

$$P = \frac{1}{x \log x} \text{ and } Q = \frac{2}{x^2} \quad (1)$$

$$\therefore \text{IF} = e^{\int \frac{1}{x \log x} dx} = e^{\log \log x}$$

$$\left[\begin{array}{l} \text{for } \int \frac{1}{x \log x} dx \Rightarrow \text{put } \log x = t \Rightarrow \frac{1}{x} dx = dt \\ \therefore \int \frac{1}{t} dt = \log |t| = \log |\log x| \end{array} \right]$$
$$\Rightarrow \text{IF} = \log x \quad [\because e^{\log x} = x] \quad (1)$$

Now, solution of above equation is given by

$$y \times \text{IF} = \int (Q \times \text{IF}) dx + C \quad \dots(iii)$$

On putting IF = log x and $Q = \frac{2}{x^2}$ in Eq. (iii),

we get

$$y \log x = \int \frac{2}{x^2} \log x \, dx$$

$$\Rightarrow y \log x = \log x \int \frac{2}{x^2} \, dx - \int \left(\frac{d}{dx} (\log x) \cdot \int \frac{2}{x^2} \, dx \right) dx$$

[using integration by parts]

$$\Rightarrow y \log x = \log x \cdot 2 \left(-\frac{1}{x} \right) - \int \frac{1}{x} \cdot 2 \left(-\frac{1}{x} \right) dx \quad (1)$$

$$\Rightarrow y \log x = -\frac{2}{x} \log x - \int \frac{2}{x} \left(-\frac{1}{x} \right) dx$$

$$\Rightarrow y \log x = -\frac{2}{x} \log x + \int \frac{2}{x^2} \, dx$$

$$\therefore y \log x = -\frac{2}{x} \log x - \frac{2}{x} + C \quad (1)$$

which is the required solution.

4. Find the general solution of the differential equation $(x - y) \frac{dy}{dx} = x + 2y$.

Delhi 2014C; All India 2010

Given differential equation is

$$(x - y) \frac{dy}{dx} = x + 2y$$

$$\Rightarrow \frac{dy}{dx} = \frac{x + 2y}{x - y} \quad \dots(i) \quad (1)$$

which is a homogeneous equation.

On putting $y = vx$

$$\Rightarrow \frac{dy}{dx} = v + x \frac{dv}{dx} \quad \dots(ii)$$

in Eq. (i), we get

$$v + x \frac{dv}{dx} = \frac{x + 2vx}{x - vx} = \frac{1 + 2v}{1 - v}$$

$$\Rightarrow x \frac{dv}{dx} = \frac{1 + 2v}{1 - v} - v = \frac{1 + 2v - v + v^2}{1 - v}$$

$$\Rightarrow x \frac{dv}{dx} = \frac{1 + v + v^2}{1 - v}$$

$$\Rightarrow \frac{1 - v}{v^2 + v + 1} dv = \frac{dx}{x}$$

On integrating both sides, we get

$$\int \frac{1 - v}{v^2 + v + 1} dv = \int \frac{dx}{x} \quad (1)$$

$$\Rightarrow I = \log |x| + C \quad \dots(iii)$$

where, $I = \int \frac{1 - v}{v^2 + v + 1} dv$

Let $1 - v = A \cdot \frac{d}{dv}(v^2 + v + 1) + B$

$$\Rightarrow 1 - v = A(2v + 1) + B$$

On comparing coefficients of v and constant term from both sides, we get

$$2A = -1 \Rightarrow A = -\frac{1}{2} \quad \text{and} \quad A + B = 1$$

$$\Rightarrow -\frac{1}{2} + B = 1 \Rightarrow B = 1 + \frac{1}{2} \Rightarrow B = \frac{3}{2}$$

$$\text{So, we write } 1 - v = -\frac{1}{2}(2v + 1) + \frac{3}{2}$$

1

3

$$\text{Then, } I = \int \frac{-\frac{1}{2}(2v+1) + \frac{\sqrt{3}}{2}}{v^2 + v + 1} dv$$

$$\Rightarrow I = -\frac{1}{2} \int \frac{2v+1}{v^2 + v + 1} dv + \frac{3}{2} \int \frac{dv}{v^2 + v + 1}$$

$$\Rightarrow I = -\frac{1}{2} \log|v^2 + v + 1| + \frac{3}{2} \int \frac{dv}{v^2 + v + 1 + \frac{1}{4} - \frac{1}{4}}$$

$$\left[\begin{array}{l} \because \int \frac{2v+1}{v^2 + v + 1} dv \Rightarrow \text{put } v^2 + v + 1 = t \\ (2v+1) dv = dt \\ \therefore \int \frac{dt}{t} = \log|t| + c = \log|v^2 + v + 1| + c \end{array} \right]$$

$$\Rightarrow I = -\frac{1}{2} \log|v^2 + v + 1| + \frac{3}{2} \int \frac{dv}{\left(v + \frac{1}{2}\right)^2 + \frac{3}{4}} \quad (1)$$

$$\Rightarrow I = -\frac{1}{2} \log|v^2 + v + 1| + \frac{3}{2} \int \frac{dv}{\left(v + \frac{1}{2}\right)^2 + \left(\frac{\sqrt{3}}{2}\right)^2}$$

$$\Rightarrow I = -\frac{1}{2} \log|v^2 + v + 1| + \frac{3}{2} \times \frac{2}{\sqrt{3}} \tan^{-1} \left(\frac{v + \frac{1}{2}}{\frac{\sqrt{3}}{2}} \right) + C$$

$$\left[\because \int \frac{dx}{x^2 + a^2} = \frac{1}{a} \tan^{-1} \frac{x}{a} + C \right]$$

$$\Rightarrow I = -\frac{1}{2} \log |v^2 + v + 1| + \frac{3}{\sqrt{3}} \tan^{-1} \left(\frac{2v+1}{\sqrt{3}} \right) + C$$

On putting $v = \frac{y}{x}$, we get

$$I = -\frac{1}{2} \log \left| \frac{y^2}{x^2} + \frac{y}{x} + 1 \right| + \sqrt{3} \tan^{-1} \left(\frac{\frac{2y}{x} + 1}{\sqrt{3}} \right) + C$$

$$\left[\because y = vx \therefore v = \frac{y}{x} \right]$$

$$\Rightarrow I = -\frac{1}{2} \log \left| \frac{y^2 + xy + x^2}{x^2} \right| + \sqrt{3} \tan^{-1} \left(\frac{2y + x}{\sqrt{3}x} \right) + C$$

On putting the value of I in Eq. (iii), we get

$$-\frac{1}{2} \log \left| \frac{y^2 + xy + x^2}{x^2} \right| + \sqrt{3} \tan^{-1} \left(\frac{2y + x}{\sqrt{3}x} \right) = \log |x| + C$$

which is the required solution. (1)

5. Find the particular solution of the differential equation $\left\{ x \sin^2 \left(\frac{y}{x} \right) - y \right\} dx + x dy = 0$, given

that $y = \frac{\pi}{4}$, when $x = 1$.

All India 2014C

Given differential equation is

$$\left[x \sin^2\left(\frac{y}{x}\right) - y \right] dx + x dy = 0$$

$$\Rightarrow \frac{dy}{dx} = \frac{y - x \sin^2\left(\frac{y}{x}\right)}{x} \quad \dots(i)$$

which is a homogeneous differential equation.

Put $y = vx \Rightarrow \frac{dy}{dx} = v + x \frac{dv}{dx}$ in Eq. (i), we get

$$\begin{aligned} v + x \frac{dv}{dx} &= \frac{vx - x \sin^2\left(\frac{vx}{x}\right)}{x} \\ \Rightarrow v + x \frac{dv}{dx} &= v - \sin^2 v \Rightarrow x \frac{dv}{dx} = -\sin^2 v \\ \Rightarrow \operatorname{cosec}^2 v \, dv &= -\frac{dx}{x} \quad (1) \end{aligned}$$

On integrating both sides, we get

$$\begin{aligned} \int \operatorname{cosec}^2 v \, dv + \int \frac{dx}{x} &= 0 \\ \Rightarrow -\cot v + \log|x| &= C \\ \Rightarrow -\cot\left(\frac{y}{x}\right) + \log|x| &= C \quad \left[\because v = \frac{y}{x} \right] \quad \dots(ii) \end{aligned}$$

(1)

Also, given that $y = \frac{\pi}{4}$, when $x = 1$.

On putting $x = 1$ and $y = \frac{\pi}{4}$ in Eq. (ii), we get

$$-\cot\left(\frac{\pi}{4}\right) + \log 1 = C$$

$$\Rightarrow C = -1 \quad \left[\because \cot \frac{\pi}{4} = 1 \right] \quad (1)$$

On putting this value of C in Eq. (ii), we get

$$-\cot\left(\frac{y}{x}\right) + \log |x| = 1$$

$$\Rightarrow 1 + \log |x| - \cot\left(\frac{y}{x}\right) = 0$$

which is the required particular solution of given differential equation. (1)

6. Find the particular solution of the differential equation

$$\frac{dy}{dx} = \frac{x(2\log x + 1)}{\sin y + y \cos y}, \text{ given that } y = \frac{\pi}{2}, \text{ when } x = 1.$$

Delhi 2014

Given differential equation is

$$\frac{dy}{dx} = \frac{x(2 \log x + 1)}{\sin y + y \cos y}$$

On separating the variables, we get

$$(\sin y + y \cos y) dy = x(2 \log x + 1) dx$$

$$\Rightarrow \sin y dy + y \cos y dy = 2x \log x dx + x dx \quad (1)$$

On integrating both sides, we get

$$\begin{aligned} \int \sin y dy + \int y \cos y dy \\ = 2 \int x \log x dx + \int x dx \end{aligned}$$

$$\begin{aligned} \Rightarrow -\cos y + \left[y \int \cos y dy \right. \\ \left. - \int \left\{ \frac{d}{dy} (y) \int \cos y dy \right\} dy \right] \\ = 2 \left[\log x \int x dx - \int \left\{ \frac{d}{dx} (\log x) \int x dx \right\} dx \right] + \frac{x^2}{2} \end{aligned} \quad (1)$$

$$\begin{aligned} \Rightarrow -\cos y + y \sin y - \int \sin y dy \\ = 2 \left[\frac{x^2}{2} \log x - \int \left\{ \frac{1}{x} \frac{x^2}{2} \right\} dx \right] + \frac{x^2}{2} \end{aligned}$$

$$\begin{aligned} \Rightarrow -\cos y + y \sin y + \cos y \\ = x^2 \log x - \int x dx + \frac{x^2}{2} \end{aligned}$$

$$\Rightarrow y \sin y = x^2 \log x - \frac{x^2}{2} + \frac{x^2}{2} + C$$

$$\Rightarrow y \sin y = x^2 \log x + C \quad \dots(i) \quad (1)$$

Also, given that $y = \frac{\pi}{2}$, when $x = 1$.

On putting $y = \frac{\pi}{2}$ and $x = 1$ in Eq. (i), we get

$$\frac{\pi}{2} \sin \left(\frac{\pi}{2} \right) = (1)^2 \log (1) + C$$

$$\Rightarrow C = \frac{\pi}{2} \quad \left[\because \sin \frac{\pi}{2} = 1, \log 1 = 0 \right]$$

On substituting the value of C in Eq. (i), we get

$$y \sin y = x^2 \log x + \frac{\pi}{2}$$

which is the required particular solution. **(1)**

7. Solve the following differential equation

$$(x^2 - 1) \frac{dy}{dx} + 2xy = \frac{2}{x^2 - 1}.$$

Delhi 2014; All India 2014C

💡 Firstly, divide the given differential equation by $(x^2 - 1)$ to convert it into the form of linear differential equation and then solve it.

Given differential equation is

$$(x^2 - 1) \frac{dy}{dx} + 2xy = \frac{2}{x^2 - 1}$$

On dividing both sides by $(x^2 - 1)$, we get

$$\frac{dy}{dx} + \frac{2x}{x^2 - 1} y = \frac{2}{(x^2 - 1)^2}$$

which is a linear differential equation. (1)

On comparing with the form $\frac{dy}{dx} + Py = Q$, we

get $P = \frac{2x}{x^2 - 1}, Q = \frac{2}{(x^2 - 1)^2}$

$$\therefore \text{IF} = e^{\int \frac{2x}{x^2 - 1} dx} \quad (1)$$

$$= e^{\log|x^2 - 1|} = x^2 - 1$$

$$\left[\begin{array}{l} \text{put } x^2 - 1 = t \Rightarrow 2x dx = dt \text{ in } \int \frac{2x}{x^2 - 1} dx, \text{ then} \\ \int \frac{2x}{x^2 - 1} dx = \int \frac{1}{t} dt = \log t = \log(x^2 - 1) \end{array} \right]$$

Hence, the required general solution is

$$y \cdot \text{IF} = \int Q \times \text{IF} dx + C$$

$$\Rightarrow y(x^2 - 1) = \int \frac{2}{(x^2 - 1)^2} \times (x^2 - 1) dx + C \quad (1)$$

$$\Rightarrow y(x^2 - 1) = \int \frac{2}{x^2 - 1} dx + C$$

$$\Rightarrow y(x^2 - 1) = \log \left| \frac{x-1}{x+1} \right| + C$$

$$\left[\because \int \frac{1}{x^2 - a^2} dx = \frac{1}{2a} \log \left| \frac{x-a}{x+a} \right| \right]$$

which is the required differential equation. (1)

- 8.** Find the particular solution of the differential equation $e^x \sqrt{1-y^2} dx + \frac{y}{x} dy = 0$, given that

$y = 1$, when $x = 0$.

Delhi 2014

Given differential equation is

$$e^x \sqrt{1-y^2} dx + \frac{y}{x} dy = 0$$

$$\Rightarrow e^x \sqrt{1-y^2} dx = \frac{-y}{x} dy$$

On separating the variables, we get

$$\frac{-y}{\sqrt{1-y^2}} dy = x e^x dx \quad (1)$$

On integrating both sides, we get

$$\int \frac{-y}{\sqrt{1-y^2}} dy = \int x e^x dx$$

On putting $1-y^2 = t \Rightarrow -y dy = \frac{dt}{2}$ in LHS, we get

$$\int \frac{1}{2\sqrt{t}} dt = \int x e^x dx$$

$$\Rightarrow \frac{1}{2} [2\sqrt{t}] = x \int e^x dx - \int \left[\frac{d}{dx}(x) \int e^x dx \right] dx$$

$$\Rightarrow \sqrt{1-y^2} = x e^x - \int e^x dx \quad [\because t = 1-y^2] \quad (1)$$

$$\Rightarrow \sqrt{1-y^2} = x e^x - e^x + C \quad \dots(i)$$

Also, given that $y = 1$, when $x = 0$

On putting $y = 1$ and $x = 0$ in Eq. (i), we get

$$\sqrt{1-1} = 0 - e^0 + C$$

$$\Rightarrow C = 1 \quad [\because e^0 = 1] \quad (1)$$

On substituting the value of C in Eq. (i), we get

$$\sqrt{1-y^2} = x e^x - e^x + 1$$

which is the required particular solution of given differential equation. (1)

9. Solve the following differential equation

$$\operatorname{cosec} x \log y \frac{dy}{dx} + x^2 y^2 = 0. \quad \text{Delhi 2014}$$



Firstly, separate the variables, then integrate by using integration by parts.

Given differential equation is

$$\operatorname{cosec} x \log y \frac{dy}{dx} + x^2 y^2 = 0 \quad \dots(i)$$

It can be rewritten as

$$\operatorname{cosec} x \log y \frac{dy}{dx} = -x^2 y^2$$

On separating the variables, we get

$$\frac{\log y}{y^2} dy = \frac{-x^2}{\operatorname{cosec} x} dx$$

On integrating both sides, we get

$$\int \frac{\log y}{y^2} dy = - \int \frac{x^2}{\operatorname{cosec} x} dx \Rightarrow I_1 = I_2 \quad \dots(ii)$$

(1)

where, $I_1 = \int \frac{\log y}{y^2} dy$

Put $\log y = t \Rightarrow y = e^t$, then $\frac{dy}{y} = dt$

$$\begin{aligned} \therefore I_1 &= \int t e^{-t} dt \\ &= t \int e^{-t} dt - \int \left[\frac{d}{dt}(t) \int e^{-t} dt \right] dt \\ &= -t e^{-t} - \int (-e^{-t}) dt \\ &= -t e^{-t} + \int e^{-t} dt = -t e^{-t} - e^{-t} + C_1 \\ &= -\frac{\log y}{y} - \frac{1}{y} + C_1 \quad \dots(iii) \quad (1) \end{aligned}$$

$$\left[\because t = \log y \text{ and } e^{-t} = \frac{1}{y} \right]$$

$$\left[\frac{1}{y} - \log y, \text{ and } \frac{1}{y} - y \right]$$

$$\begin{aligned} \text{and } I_2 &= -\int \frac{x^2}{\operatorname{cosec} x} dx \\ &= -\int x^2 \sin x dx \\ &= -x^2 \int \sin x dx - \int \left[\frac{d}{dx}(x^2) \int \sin x dx \right] dx \\ &= -x^2 (-\cos x) - \int [2x(-\cos x)] dx \\ &= x^2 \cos x + 2 \int x \cos x dx \\ &= x^2 \cos x + 2 \left[x \int \cos x dx \right. \\ &\quad \left. - \int \left\{ \frac{d}{dx}(x) \int \cos x dx \right\} dx \right] \\ &= x^2 \cos x + 2 [x \sin x - \int \sin x dx] \\ &= x^2 \cos x + 2x \sin x + 2 \cos x + C_2 \quad \dots \text{(iv)} \end{aligned}$$

(1)

On putting the values of I_1 and I_2 from Eqs.(iii) and (iv) in Eq. (ii), we get

$$\begin{aligned} -\frac{\log y}{y} - \frac{1}{y} + C_1 &= x^2 \cos x + 2x \sin x \\ &\quad + 2 \cos x + C_2 \\ \Rightarrow -\frac{(1 + \log y)}{y} &= x^2 \cos x + 2x \sin x \\ &\quad + 2 \cos x + C_2 - C_1 \\ \Rightarrow -\frac{(1 + \log y)}{y} &= x^2 \cos x + 2x \sin x \\ &\quad + 2 \cos x + C \end{aligned}$$

where, $C = C_2 - C_1$

which is the required solution of given differential equation. **(1)**

- 10.** Find the particular solution of the differential equation $x(1 + y^2) dx - y(1 + x^2) dy = 0$, given that $y = 1$, when $x = 0$. All India 2014

Given differential equation is

$$\begin{aligned} x(1 + y^2) dx - y(1 + x^2) dy &= 0 \\ \Rightarrow x(1 + y^2) dx &= y(1 + x^2) dy \end{aligned}$$

On separating the variables, we get

$$\frac{y}{(1 + y^2)} dy = \frac{x}{(1 + x^2)} dx \quad (1)$$

On integrating both sides, we get

$$\begin{aligned} \int \frac{y}{1 + y^2} dy &= \int \frac{x}{(1 + x^2)} dx \\ \Rightarrow \frac{1}{2} \log |1 + y^2| &= \frac{1}{2} \log |1 + x^2| + C \quad \dots(i) \end{aligned}$$

$$\left[\begin{array}{l} \text{let } 1 + y^2 = u \Rightarrow 2y dy = du, \\ \text{then } \int \frac{y}{1 + y^2} dy = \int \frac{1}{2u} du = \frac{1}{2} \log |u| \\ \text{and let } 1 + x^2 = v \Rightarrow 2x dx = dv, \\ \text{then } \int \frac{x}{1 + x^2} dx = \frac{1}{2} \int \frac{1}{v} dv = \frac{1}{2} \log |v| \end{array} \right]$$

Also, given that $y = 1$, when $x = 0$. (1)

On substituting the values of x and y in Eq. (i), we get

$$\begin{aligned} \frac{1}{2} \log |1 + (1)^2| &= \frac{1}{2} \log |1 + (0)^2| + C \\ \Rightarrow \frac{1}{2} \log 2 &= C \quad [\because \log 1 = 0] \end{aligned}$$

On putting $C = \frac{1}{2} \log 2$ in Eq. (i), we get

$$\frac{1}{2} \log |1 + y^2| = \frac{1}{2} \log |1 + x^2| + \frac{1}{2} \log 2$$

$$\Rightarrow \log |1 + y^2| = \log |1 + x^2| + \log 2 \quad (1)$$

$$\Rightarrow \log |1 + y^2| - \log |1 + x^2| = \log 2$$

$$\Rightarrow \log \left| \frac{1+y^2}{1+x^2} \right| = \log 2 \left[\because \log m - \log n = \log \frac{m}{n} \right]$$

$$\Rightarrow \frac{1+y^2}{1+x^2} = 2$$

$$\Rightarrow 1 + y^2 = 2 + 2x^2 \Rightarrow y^2 - 2x^2 - 1 = 0$$

which is the required particular solution of given differential equation. (1)

11. Find the particular solution of the differential equation $\log \left(\frac{dy}{dx} \right) = 3x + 4y$ equation, given

that $y = 0$, when $x = 0$.

All India 2014

Given differential equation is

$$\log \left(\frac{dy}{dx} \right) = 3x + 4y$$

$$\Rightarrow \frac{dy}{dx} = e^{3x+4y}$$

$$[\because \log m = n \Rightarrow e^n = m]$$

$$\Rightarrow \frac{dy}{dx} = e^{3x} e^{4y} \quad (1)$$

On separating the variables, we get

$$\frac{1}{e^{4y}} dy = e^{3x} dx$$

On integrating both sides, we get

$$\int e^{-4y} dy = \int e^{3x} dx$$

$$\Rightarrow \frac{e^{-4y}}{-4} = \frac{e^{3x}}{3} + C \quad \dots(i) \quad (1)$$

Also, given that $y = 0$, when $x = 0$.

On putting $y = 0$ and $x = 0$ in Eq. (i), we get

$$\frac{e^{-4(0)}}{-4} = \frac{e^{3(0)}}{3} + C$$

$$\Rightarrow -\frac{1}{4} = \frac{1}{3} + C \quad [\because e^{-0} = e^0 = 1]$$

$$\Rightarrow C = -\frac{1}{4} - \frac{1}{3}$$

$$\therefore C = \frac{-7}{12} \quad (1)$$

On substituting the value of C in Eq. (i), we get

$$\frac{e^{-4y}}{-4} = \frac{e^{3x}}{3} - \frac{7}{12}$$

which is the required particular solution of given differential equation. (1)

12. Solve the differential equation

$$(1+x^2) \frac{dy}{dx} + y = e^{\tan^{-1} x}.$$

All India 2014

Given differential equation is

$$(1+x^2) \frac{dy}{dx} + y = e^{\tan^{-1} x}$$

On dividing both sides by $(1+x^2)$, we get

$$\frac{dy}{dx} + \frac{1}{(1+x^2)} y = \frac{e^{\tan^{-1} x}}{1+x^2}$$

It is a linear differential equation of the form

$$\frac{dy}{dx} + Py = Q$$

On comparing, we get

$$P = \frac{1}{1+x^2} \text{ and } Q = \frac{e^{\tan^{-1} x}}{1+x^2}$$

$$\begin{aligned} \therefore \text{IF} &= e^{\int P dx} = e^{\int \frac{1}{1+x^2} dx} = e^{\tan^{-1} x} \\ &\left[\because \int \frac{1}{1+x^2} dx = \tan^{-1} x \right] \quad (1) \end{aligned}$$

Then, required solution is

$$\begin{aligned} (y \cdot \text{IF}) &= \int (Q \cdot \text{IF}) dx + C \\ \therefore y e^{\tan^{-1} x} &= \int \frac{e^{\tan^{-1} x} \cdot e^{\tan^{-1} x}}{1+x^2} dx + C \\ \Rightarrow y e^{\tan^{-1} x} &= \int \frac{e^{2 \tan^{-1} x}}{1+x^2} dx + C \\ \Rightarrow y e^{\tan^{-1} x} &= I + C \quad \dots(i) \quad (1) \end{aligned}$$

$$\text{where, } I = \int \frac{e^{2 \tan^{-1} x}}{1+x^2} dx$$

$$\text{Put } \tan^{-1} x = t \Rightarrow \frac{1}{1+x^2} dx = dt$$

$$\begin{aligned} \therefore I &= \int_{-2t} e^{2t} dt \\ &\quad \quad \quad -2 \tan^{-1} x \end{aligned}$$

$$\Rightarrow I = \frac{e^{-}}{2} \Rightarrow I = \frac{e^{-2 \tan^{-1} x}}{2} \quad (1)$$

On putting the value of I in Eq. (i), we get

$$y e^{\tan^{-1} x} = \frac{e^{2 \tan^{-1} x}}{2} + C$$

which is the required general solution of given differential equation. (1)

13. Find a particular solution of the differential equation $\frac{dy}{dx} + 2y \tan x = \sin x$, given that

$$y = 0, \text{ when } x = \frac{\pi}{3}. \quad \text{Foreign 2014}$$

Given differential equation is

$$\frac{dy}{dx} + 2y \tan x = \sin x$$

which is a linear differential equation of the form $\frac{dy}{dx} + Py = Q$.

On comparing, we get

$$P = 2 \tan x \text{ and } Q = \sin x$$

$$\therefore \text{IF} = e^{2 \int \tan x \, dx} = e^{2 \log |\sec x|} \quad (1)$$

$$= e^{\log \sec^2 x} \quad [\because m \log n = \log n^m]$$

$$= \sec^2 x \quad [\because e^{\log x} = x]$$

The general solution is given by

$$Y \cdot \text{IF} = \int Q \times \text{IF} \, dx + C \quad \dots(i) \quad (1)$$

$$\Rightarrow y \sec^2 x = \int (\sin x \cdot \sec^2 x) \, dx + C$$

$$\Rightarrow y \sec^2 x = \int \sin x \cdot \frac{1}{\cos^2 x} \, dx + C$$

$$\Rightarrow y \sec^2 x = \int \tan x \sec x \, dx + C$$

$$\Rightarrow y \sec^2 x = \sec x + C \quad \dots(ii)$$

Also, given that $y = 0$, when $x = \frac{\pi}{3}$. On putting

$y = 0$ and $x = \frac{\pi}{3}$ in Eq. (ii), we get

$$0 \times \sec^2 \frac{\pi}{3} = \sec \frac{\pi}{3} + C$$

$$\Rightarrow 0 = 2 + C \Rightarrow C = -2 \quad (1)$$

On putting the value of C in Eq. (ii), we get

$$y \sec^2 x = \sec x - 2$$

$$\Rightarrow y = \cos x - 2 \cos^2 x$$

which is the required solution of the given differential equation. (1)

14. Solve the following differential equation

$$x \cos \left(\frac{y}{x} \right) \frac{dy}{dx} = y \cos \left(\frac{y}{x} \right) + x; x \neq 0.$$

All India 2014C

Given differential equation is

$$x \cos\left(\frac{y}{x}\right) \frac{dy}{dx} = y \cos\left(\frac{y}{x}\right) + x \quad \dots(i)$$

which is a homogeneous differential equation.

On putting $y = vx \Rightarrow \frac{dy}{dx} = v + x \frac{dv}{dx}$ in

Eq. (i), we get

$$x \cos v \left[v + x \frac{dv}{dx} \right] = vx \cos v + x$$

$$\Rightarrow v + x \frac{dv}{dx} = \frac{x(v \cos v + 1)}{x \cos v} \quad (1)$$

$$\Rightarrow x \frac{dv}{dx} = \frac{v \cos v + 1}{\cos v} - v$$

$$\Rightarrow x \frac{dv}{dx} = \frac{v \cos v + 1 - v \cos v}{\cos v}$$

$$\Rightarrow x \frac{dv}{dx} = \frac{1}{\cos v} \Rightarrow \cos v dv = \frac{dx}{x} \quad (1)$$

On integrating both sides, we get

$$\int \cos v dv = \int \frac{dx}{x}$$

$$\Rightarrow \sin v = \log x + C \quad (1)$$

$$\Rightarrow \sin\left(\frac{y}{x}\right) = \log x + C \left[\because y = vx \Rightarrow v = \frac{y}{x} \right]$$

which is the required solution of given differential equation. (1)

15. If $y(x)$ is a solution of the differential equation

$$\left(\frac{2 + \sin x}{1 + y} \right) \frac{dy}{dx} = -\cos x \text{ and } y(0) = 1, \text{ then find}$$

the value of $y\left(\frac{\pi}{2}\right)$.

Delhi 2014C

Given differential equation is

$$\left(\frac{2 + \sin x}{1 + y} \right) \frac{dy}{dx} = -\cos x$$
$$\Rightarrow \frac{1}{1 + y} dy = -\frac{\cos x}{2 + \sin x} dx \quad (1)$$

Now, on integrating both sides, we get

$$\int \frac{1}{1 + y} dy = - \int \frac{\cos x}{2 + \sin x} dx$$
$$\Rightarrow \log |1 + y| = -\log |2 + \sin x| + \log C$$

$$\left[\begin{array}{l} \text{for } \int \frac{\cos x}{2 + \sin x} dx, \text{ let } 2 + \sin x = t \\ \Rightarrow \cos x dx = dt, \\ \text{then } \int \frac{\cos x}{2 + \sin x} dx = \int \frac{dt}{t} = \log t + C \\ = \log |2 + \sin x| + C \end{array} \right]$$

$$\Rightarrow \log (1 + y) + \log (2 + \sin x) = \log C$$

$$\Rightarrow \log (1 + y) (2 + \sin x) = \log C$$

$$\Rightarrow (1 + y) (2 + \sin x) = C \quad \dots(i)$$

Also, given that at $x = 0, y(0) = 1$

On putting $x = 0$ and $y = 1$ in Eq. (i), we get

$$(1 + 1) (2 + \sin 0) = C$$
$$\Rightarrow C = 4 \quad (1)$$

On putting $C = 4$ in Eq. (i), we get

$$(1 + y) (2 + \sin x) = 4$$

$$\Rightarrow 1 + y = \frac{4}{2 + \sin x}$$

$$\Rightarrow y = \frac{4}{2 + \sin x} - 1$$

$$\Rightarrow y = \frac{4 - 2 - \sin x}{2 + \sin x}$$

$$\Rightarrow y = \frac{2 - \sin x}{2 + \sin x} \quad (1)$$

$$\text{Now, at } x = \frac{\pi}{2}, y\left(\frac{\pi}{2}\right) = \frac{2 - \sin \frac{\pi}{2}}{2 + \sin \frac{\pi}{2}}$$

$$\therefore y\left(\frac{\pi}{2}\right) = \frac{1}{3} \quad \left[\because \sin \frac{\pi}{2} = 1 \right] \quad (1)$$

16. Solve the differential equation

$$x \frac{dy}{dx} + y = x \cos x + \sin x, \text{ given } y\left(\frac{\pi}{2}\right) = 1.$$

All India 2014C

Given differential equation is

$$x \frac{dy}{dx} + y = x \cos x + \sin x$$

$$\Rightarrow \frac{dy}{dx} + \frac{y}{x} = \cos x + \frac{\sin x}{x}$$

[dividing on both sides by x]

which is a linear differential equation.

On comparing with the form $\frac{dy}{dx} + Py = Q$,

$$\text{we get } P = \frac{1}{x} \text{ and } Q = \cos x + \frac{\sin x}{x}$$

$$\therefore \text{IF} = e^{\int P dx} = e^{\int \frac{1}{x} dx} = e^{\log x} = x$$

The general solution is given by

$$y \cdot \text{IF} = \int Q \times \text{IF} dx + C \quad (1)$$

$$\Rightarrow yx = \int x \left(\cos x + \frac{\sin x}{x} \right) dx + C$$

$$\Rightarrow yx = \int (x \cos x + \sin x) dx + C$$

$$\Rightarrow xy = \int x \cos x dx + \int \sin x dx + C$$

$$\Rightarrow xy = x \int \cos x dx - \int \left[\frac{d}{dx}(x) \int \cos x dx \right] dx + \int \sin x dx + C$$

$$\Rightarrow xy = x \sin x + \cos x - \cos x + C$$

$$\Rightarrow xy = x \sin x + C$$

$$\Rightarrow y = \sin x + C \cdot \frac{1}{x} \quad \dots(i) \quad (1)$$

Also, given that at $x = \frac{\pi}{2}; y = 1$

On putting $x = \frac{\pi}{2}$ and $y = 1$ in Eq. (i), we get

$$1 = 1 + C \cdot \frac{2}{\pi} \Rightarrow C = 0 \quad (1)$$

On putting the value of C in Eq. (i), we get

$$y = \sin x$$

which is the required solution of given differential equation. (1)

17. Solve the differential equation

$$\frac{dy}{dx} + y \cot x = 2 \cos x, \text{ given that } y = 0, \text{ when}$$

$$x = \frac{\pi}{2}.$$

Foreign 2014

Given differential equation is

$$\frac{dy}{dx} + y \cot x = 2 \cos x$$

which is a linear differential equation of the form

$$\frac{dy}{dx} + Py = Q$$

Here, $P = \cot x$ and $Q = 2 \cos x$

$$\therefore \text{IF} = e^{\int P dx} = e^{\int \cot x dx} = e^{\log \sin x}$$

$$\Rightarrow \text{IF} = \sin x \quad (1)$$

The general solution is given by

$$Y \times \text{IF} = \int \text{IF} \times Q dx + C$$

$$\Rightarrow y \sin x = \int 2 \sin x \cos x dx + C$$

$$\Rightarrow y \sin x = \int \sin 2x dx + C$$

$$\Rightarrow y \sin x = -\frac{\cos 2x}{2} + C \quad \dots(i) \quad (1)$$

Also, given that $y = 0$, when $x = \frac{\pi}{2}$.

On putting $x = \frac{\pi}{2}$ and $y = 0$ in Eq. (i), we get

$$0 \sin \frac{\pi}{2} = -\frac{\cos 2 \frac{\pi}{2}}{2} + C$$

$$\Rightarrow C - \frac{\cos \pi}{2} = 0 \Rightarrow C + \frac{1}{2} = 0$$

$$\therefore C = -\frac{1}{2} \quad (1)$$

On putting the value of C in Eq. (i), we get

$$y \sin x = -\cos \frac{2x}{2} - \frac{1}{2}$$

$$\Rightarrow 2y \sin x + \cos 2x + 1 = 0$$

which is the required solution. (1)

- 18.** Solve the differential equation
 $(x^2 - yx^2) dy + (y^2 + x^2y^2) dx = 0$, given that
 $y = 1$, when $x = 1$. Foreign 2014

Direction (Q. Nos. 19-22) Solve the following differential equations.

Given differential equation is

$$(x^2 - yx^2)dy + (y^2 + x^2y^2) dx = 0$$

On dividing both sides by dx , we get

$$(x^2 - yx^2) \frac{dy}{dx} + (y^2 + x^2y^2) = 0$$

$$\Rightarrow x^2 (1 - y) \frac{dy}{dx} + y^2 (1 + x^2) = 0$$

$$\Rightarrow -x^2 (1 - y) \frac{dy}{dx} = y^2 (1 + x^2)$$

$$\Rightarrow x^2 (y - 1) \frac{dy}{dx} = y^2 (1 + x^2)$$

$$\Rightarrow \frac{y-1}{y^2} dy = \frac{1+x^2}{x^2} dx \quad (1)$$

On integrating both sides, we get

$$\int \frac{y-1}{y^2} dy = \int \frac{1+x^2}{x^2} dx$$

$$\Rightarrow \frac{1}{2} \int \frac{2y}{y^2} dy - \int \frac{1}{y^2} dy = \int \frac{1}{x^2} dx + \int 1 \cdot dx \quad (1)$$

On putting $y^2 = t \Rightarrow 2y dy = dt$ in first integral, we get

$$\frac{1}{2} \int \frac{dt}{t} + \frac{1}{y} = -\frac{1}{x} + x$$

$$\Rightarrow \frac{1}{2} \log |y^2| + \frac{1}{y} = -\frac{1}{x} + x + C \quad \dots(i)$$

$[\because t = y^2]$

Also, given that $y = 1$, when $x = 1$.

On putting $y = 1$ and $x = 1$ in Eq.(i), we get

$$\frac{1}{2} \log |1| + \frac{1}{1} = -\frac{1}{1} + 1 + C$$

$$\Rightarrow \frac{1}{2} \log |1| + 1 = -1 + 1 + C$$

$$\Rightarrow C = 1 \quad [\because \log 1 = 0] \quad (1)$$

On putting the value of C in Eq. (i), we get

$$\frac{1}{2} \log |y^2| + \frac{1}{y} = -\frac{1}{x} + x + 1$$

which is the required solution. (1)

19. $\frac{dy}{dx} + y \sec x = \tan x$ All India 2012C; Delhi 2008C

Given differential equation is

$$\frac{dy}{dx} + y \sec x = \tan x \quad \dots(i)$$

which is a linear differential equation of first order and is of the form

$$\frac{dy}{dx} + Py = Q \quad \dots(ii)$$

On comparing Eqs. (i) and (ii), we get

$$P = \sec x \text{ and } Q = \tan x \quad (1)$$

$$\therefore IF = e^{\int \sec x dx} = e^{\log |\sec x + \tan x|}$$

$$[\because \int \sec x dx = \log |\sec x + \tan x|]$$

$$\Rightarrow IF = \sec x + \tan x \quad (1)$$

The general solution is

$$y \times IF = \int Q \cdot IF dx + C$$

$$y (\sec x + \tan x) = \int \tan x \cdot (\sec x + \tan x) dx$$

$$\Rightarrow y (\sec x + \tan x) = \int \sec x \tan x dx + \int \tan^2 x dx$$

$$\Rightarrow y (\sec x + \tan x) = \sec x + \int (\sec^2 x - 1) dx \quad (1)$$

$$\Rightarrow y (\sec x + \tan x) = (\sec x + \tan x) - x + C$$
$$[\because \int \sec^2 x dx = \tan x + C]$$

On dividing both sides by $(\sec x + \tan x)$, we get the required solution as

$$y = 1 - \frac{x}{\sec x + \tan x} + \frac{C}{\sec x + \tan x} \quad (1)$$

20. $2x^2 \frac{dy}{dx} - 2xy + y^2 = 0$

Delhi 2012

Given differential equation is

$$2x^2 \frac{dy}{dx} - 2xy + y^2 = 0$$

$$\Rightarrow \frac{dy}{dx} = \frac{2xy - y^2}{2x^2} \quad \dots(i) \quad (1)$$

which is a homogeneous differential equation.

On putting $y = vx \Rightarrow \frac{dy}{dx} = v + x \frac{dv}{dx}$ in Eq. (i),

we get

$$v + x \frac{dv}{dx} = \frac{2vx^2 - v^2x^2}{2x^2} \quad (1)$$

$$\Rightarrow v + x \frac{dv}{dx} = \frac{2v - v^2}{2}$$

$$\Rightarrow x \frac{dv}{dx} = \frac{2v - v^2}{2} - v$$

$$\Rightarrow x \frac{dv}{dx} = \frac{2v - v^2 - 2v}{2}$$

$$\Rightarrow x \frac{dv}{dx} = \frac{-v^2}{2}$$

$$\Rightarrow \frac{2dv}{v^2} = -\frac{1}{x} dx \quad (1)$$

On integrating both sides, we get

$$\int \frac{2dv}{v^2} = \int \frac{-dx}{x} + C$$

$$\Rightarrow 2 \int v^{-2} dv = -\log|x| + C$$

$$\Rightarrow \frac{2v^{-1}}{-1} = -\log|x| + C$$

$$\Rightarrow \frac{-2}{v} = -\log|x| + C$$

$$\Rightarrow \frac{-2x}{y} = -\log|x| + C$$

$$\left[\because y = vx \Rightarrow v = \frac{y}{x} \right]$$

$$\Rightarrow -2x = y(-\log|x| + C)$$

$$\Rightarrow y = \frac{-2x}{-\log|x| + C}$$

which is the required solution.

(1)

21. $\frac{dy}{dx} = 1 + x^2 + y^2 + x^2y^2$, given that $y = 1$,
when $x = 0$.

Delhi 2012

Given differential equation is

$$\frac{dy}{dx} = 1 + x^2 + y^2 + x^2y^2 \quad (1)$$

$$\Rightarrow \frac{dy}{dx} = (1 + x^2) + y^2(1 + x^2)$$

$$\Rightarrow \frac{dy}{dx} = (1 + x^2)(1 + y^2)$$

$$\Rightarrow \frac{dy}{1 + y^2} = (1 + x^2) dx \quad (1)$$

On integrating both sides, we get

$$\int \frac{dy}{1 + y^2} = \int (1 + x^2) dx$$

$$\Rightarrow \tan^{-1} y = x + \frac{x^3}{3} + C \quad \dots(i)$$

Also, given that $y = 0$, when $x = 2$.

On putting $x = 0$ and $y = 1$ in Eq. (i), we get

$$\tan^{-1} 1 = C$$

$$\Rightarrow \tan^{-1}(\tan \pi/4) = C \quad \left[\because \tan \frac{\pi}{4} = 1 \right]$$

$$\Rightarrow C = \pi/4 \quad (1)$$

On putting the value of C in Eq. (i), we get

$$\tan^{-1} y = x + \frac{x^3}{3} + \frac{\pi}{4}$$

$$\Rightarrow y = \tan \left(x + \frac{x^3}{3} + \frac{\pi}{4} \right)$$

which is the required solution. (1)

22. $x(x^2 - 1) \frac{dy}{dx} = 1$, $y = 0$, when $x = 2$.

All India 2012

Given differential equation is

$$\begin{aligned}x(x^2 - 1) \frac{dy}{dx} &= 1 \\ \Rightarrow \frac{dy}{dx} &= \frac{1}{x(x^2 - 1)} \\ \Rightarrow \frac{dy}{dx} &= \frac{1}{x(x-1)(x+1)} \\ &[\because a^2 - b^2 = (a-b)(a+b)] \\ \Rightarrow dy &= \frac{dx}{x(x-1)(x+1)}\end{aligned}$$

On integrating both sides, we get

$$\begin{aligned}\int dy &= \int \frac{dx}{x(x-1)(x+1)} + C \\ \Rightarrow y &= I + C \quad \dots(i) \\ \text{where, } I &= \int \frac{dx}{x(x-1)(x+1)} \quad (1)\end{aligned}$$

$$\text{Let } \frac{1}{x(x-1)(x+1)} = \frac{A}{x} + \frac{B}{x-1} + \frac{C}{x+1}$$

$$\Rightarrow 1 = A(x-1)(x+1) + Bx(x+1) + Cx(x-1)$$

On comparing coefficients of x^2 , x and constant terms from both sides, we get

$$A + B + C = 0 \quad \dots(ii)$$

$$B - C = 0 \quad \dots(iii)$$

$$\text{and } -A = 1$$

$$\Rightarrow A = -1$$

On putting $A = -1$ in Eq. (ii), we get

$$B + C = 1 \quad \dots(iv)$$

Now, on adding Eqs. (iii) and (iv), we get

$$2B = 1 \Rightarrow B = \frac{1}{2}$$

On putting $B = \frac{1}{2}$ in Eq. (iii), we get

$$\frac{1}{2} - C = 0 \Rightarrow C = \frac{1}{2}$$

$$\therefore A = -1, B = \frac{1}{2} \text{ and } C = \frac{1}{2},$$

$$\text{then } \frac{1}{x(x-1)(x+1)} = \frac{-1}{x} + \frac{1/2}{x-1} + \frac{1/2}{x+1} \quad (1)$$

On integrating both sides w.r.t. x , we get

$$I = \int \frac{1}{x(x-1)(x+1)} dx = \int \frac{-1}{x} dx$$

$$+ \frac{1}{2} \int \frac{dx}{x-1} + \frac{1}{2} \int \frac{dx}{x+1}$$

$$\Rightarrow I = -\log|x| + \frac{1}{2} \log|x-1| + \frac{1}{2} \log|x+1|$$

On putting the value of I in Eq. (i), we get

$$y = -\log|x| + \frac{1}{2} \log|x-1| + \frac{1}{2} \log|x+1| + C$$

... (v)

Also, given that $y = 0$, when $x = 2$.

On putting $y = 0$ and $x = 2$ in Eq. (v), we get

$$0 = -\log 2 + \frac{1}{2} \log 1 + \frac{1}{2} \log 3 + C$$

$$\Rightarrow C = \log 2 - \frac{1}{2} \log 1 - \frac{1}{2} \log 3$$

$$\Rightarrow C = \log 2 - \log \sqrt{3} \quad [\because \log 1 = 0]$$

$$\Rightarrow C = \log \frac{2}{\sqrt{3}} \quad (1)$$

On putting the value of C in Eq. (v), we get

$$y = -\log|x| + \frac{1}{2} \log|x-1|$$

$$+ \frac{1}{2} \log|x+1| + \log \frac{2}{\sqrt{3}} \quad (1)$$

which is the required solution.

23. Solve the following differential equation

$$\frac{dy}{dx} + y \cot x = 4x \operatorname{cosec} x, \text{ given that } y = 0,$$

$$\text{when } x = \frac{\pi}{2}.$$

Delhi 2012C; Foreign 2011

Given differential equation is

$$\frac{dy}{dx} + y \cot x = 4x \operatorname{cosec} x$$

which is a linear differential equation.

On comparing with general form of linear differential equation of 1st order

$$\frac{dy}{dx} + Py = Q, \text{ we get}$$

$$P = \cot x \text{ and } Q = 4x \operatorname{cosec} x \quad (1)$$

$$\begin{aligned} \therefore \text{IF} &= e^{\int P dx} = e^{\int \cot x dx} \\ &= e^{\log \sin x} = \sin x \quad [\because e^{\log x} = x] \end{aligned}$$

$$\Rightarrow \text{IF} = \sin x \quad (1)$$

Now, solution of linear differential equation is given by

$$y \times \text{IF} = \int (Q \times \text{IF}) dx + C$$

On putting $\text{IF} = \sin x$ and $Q = 4x \operatorname{cosec} x$, we get

$$y \times \sin x = \int 4x \operatorname{cosec} x \cdot \sin x dx + C$$

$$\Rightarrow y \sin x = \int 4x \cdot \frac{1}{\sin x} \cdot \sin x dx + C$$

$$\Rightarrow y \sin x = \int 4x dx + C$$

$$\Rightarrow y \sin x = 2x^2 + C \quad \dots(i) \quad (1)$$

Also, given that $y = 0$, when $x = \frac{\pi}{2}$.

On putting $y = 0$ and $x = \frac{\pi}{2}$ in Eq. (i), we get

$$0 = 2 \times \frac{\pi^2}{4} + C \Rightarrow C = \frac{-\pi^2}{2}$$

On putting $C = -\frac{\pi^2}{2}$ in Eq. (i), we get

$$y \sin x = 2x^2 - \frac{\pi^2}{2}$$

$$\Rightarrow y = 2x^2 \operatorname{cosec} x - \frac{\pi^2}{2} \operatorname{cosec} x \quad (1)$$

which is the required solution.

24. Solve the following differential equation
 $(1+x^2) dy + 2xy dx = \cot x dx$, where $x \neq 0$.

All India 2012C, 2011

Given differential equation is

$$(1+x^2) dy + 2xy dx = \cot x dx \quad [\because x \neq 0]$$

$$\Rightarrow (1+x^2) dy = (\cot x - 2xy) dx$$

On dividing both sides by $1+x^2$, we get

$$dy = \frac{\cot x - 2xy}{1+x^2} dx$$

$$\Rightarrow \frac{dy}{dx} + \frac{2xy}{1+x^2} = \frac{\cot x}{1+x^2} \quad \dots(i) \quad (1)$$

which is a linear differential equation of 1st order and is of the form

$$\frac{dy}{dx} + Py = Q \quad \dots(ii)$$

On comparing Eqs. (i) and (ii), we get

$$P = \frac{2x}{1+x^2} \text{ and } Q = \frac{\cot x}{1+x^2}$$

$$\begin{aligned} \therefore \text{IF} &= e^{\int \frac{2x}{1+x^2} dx} \\ &= e^{\log|1+x^2|} = 1+x^2 \end{aligned} \quad (1)$$

$$\left[\text{for } \int \frac{2x}{1+x^2} dx, \text{ put } 1+x^2 = t \Rightarrow 2x dx = dt \right]$$

$$\left[\int \frac{dt}{t} = \log |t| = \log |1 + x^2| + C \right]$$

Now, solution of linear differential equation is given by

$$\begin{aligned} y \times IF &= \int (Q \times IF) dx + C \\ \therefore y(1 + x^2) &= \int \frac{\cot x}{1 + x^2} \times (1 + x^2) dx + C \\ \Rightarrow y(1 + x^2) &= \int \cot x dx + C \end{aligned} \quad (1)$$

$$\begin{aligned} \Rightarrow y(1 + x^2) &= \log |\sin x| + C \\ [\because \int \cot x dx &= \log |\sin x| + C] \end{aligned}$$

On dividing both sides by $1 + x^2$, we get

$$y = \frac{\log |\sin x|}{1 + x^2} + \frac{C}{1 + x^2}$$

which is the required solution. (1)

25. Find the particular solution of the differential equation

$$(1 + e^{2x})dy + (1 + y^2)e^x dx = 0, \text{ given that } y = 1, \text{ when } x = 0. \quad \text{Foreign 2011; All India 2008C}$$

Given differential equation is

$$(1 + e^{2x}) dy + (1 + y^2)e^x dx = 0$$

Above equation may be written as

$$\frac{dy}{1 + y^2} = \frac{-e^x}{1 + e^{2x}} dx \quad (1)$$

On integrating both sides, we get

$$\int \frac{dx}{1+y^2} = -\int \frac{e^x}{1+e^{2x}} dx$$

On putting $e^x = t \Rightarrow e^x dx = dt$ in RHS, we get

$$\tan^{-1} y = -\int \frac{1}{1+t^2} dt$$

$$\Rightarrow \tan^{-1} y = -\tan^{-1} t + C$$

$$\Rightarrow \tan^{-1} y = -\tan^{-1}(e^x) + C \quad \dots(i)$$

$[\because t = e^x](1\frac{1}{2})$

Now, given that $y = 1$, when $x = 0$.

On putting above values in Eq. (i), we get

$$\tan^{-1} 1 = -\tan^{-1}(e^0) + C$$

$$\Rightarrow \tan^{-1}\left(\tan \frac{\pi}{4}\right) = -\tan^{-1} 1 + C \quad [\because e^0 = 1]$$

$$\Rightarrow \frac{\pi}{4} = -\tan^{-1}\left(\tan \frac{\pi}{4}\right) + C$$

$$\Rightarrow \frac{\pi}{4} = -\frac{\pi}{4} + C$$

$$\Rightarrow C = \frac{\pi}{4} + \frac{\pi}{4} \Rightarrow C = \frac{\pi}{2}$$

On putting $C = \frac{\pi}{2}$ in Eq. (i), we get

$$\tan^{-1} y = -\tan^{-1} e^x + \frac{\pi}{2}$$

$$\Rightarrow y = \tan\left[\frac{\pi}{2} - \tan^{-1}(e^x)\right] = \cot[\tan^{-1}(e^x)]$$

$$= \cot\left[\cot^{-1}\left(\frac{1}{e^x}\right)\right] \quad \left[\because \tan^{-1} x = \cot^{-1} \frac{1}{x}\right]$$

$$\Rightarrow y = \frac{1}{e^x}$$

which is the required solution. (1½)

26. Solve the following differential equation

$$(1 + x^2) \frac{dy}{dx} + 2xy = \frac{1}{1 + x^2}, \text{ given that } y = 0,$$

when $x = 1$.

Foreign 2011

Given differential equation is

$$(1 + x^2) \frac{dy}{dx} + 2xy = \frac{1}{1 + x^2}$$

On dividing both sides by $(1 + x^2)$, we get

$$\frac{dy}{dx} + \frac{2xy}{1 + x^2} = \frac{1}{(1 + x^2)^2} \quad \dots(i)$$

which is a linear differential equation of the form

$$\frac{dy}{dx} + Py = Q \quad \dots(ii)$$

On comparing Eqs. (i) and (ii), we get

$$P = \frac{2x}{1+x^2} \text{ and } Q = \frac{1}{(1+x^2)^2} \quad (1)$$

$$\therefore \text{IF} = e^{\int \frac{2x}{1+x^2} dx} = e^{\log|1+x^2|} \quad (1)$$

$$\Rightarrow \text{IF} = 1+x^2 \quad [\because e^{\log x} = x]$$

$$\left[\begin{array}{l} \because \int \frac{2x}{1+x^2} dx, \text{ put } 1+x^2 = t \Rightarrow 2x dx = dt \\ \therefore \int \frac{dt}{t} = \log|t| = \log|1+x^2| \end{array} \right]$$

Now, solution of linear equation is given by

$$y \times \text{IF} = \int (Q \times \text{IF}) dx + C \quad \dots(iii)$$

$$\therefore y(1+x^2) = \int \frac{1}{(1+x^2)^2} \times (1+x^2) dx + C$$

$$\Rightarrow y(1+x^2) = \int \frac{1}{1+x^2} dx + C$$

$$\Rightarrow y(1+x^2) = \tan^{-1} x + C \quad \dots(iv) \quad (1)$$

$$\left[\because \int \frac{1}{1+x^2} dx = \tan^{-1} x + C \right]$$

Also, given that $y = 0$, when $x = 1$.

On putting $y = 0$ and $x = 1$ in Eq. (iv), we get

$$0 = \tan^{-1} 1 + C$$

$$\Rightarrow 0 = \tan^{-1} \left(\tan \frac{\pi}{4} \right) + C \quad \left[\because 1 = \tan \frac{\pi}{4} \right]$$

$$\Rightarrow 0 = \frac{\pi}{4} + C \Rightarrow C = -\frac{\pi}{4}$$

On putting $C = -\frac{\pi}{4}$ in Eq. (iv), we get

$$y(1+x^2) = \tan^{-1} x - \frac{\pi}{4}$$

$$\Rightarrow y = \frac{\tan^{-1} x}{1+x^2} - \frac{\pi}{4(1+x^2)} \quad (1)$$

which is the required solution.

27. Solve the following differential equation

$$x dy - y dx = \sqrt{x^2 + y^2} dx.$$

All India 2011

Given differential equation is

$$x dy - y dx = \sqrt{x^2 + y^2} dx$$

$$\Rightarrow (y + \sqrt{x^2 + y^2}) dx = x dy$$

$$\Rightarrow \frac{dy}{dx} = \frac{y + \sqrt{x^2 + y^2}}{x} \quad \dots(i) \quad (1)$$

which is a homogeneous differential equation because each term have same degree.

$$\text{On putting } y = vx \Rightarrow \frac{dy}{dx} = v + x \frac{dv}{dx} \quad (1)$$

in Eq. (i), we get

$$v + x \frac{dv}{dx} = \frac{vx + \sqrt{x^2 + v^2 x^2}}{x} = \frac{vx + x\sqrt{1 + v^2}}{x}$$

$$\Rightarrow v + x \frac{dv}{dx} = v + \sqrt{1 + v^2}$$

$$\Rightarrow x \frac{dv}{dx} = \sqrt{1 + v^2} \Rightarrow \frac{dv}{\sqrt{1 + v^2}} = \frac{dx}{x}$$

On integrating both sides, we get

$$\int \frac{dv}{\sqrt{1 + v^2}} = \int \frac{dx}{x}$$

$$\Rightarrow \log |v + \sqrt{1 + v^2}| = \log |x| + C$$

$$\left[\because \int \frac{dx}{\sqrt{a^2 + x^2}} = \log |x + \sqrt{x^2 + a^2}| \right]$$

$$\text{and } \int \frac{dx}{x} = \log |x| + C \quad (1)$$

$$\Rightarrow \log \left| \frac{y}{x} + \sqrt{1 + \frac{y^2}{x^2}} \right| = \log |x| + C \quad \left[\because y = vx \right]$$

$$\begin{aligned}
 & \log \left| \frac{y + \sqrt{x^2 + y^2}}{x} \right| - \log |x| = C \quad \left[\because v = \frac{y}{x} \right] \\
 \Rightarrow & \log \frac{y + \sqrt{x^2 + y^2}}{x} = C \\
 & \left[\because \log m - \log n = \log \left(\frac{m}{n} \right) \right] \\
 \Rightarrow & \frac{y + \sqrt{x^2 + y^2}}{x^2} = e^C \quad \left[\because \text{if } \log y = x, \right. \\
 & \quad \left. \text{then } y = e^x \right] \\
 \Rightarrow & y + \sqrt{x^2 + y^2} = x^2 \cdot e^C \\
 \therefore y + \sqrt{x^2 + y^2} = Ax^2 \quad [\text{where, } A = e^C] \quad (1)
 \end{aligned}$$

which is the required solution.

28. Solve the following differential equation

$$(y + 3x^2) \frac{dx}{dy} = x. \quad \text{All India 2011}$$

Given differential equation is

$$\begin{aligned}
 (y + 3x^2) \frac{dx}{dy} = x & \Rightarrow \frac{dy}{dx} = \frac{y}{x} + 3x \\
 \Rightarrow \frac{dy}{dx} - \frac{y}{x} & = 3x \quad \dots(i) \quad (1)
 \end{aligned}$$

which is a linear differential equation of the form

$$\frac{dy}{dx} + Py = Q \quad \dots(ii)$$

On comparing Eqs. (i) and (ii), we get

$$P = \frac{-1}{x} \text{ and } Q = 3x \quad (1)$$

$$\begin{aligned} \therefore \quad IF &= e^{\int -\frac{1}{x} dx} = e^{-\log|x|} = e^{\log x^{-1}} = x^{-1} \\ \Rightarrow \quad IF &= x^{-1} = \frac{1}{x} \end{aligned}$$

Now, solution of linear differential equation is given by

$$y \times IF = \int (Q \times IF) dx + C$$

$$\therefore \quad y \times \frac{1}{x} = \int 3x \times \frac{1}{x} dx \quad (1)$$

$$\Rightarrow \quad \frac{y}{x} = \int 3 dx \Rightarrow \frac{y}{x} = 3x + C$$

$$\Rightarrow \quad y = 3x^2 + Cx$$

which is the required solution. (1)

29. Solve the following differential equation

$$x dy - (y + 2x^2) dx = 0.$$

All India 2011

Given differential equation is

$$x dy - (y + 2x^2) dx = 0$$

$$\Rightarrow \frac{dy}{dx} = \frac{y + 2x^2}{x}$$

$$\Rightarrow \frac{dy}{dx} - \frac{y}{x} = 2x \quad \dots(i) \quad (1)$$

which is a linear differential equation of the form

$$\frac{dy}{dx} + Py = Q \quad \dots(ii)$$

On comparing Eqs. (i) and (ii), we get

$$P = \frac{-1}{x} \text{ and } Q = 2x \quad (1)$$

$$\therefore \text{IF} = e^{\int -\frac{1}{x} dx} = e^{-\log|x|} = x^{-1} = \frac{1}{x} \quad (1)$$

Now, solution of linear differential equation is given by

$$y \times \text{IF} = \int (Q \times \text{IF}) dx + C$$

$$\therefore \frac{y}{x} = \int \left(2x \times \frac{1}{x}\right) dx + C$$

$$\Rightarrow \frac{y}{x} = \int 2 dx + C \Rightarrow \frac{y}{x} = 2x + C$$

$$\Rightarrow y = 2x^2 + Cx$$

which is the required solution. (1)

30. Solve the following differential equation

$$x dy + (y - x^3) dx = 0.$$

All India 2011

Given differential equation is

$$x dy + (y - x^3) dx = 0$$

$$\Rightarrow \frac{dy}{dx} = \frac{x^3 - y}{x}$$

$$\Rightarrow \frac{dy}{dx} = x^2 - \frac{y}{x}$$

$$\Rightarrow \frac{dy}{dx} + \frac{y}{x} = x^2 \quad \dots(i) \quad (1)$$

which is a linear differential equation of the form

$$\frac{dy}{dx} + Py = Q \quad \dots(ii)$$

On comparing Eqs. (i) and (ii), we get

$$P = \frac{1}{x} \text{ and } Q = x^2 \quad (1)$$

$$\therefore \text{IF} = e^{\int \frac{1}{x} dx} = e^{\log|x|} = x \quad (1)$$

Solution of linear differential equation is given by

$$y \times \text{IF} = \int (Q \times \text{IF}) dx + C$$

$$\therefore y \times x = \int x^2 \times x dx + C$$

$$\Rightarrow yx = \int x^3 dx + C$$

$$\Rightarrow yx = \frac{x^4}{4} + C \Rightarrow y = \frac{x^3}{4} + \frac{C}{x}$$

which is the required solution. (1)

31. Solve the following differential equation

$$e^x \tan y dx + (1 - e^x) \sec^2 y dy = 0. \quad \text{Delhi 2011}$$

Given differential equation is

$$e^x \tan y \, dx + (1 - e^x) \sec^2 y \, dy = 0$$

$$\Rightarrow \frac{e^x}{e^x - 1} dx = \frac{\sec^2 y}{\tan y} dy \quad (1)$$

On integrating both sides, we get

$$\int \frac{e^x}{e^x - 1} dx = \int \frac{\sec^2 y}{\tan y} dy$$

On putting $e^x - 1 = t$ and $\tan y = z$

$$\Rightarrow e^x dx = dt \text{ and } \sec^2 y \, dy = dz$$

$$\therefore \int \frac{dt}{t} = \int \frac{dz}{z} \quad (1)$$

$$\Rightarrow \log |t| = \log |z| + \log C \quad [\because \int \frac{1}{x} dx = \log |x|]$$

$$\Rightarrow \log |e^x - 1| = \log |\tan y| + \log C$$

$$\Rightarrow \log |e^x - 1| = \log |C \cdot \tan y|$$

$$[\because \log m + \log n = \log mn]$$

$$\Rightarrow e^x - 1 = C \tan y \quad (1)$$

$$\Rightarrow \tan y = \frac{e^x - 1}{C} \Rightarrow y = \tan^{-1} \left(\frac{e^x - 1}{C} \right)$$

which is the required solution. (1)

32. Solve the following differential equation

$$(1 + y^2) (1 + \log x) \, dx + x \, dy = 0. \quad \text{Delhi 2011}$$

Given differential equation is

$$(1 + y^2)(1 + \log x) dx + x dy = 0$$

$$\Rightarrow \frac{1 + \log x}{x} dx = \frac{-dy}{1 + y^2} \quad (1)$$

On integrating both sides, we get

$$\int \frac{1 + \log x}{x} dx = - \int \frac{dy}{1 + y^2}$$

$$\Rightarrow \int \frac{1}{x} dx + \int \frac{\log x}{x} dx = - \int \frac{dy}{1 + y^2} \quad (1\frac{1}{2})$$

$$\Rightarrow \log |x| + \frac{(\log x)^2}{2} + C = - \tan^{-1} y$$

$$\left[\begin{array}{l} \text{for } \int \frac{\log x}{x} dx \Rightarrow \text{put } \log x = t \Rightarrow \frac{1}{x} dx = dt \\ \therefore \int t dt = \frac{t^2}{2} + C = \frac{(\log x)^2}{2} + C \end{array} \right]$$

$$\Rightarrow \tan^{-1} y = - \left[\log |x| + \frac{(\log x)^2}{2} + C \right]$$

$$\Rightarrow y = \tan \left[- \log |x| - \frac{(\log x)^2}{2} - C \right]$$

which is the required solution. (1\frac{1}{2})

33. Solve the following differential equation

$$\left[x \sin^2 \left(\frac{y}{x} \right) - y \right] dx + x dy = 0. \quad \text{Delhi 2011C}$$

Given differential equation is

$$\left[x \sin^2\left(\frac{y}{x}\right) - y \right] dx + x dy = 0$$

which is a homogeneous differential equation.

This equation can be written as

$$\left[x \sin^2\left(\frac{y}{x}\right) - y \right] dx = -x dy$$

$$\Rightarrow \frac{dy}{dx} = \frac{y - x \sin^2\left(\frac{y}{x}\right)}{x} \quad \dots(i)$$

On putting $y = vx \Rightarrow \frac{dy}{dx} = v + x \frac{dv}{dx}$ in

Eq. (i), we get (1)

$$v + x \frac{dv}{dx} = \frac{vx - x \sin^2\left(\frac{vx}{x}\right)}{x} = v - \sin^2 v$$

$$\Rightarrow x \frac{dv}{dx} = -\sin^2 v$$

$$\Rightarrow \frac{dv}{\sin^2 v} = -\frac{dx}{x} \quad (1)$$

On integrating both sides, we get

$$\int \frac{dv}{\sin^2 v} = -\int \frac{dx}{x}$$

$$\Rightarrow \int \operatorname{cosec}^2 v \, dv = -\int \frac{dx}{x} \left[\because \frac{1}{\sin^2 v} = \operatorname{cosec}^2 v \right]$$

$$\Rightarrow -\cot v = -\log x + C$$

$$\left[\because \int \operatorname{cosec}^2 v \, dv = -\cot v + C \right] \quad (1)$$

$$\Rightarrow -\cot \left(\frac{y}{x} \right) = -\log x + C \left[\because y = vx \therefore v = \frac{y}{x} \right]$$

$$\Rightarrow \cot \left(\frac{y}{x} \right) = \log x - C$$

$$\Rightarrow \frac{y}{x} = \cot^{-1}(\log x - C) \quad (1)$$

$$\Rightarrow y = x \cdot \cot^{-1}(\log x - C)$$

which is the required solution.

34. Solve the following differential equation

$$x \frac{dy}{dx} + y - x + xy \cot x = 0, \quad x \neq 0. \quad \text{Delhi 2011C}$$

Given differential equation is

$$x \frac{dy}{dx} + y - x + xy \cot x = 0, x \neq 0$$

Above equation can be written as

$$x \frac{dy}{dx} + y(1 + x \cot x) = x$$

On dividing both sides by x , we get

$$\frac{dy}{dx} + y \left(\frac{1 + x \cot x}{x} \right) = 1$$

$$\Rightarrow \frac{dy}{dx} + y \left(\frac{1}{x} + \cot x \right) = 1 \quad \dots(i) \quad (1)$$

which is a linear differential equation of the form

$$\frac{dy}{dx} + Py = Q \quad \dots(ii)$$

On comparing Eqs. (i) and (ii), we get

$$P = \frac{1}{x} + \cot x \text{ and } Q = 1$$

$$\therefore IF = e^{\int P dx} = e^{\int \left(\frac{1}{x} + \cot x \right) dx} = e^{\log|x| + \log \sin x}$$

$$\left[\because \int \frac{1}{x} dx = \log|x| \text{ and } \int \cot x dx = \log|\sin x| \right]$$

$$= e^{\log|x \sin x|}$$

$$[\because \log m + \log n = \log mn]$$

$$\Rightarrow IF = x \sin x \quad (1/2)$$

$$y \times IF = \int (Q \times IF) dx + C \quad (1/2)$$

$$y \times x \sin x = \int 1 \times x \sin x dx + C$$

$$\Rightarrow y x \sin x = \int \underset{I}{x} \underset{II}{\sin x} dx + C$$

$$\Rightarrow y x \sin x = x \int \sin x dx - \int \left(\frac{d}{dx}(x) \cdot \int \sin x dx \right) dx + C$$

$$[\text{using integration by parts in } \int x \sin x dx]$$

$$\Rightarrow y x \sin x = -x \cos x - \int 1(-\cos x) dx + C \quad (1)$$

$$\Rightarrow y x \sin x = -x \cos x + \int \cos x dx + C$$

$$\Rightarrow y x \sin x = -x \cos x + \sin x + C$$

On dividing both sides by $x \sin x$, we get

$$y = \frac{-x \cos x + \sin x + C}{x \sin x}$$

$$\Rightarrow y = -\cot x + \frac{1}{x} + \frac{C}{x \sin x}$$

which is the required solution. (1)

35. Show that the following differential equation is homogeneous and then solve it.

$$y dx + x \log \left(\frac{y}{x} \right) dy - 2x dy = 0$$

HOTS; All India 2011C

Let the value of $\frac{dy}{dx}$ be $f(x, y)$. Now, put $x = \lambda x$ and $y = \lambda y$ and verify whether $f(\lambda x, \lambda y) = \lambda^n f(x, y)$ $n \in \mathbb{Z}$. If above equation is satisfied, then given equation is said to be homogeneous equation. Then, we use the substitution $y = vx$ to solve the equation.

Given differential equation is

$$y dx + x \log \left(\frac{y}{x} \right) dy - 2x dy = 0$$

$$\Rightarrow y dx = \left[2x - x \log \left(\frac{y}{x} \right) \right] dy$$

$$\Rightarrow \frac{dy}{dx} = \frac{y}{2x - x \log \left(\frac{y}{x} \right)} \quad \dots(i) \quad (1/2)$$

$$\text{Now, let } f(x, y) = \frac{y}{2x - x \log \left(\frac{y}{x} \right)}$$

On replace x by λx and y by λy both sides, we get

$$f(\lambda x, \lambda y) = \frac{\lambda y}{2\lambda x - \lambda x \log \left(\frac{\lambda y}{\lambda x} \right)}$$

$$= \frac{\lambda y}{\lambda \left[2x - x \log \left(\frac{y}{x} \right) \right]}$$

$$\Rightarrow f(\lambda x, \lambda y) = \lambda^0 \frac{y}{2x - x \log \left(\frac{y}{x} \right)} = \lambda^0 f(x, y)$$

So, given differential equation is homogeneous. (1/2)

dv

dv

On putting $y = vx \Rightarrow \frac{dy}{dx} = v + x \frac{dv}{dx}$

in Eq. (i), we get

$$v + x \frac{dv}{dx} = \frac{vx}{2x - x \log \left(\frac{vx}{x} \right)} = \frac{v}{2 - \log v}$$

$$\Rightarrow x \frac{dv}{dx} = \frac{v}{2 - \log v} - v = \frac{v - 2v + v \log v}{2 - \log v}$$

$$\Rightarrow x \frac{dv}{dx} = \frac{-v + v \log v}{2 - \log v}$$

$$\Rightarrow \frac{2 - \log v}{v \log v - v} dv = \frac{dx}{x} \quad (1)$$

On integrating both sides, we get

$$\int \frac{2 - \log v}{v(\log v - 1)} dv = \int \frac{dx}{x}$$

On putting $\log v = t \Rightarrow \frac{1}{v} dv = dt$

Then, $\int \frac{2 - t}{t - 1} dt = \log |x| + C$

$$\Rightarrow \int \left(\frac{1}{t - 1} - 1 \right) dt = \log |x| + C \quad (1)$$

$$\left[\begin{array}{l} \because t - 1 \overline{) 2 - t} \begin{array}{r} -1 \\ 1 - t \\ \hline 1 \end{array} \\ \text{and use } \int \left(\frac{R}{D} + Q \right) dt \end{array} \right]$$

$$\Rightarrow \log |t - 1| - t = \log |x| + C$$

$$\Rightarrow \log |\log v - 1| - \log v = \log |x| + C$$

$$\Rightarrow \log \left| \frac{\log v - 1}{v} \right| = \log |x| + C$$

$$\left[\because \log m - \log n = \log \left(\frac{m}{n} \right) \right]$$

$$\Rightarrow \log \left| \frac{\log v - 1}{v} \right| - \log |x| = C$$

$$\Rightarrow \log \left| \frac{\log v - 1}{vx} \right| = C$$

$$\therefore \log \left| \frac{\log \frac{y}{x} - 1}{\frac{y}{x}} \right| = C \quad \left[\because y = vx \Rightarrow v = \frac{y}{x} \right]$$

which is the required solution. (1)

36. Solve the following differential equation

$$\left(x \cos \frac{y}{x} + y \sin \frac{y}{x} \right) y - \left(y \sin \frac{y}{x} - x \cos \frac{y}{x} \right) x \frac{dy}{dx} = 0.$$

All India 2010C



Firstly, convert the given differential equation in homogeneous and then put $y = vx$.

$$\Rightarrow \frac{dy}{dx} = v + x \frac{dv}{dx}$$

Further, separate the variables and integrate it.

Given differential equation is

$$\left(x \cos \frac{y}{x} + y \sin \frac{y}{x} \right) \cdot y$$

$$- \left(y \sin \frac{y}{x} - x \cos \frac{y}{x} \right) \cdot x \frac{dy}{dx} = 0$$

which is a homogeneous differential equation.

It can be written as

$$\left(x \cos \frac{y}{x} + y \sin \frac{y}{x} \right) \cdot y$$

$$= \left(y \sin \frac{y}{x} - x \cos \frac{y}{x} \right) \cdot x \frac{dy}{dx}$$

$$\Rightarrow \frac{dy}{dx} = \frac{\left[x \cos \left(\frac{y}{x} \right) + y \sin \left(\frac{y}{x} \right) \right] \cdot y}{\left(y \sin \frac{y}{x} - x \cos \frac{y}{x} \right) \cdot x} \quad \dots(i)$$

On putting $y = vx$

$$\Rightarrow \frac{dy}{dx} = v + x \frac{dv}{dx} \text{ in Eq. (i), we get} \quad (1)$$

$$v + x \frac{dv}{dx} = \frac{(x \cos v + vx \sin v) \cdot vx}{(vx \sin v - x \cos v) \cdot x}$$

$$\Rightarrow v + x \frac{dv}{dx} = \frac{v \cos v + v^2 \sin v}{v \sin v - \cos v}$$

$$\Rightarrow x \frac{dv}{dx} = \frac{v \cos v + v^2 \sin v}{v \sin v - \cos v} - v$$

$$\Rightarrow x \frac{dv}{dx} = \frac{v \cos v + v^2 \sin v - v^2 \sin v + v \cos v}{v \sin v - \cos v}$$

(1)

$$\Rightarrow x \frac{dv}{dx} = \frac{2v \cos v}{v \sin v - \cos v}$$

$$\Rightarrow \frac{v \sin v - \cos v}{v \cos v} dv = 2 \frac{dx}{x}$$

On integrating both sides, we get

$$\int \frac{v \sin v - \cos v}{v \cos v} dv = 2 \int \frac{dx}{x}$$

$$\Rightarrow \int \left(\frac{v \sin v}{v \cos v} - \frac{\cos v}{v \cos v} \right) dv = 2 \int \frac{dx}{x}$$

$$\Rightarrow \int \left(\tan v - \frac{1}{v} \right) dv = 2 \int \frac{dx}{x}$$

$$\Rightarrow \log |\sec v| - \log |v| = 2 \log |x| + C \quad (1)$$

$$\left[\because \int \tan v dv = \log |\sec v| \text{ and } \int \frac{1}{x} dx = \log |x| \right]$$

$$\Rightarrow \log |\sec v| - \log |v| - 2 \log |x| = C$$

$$\Rightarrow \log |\sec v| - [\log |v| + \log |x|^2] = C$$

$$[\because \log m^n = n \log m]$$

$$\Rightarrow \log |\sec v| - \log |vx^2| = C$$

$$[\because \log m + \log n = \log mn]$$

$$\Rightarrow \log \left| \frac{\sec v}{vx^2} \right| = C$$

$$\left[\because \log m - \log n = \log \left(\frac{m}{n} \right) \right]$$

$$\Rightarrow \log \left| \frac{\sec \frac{y}{x}}{\frac{y}{x} \cdot x^2} \right| = C \quad \left[\begin{array}{l} \because y = vx \\ \therefore v = \frac{y}{x} \end{array} \right]$$

$$\Rightarrow \log \left| \frac{\sec \frac{y}{x}}{xy} \right| = C$$

which is the required solution

(1)

which is the required solution.

37. Solve the following differential equation

$$xy \log \left(\frac{y}{x} \right) dx + \left[y^2 - x^2 \log \left(\frac{y}{x} \right) \right] dy = 0. \quad \text{Delhi 2010C}$$

Given differential equation is

$$xy \log \left(\frac{y}{x} \right) dx + \left[y^2 - x^2 \log \left(\frac{y}{x} \right) \right] dy = 0$$

which is a homogeneous differential equation. This equation can be written as

$$xy \log \left(\frac{y}{x} \right) dx = \left[x^2 \log \left(\frac{y}{x} \right) - y^2 \right] dy$$

$$\Rightarrow \frac{dy}{dx} = \frac{xy \log \left(\frac{y}{x} \right)}{x^2 \log \left(\frac{y}{x} \right) - y^2} \quad \dots(i)$$

$$\text{Now, put } y = vx \Rightarrow \frac{dy}{dx} = v + x \frac{dv}{dx} \quad (1)$$

in Eq. (i), we get

$$v + x \frac{dv}{dx} = \frac{vx^2 \log \left(\frac{vx}{x} \right)}{x^2 \log \left(\frac{vx}{x} \right) - v^2 x^2} = \frac{v \log v}{\log v - v^2}$$

$$\Rightarrow x \frac{dv}{dx} = \frac{v \log v}{\log v - v^2} - v$$

$$\Rightarrow x \frac{dv}{dx} = \frac{v \log v - v \log v + v^3}{\log v - v^2} = \frac{v^3}{\log v - v^2}$$

$$\Rightarrow \frac{\log v - v^2}{v^3} dv = \frac{dx}{x} \quad (1)$$

On integrating both sides, we get

$$\int \frac{\log v - v^2}{v^3} dv = \int \frac{dx}{x}$$

$\int \frac{\log v}{v^3} dv - \int \frac{v^2}{v^3} dv = \int \frac{dx}{x}$

$$\Rightarrow \int \frac{-\frac{1}{v}}{v^3} dv - \int \frac{1}{v} dv = \int \frac{-1}{x}$$

$$\Rightarrow \int \frac{v^{-3}}{v} \log v dv - \log |v| = \log |x| + C$$

Using integration by parts, we get

$$\log v \int v^{-3} dv - \int \left[\frac{d}{dv} (\log v) \cdot \int v^{-3} dv \right] dv$$

$$= \log |v| + \log |x| + C$$

$$\Rightarrow \frac{v^{-2}}{-2} \log v - \int \frac{1}{v} \frac{v^{-2}}{(-2)} dv = \log |v| + \log |x| + C$$

$$\Rightarrow \frac{-1}{2v^2} \log v + \frac{1}{2} \int v^{-3} dv = \log |v| + \log |x| + C$$

$$\Rightarrow \frac{-1}{2v^2} \log v + \frac{1}{2} \cdot \frac{v^{-2}}{(-2)} = \log |v| + \log |x| + C$$

$$\left[\because \int x^n dx = \frac{x^{n+1}}{n+1} + C \right]$$

$$\Rightarrow \frac{-1}{2v^2} \log v - \frac{1}{4v^2} = \log |vx| + C \quad (1)$$

$$[\because \log m + \log n = \log mn]$$

$$\Rightarrow \frac{-1}{2} \cdot \frac{x^2}{y^2} \log \left(\frac{y}{x} \right) - \frac{1}{4} \cdot \frac{x^2}{y^2} = \log \left| \frac{y}{x} \cdot x \right| + C$$

$$\left[\because y = vx \Rightarrow v = \frac{y}{x} \right]$$

$$\Rightarrow \frac{-x^2}{2y^2} \log \left(\frac{y}{x} \right) - \frac{x^2}{4y^2} = \log |y| + C$$

$$\Rightarrow \frac{-x^2}{y^2} \left[\frac{\log \left(\frac{y}{x} \right)}{2} + \frac{1}{4} \right] = \log |y| + C$$

$$\Rightarrow \frac{x^2}{4y^2} \left[2 \log \left(\frac{y}{x} \right) + 1 \right] + \log |y| = -C$$

$$\Rightarrow x^2 \left[2 \log \left(\frac{y}{x} \right) + 1 \right] + 4y^2 \log |y| = 4y^2 k$$

[where, $k = -C$] (1)

which is the required solution.

38. Solve the following differential equation

$$(x^2 + 1) \frac{dy}{dx} + 2xy = \sqrt{x^2 + 4}. \quad \text{All India 2010, 2008}$$

Given differential equation is

$$(x^2 + 1) \frac{dy}{dx} + 2xy = \sqrt{x^2 + 4}$$

On dividing both sides by $(x^2 + 1)$, we get

$$\frac{dy}{dx} + \frac{2xy}{x^2 + 1} = \frac{\sqrt{x^2 + 4}}{x^2 + 1} \quad \dots(i)$$

which is a linear differential equation of the form $\frac{dy}{dx} + Py = Q$...(ii)

On comparing Eqs. (i) and (ii), we get

$$P = \frac{2x}{x^2 + 1} \quad \text{and} \quad Q = \frac{\sqrt{x^2 + 4}}{x^2 + 1}$$

$$\therefore \quad \text{IF} = e^{\int \frac{2x}{x^2 + 1} dx} = e^{\log|x^2 + 1|}$$

$$\Rightarrow \quad \text{IF} = x^2 + 1 \quad [\because e^{\log x} = x] \quad (1)$$

$$\left[\because \int \frac{2x}{x^2 + 1} dx \Rightarrow \text{put } x^2 + 1 = t \Rightarrow 2x dx = dt \right.$$

$$\left. \therefore \int \frac{dt}{t} = \log|t| = \log|x^2 + 1| \right]$$

Now, solution of this equation is given by

$$y \times \text{IF} = \int (Q \times \text{IF}) dx + C \quad (1)$$

$$\therefore \quad y(x^2 + 1) = \int (x^2 + 1) \cdot \frac{\sqrt{x^2 + 4}}{x^2 + 1} dx \quad (1)$$

$$\Rightarrow y(x^2 + 1) = \int \sqrt{x^2 + 4} dx$$

$$\Rightarrow y(x^2 + 1) = \int \sqrt{x^2 + (2)^2} dx$$

Now, we know that

$$\int \sqrt{x^2 + a^2} dx = \frac{x}{2} \sqrt{x^2 + a^2} + \frac{a^2}{2} \log |x + \sqrt{x^2 + a^2}| + C$$

$$\therefore y(x^2 + 1) = \frac{x}{2} \sqrt{x^2 + 4} + \frac{4}{2} \log |x + \sqrt{x^2 + 4}| + C$$

$$\Rightarrow y(x^2 + 1) = \frac{x}{2} \sqrt{x^2 + 4} + 2 \log |x + \sqrt{x^2 + 4}| + C$$

which is the required solution. (1)

39. Solve the following differential equation

$$(x^3 + x^2 + x + 1) \frac{dy}{dx} = 2x^2 + x.$$

HOTS; All India 2010



Firstly, divide given equation by $x^3 + x^2 + x + 1$, then it becomes a variable separable type differential equation and then solve it.

Given differential equation is

$$(x^3 + x^2 + x + 1) \frac{dy}{dx} = 2x^2 + x$$

$$\Rightarrow \frac{dy}{dx} = \frac{2x^2 + x}{x^3 + x^2 + x + 1}$$

It is a variable separable type differential equation.

$$dy = \frac{2x^2 + x}{x^3 + x^2 + x + 1} dx$$

On integrating both sides, we get

$$\int dy = \int \frac{2x^2 + x}{x^3 + x^2 + x + 1} dx$$

$$\Rightarrow y = \int \frac{2x^2 + x}{x^2(x+1) + 1(x+1)} dx$$

$$= \int \frac{2x^2 + x}{(x+1)(x^2+1)} dx$$

$$y = I \quad \dots(i) \quad (1)$$

where, $I = \int \frac{2x^2 + x}{(x+1)(x^2+1)} dx$

Using partial fractions, we get

$$\frac{2x^2 + x}{(x+1)(x^2+1)} = \frac{A}{x+1} + \frac{Bx+C}{x^2+1} \quad \dots(ii)$$

$$\Rightarrow \frac{2x^2 + x}{(x+1)(x^2+1)} = \frac{A(x^2+1) + (Bx+C)(x+1)}{(x+1)(x^2+1)}$$

$$\Rightarrow 2x^2 + x = A(x^2+1) + (Bx+C)(x+1)$$

Now, comparing coefficients of x^2 , x and constant term from both sides we get

constant term from both sides, we get

$$A + B = 2 \quad \dots(\text{iii})$$

$$B + C = 1 \quad \dots(\text{iv})$$

and $A + C = 0 \quad \dots(\text{v})$

On subtracting Eq. (iv) from Eq. (iii), we get

$$A - C = 1 \quad \dots(\text{vi})$$

On adding Eqs. (v) and (vi), we get

$$2A = 1 \Rightarrow A = \frac{1}{2}$$

On putting $A = \frac{1}{2}$ in Eq. (iii), we get

$$\frac{1}{2} + B = 2 \Rightarrow B = 2 - \frac{1}{2} = \frac{3}{2}$$

On putting $B = \frac{3}{2}$ in Eq. (iv), we get

$$\frac{3}{2} + C = 1 \Rightarrow C = 1 - \frac{3}{2}$$

$$\Rightarrow C = \frac{-1}{2} \quad (1)$$

On substituting the values of A , B and C in Eq. (ii), we get

$$\frac{2x^2 + x}{(x+1)(x^2+1)} = \frac{1/2}{x+1} + \frac{\frac{3}{2}x - \frac{1}{2}}{x^2+1}$$

On integrating both sides, we get

$$\begin{aligned} I &= \int \frac{2x^2 + x}{(x+1)(x^2+1)} dx = \frac{1}{2} \int \frac{dx}{x+1} \\ &\quad + \frac{3}{2} \int \frac{x}{x^2+1} dx - \frac{1}{2} \int \frac{dx}{x^2+1} \end{aligned}$$

$$\begin{aligned} \Rightarrow I &= \frac{1}{2} \log|x+1| + \frac{3}{4} \log|x^2+1| \\ &\quad - \frac{1}{2} \tan^{-1} x + C \quad (1) \end{aligned}$$

$$\left[\because \int \frac{x}{x^2+1} dx \Rightarrow \text{put } x^2+1=t \Rightarrow 2x dx = dt \right]$$

$$\Rightarrow x dx = \frac{dt}{2}, \text{ then } \int \frac{x}{x^2 + 1} dx = \frac{1}{2} \int \frac{dt}{t} = \frac{1}{2} \log t = \frac{1}{2} \log |x^2 + 1| + C$$

On putting above value of I in Eq. (i), we get

$$y = \frac{1}{2} \log |x + 1| + \frac{3}{4} \log |x^2 + 1| - \frac{1}{2} \tan^{-1} x + C$$

which is the required solution. **(1)**

40. Solve the following differential equation

$$\sqrt{1 + x^2 + y^2 + x^2 y^2} + xy \frac{dy}{dx} = 0. \text{ All India 2010}$$

Given differential equation is

$$\sqrt{1 + x^2 + y^2 + x^2 y^2} + xy \frac{dy}{dx} = 0$$

$$\Rightarrow \sqrt{(1 + x^2) + y^2(1 + x^2)} = -xy \frac{dy}{dx}$$

$$\Rightarrow \sqrt{(1 + x^2)(1 + y^2)} = -xy \frac{dy}{dx}$$

$$\Rightarrow \sqrt{1 + x^2} \cdot \sqrt{1 + y^2} = -xy \frac{dy}{dx}$$

$$\Rightarrow \frac{y}{\sqrt{1 + y^2}} dy = -\frac{\sqrt{1 + x^2}}{x} dx \quad (1)$$

On integrating both sides, we get

$$\int \frac{y}{\sqrt{1 + y^2}} dy = - \int \frac{\sqrt{1 + x^2}}{x^2} \cdot x dx$$

On putting $1 + y^2 = t$ and $1 + x^2 = u^2$

$$\Rightarrow 2y dy = dt \text{ and } 2x dx = 2u du$$

$$\Rightarrow y dy = \frac{dt}{2} \text{ and } x dx = u du \quad (1)$$

$$\therefore \frac{1}{2} \int \frac{dt}{\sqrt{t}} = - \int \frac{u}{u^2 - 1} \cdot u du$$

$$\Rightarrow \frac{1}{2} \int t^{-1/2} dt = - \int \frac{u^2}{u^2 - 1} du$$

$$\begin{aligned}
 & \Rightarrow \frac{1}{2} t^{1/2} = - \int \frac{(u^2 - 1 + 1)}{u^2 - 1} du \quad (1) \\
 & \Rightarrow t^{1/2} = - \int \frac{u^2 - 1}{u^2 - 1} du - \int \frac{1}{u^2 - 1} du \\
 & \Rightarrow \sqrt{1 + y^2} = - \int du - \int \frac{1}{u^2 - (1)^2} du \\
 & \quad [\because 1 + y^2 = t] \\
 & \Rightarrow \sqrt{1 + y^2} = -u - \frac{1}{2} \log \left| \frac{u-1}{u+1} \right| + C \\
 & \quad \left[\because \int \frac{dx}{x^2 - a^2} = \frac{1}{2a} \log \left| \frac{x-a}{x+a} \right| + C \right] \\
 & \Rightarrow \sqrt{1 + y^2} = -\sqrt{1 + x^2} - \frac{1}{2} \log \left| \frac{\sqrt{1 + x^2} - 1}{\sqrt{1 + x^2} + 1} \right| + C
 \end{aligned}$$

which is the required solution. (1)

41. Find the particular solution of the differential equation satisfying the given condition

$$x^2 dy + (xy + y^2) dx = 0, \text{ when } y(1) = 1.$$

Delhi 2010

Given differential equation is

$$x^2 dy + (xy + y^2) dx = 0$$

Since, degree of each term is same, so the above equation is a homogeneous equation.

This equation can be written as

$$x^2 dy = -(xy + y^2) dx$$

$$\Rightarrow \frac{dy}{dx} = \frac{-(xy + y^2)}{x^2} \quad \dots(i)$$

On putting $y = vx$

$$\Rightarrow \frac{dy}{dx} = v + x \frac{dv}{dx} \quad (1)$$

in Eq (i), we get

$$v + x \frac{dv}{dx} = \frac{-(vx^2 + v^2 x^2)}{x^2} = -(v + v^2)$$

$$\begin{aligned}
 \Rightarrow x \frac{dv}{dx} &= -v - v^2 - v \\
 \Rightarrow x \frac{dv}{dx} &= -v^2 - 2v \\
 \Rightarrow \frac{dv}{v^2 + 2v} &= \frac{-dx}{x} \quad (1)
 \end{aligned}$$

On integrating both sides, we get

$$\begin{aligned}
 \int \frac{dv}{v^2 + 2v} &= - \int \frac{dx}{x} \\
 \Rightarrow \int \frac{dv}{v^2 + 2v + 1 - 1} &= - \int \frac{dx}{x} \\
 \Rightarrow \int \frac{dv}{(v+1)^2 - (1)^2} &= - \int \frac{dx}{x} \\
 \Rightarrow \frac{1}{2} \log \left| \frac{v+1-1}{v+1+1} \right| &= - \log |x| + C \\
 \left[\because \int \frac{dx}{x^2 - a^2} = \frac{1}{2a} \log \left| \frac{x-a}{x+a} \right| + C \right] \\
 \Rightarrow \frac{1}{2} \log \left| \frac{v}{v+2} \right| &= - \log |x| + C \\
 \Rightarrow \frac{1}{2} \log \left| \frac{\frac{y}{x}}{\frac{y}{x} + 2} \right| &= - \log |x| + C, \quad \left[\begin{array}{l} \because y = vx \\ \therefore v = \frac{y}{x} \end{array} \right] \\
 \Rightarrow \frac{1}{2} \log \left| \frac{y}{y+2x} \right| &= - \log |x| + C \quad \dots(ii)
 \end{aligned}$$

Also, given that $y = 0$ at $x = 1, y = 1$.

On putting $x = y = 1$ in Eq. (ii), we get

$$\therefore \frac{1}{2} \log \left| \frac{1}{1+2} \right| = - \log 1 + C$$

$$\Rightarrow \frac{1}{2} \log \left| \frac{1}{3} \right| = -\log 1 + C$$

$$\Rightarrow C = \frac{1}{2} \log \frac{1}{3} \quad [\because \log 1 = 0] \quad (1)$$

On putting the value of C in Eq. (ii), we get

$$\frac{1}{2} \log \left| \frac{y}{y+2x} \right| = -\log |x| + \frac{1}{2} \log \frac{1}{3}$$

$$\Rightarrow \log \left| \frac{y}{y+2x} \right| = -2 \log |x| + \log \frac{1}{3}$$

$$\Rightarrow \log \frac{y}{y+2x} = \log x^{-2} + \log \frac{1}{3}$$

$$[\because n \log m = \log m^n]$$

$$\Rightarrow \log \frac{y}{y+2x} = \log \frac{1}{x^2} + \log \frac{1}{3}$$

$$\Rightarrow \log \left(\frac{y}{y+2x} \right) = \log \frac{1}{3x^2}$$

$$[\because \log m + \log n = \log mn]$$

$$\Rightarrow \frac{y}{y+2x} = \frac{1}{3x^2}$$

$$\Rightarrow y \cdot 3x^2 = y + 2x$$

$$\Rightarrow y(1 - 3x^2) = -2x$$

$$\therefore y = \frac{2x}{3x^2 - 1}$$

which is the required particular solution. **(1)**

42. Find the particular solution of the differential equation satisfying the given condition

$$\frac{dy}{dx} = y \tan x, \text{ given that } y = 1, \text{ when } x = 0.$$

Delhi 2010

Given differential equation is

$$\frac{dy}{dx} = y \tan x$$

It can be written as $\frac{dy}{y} = \tan x \, dx$ (1)

On integrating both sides, we get

$$\int \frac{dy}{y} = \int \tan x \, dx$$

$$\Rightarrow \log|y| = \log|\sec x| + C \quad \dots(i) \quad (1)$$

$$\left[\because \int \frac{1}{y} dy = \log|y| \text{ and } \int \tan x \, dx = \log|\sec x| \right]$$

Also, given that $y = 1$, when $x = 0$.

On putting $x = 0$ and $y = 1$ in Eq.(i), we get

$$\log 1 = \log(\sec 0^\circ) + C$$

$$\Rightarrow 0 = \log 1 + C \quad [\because \sec 0^\circ = 1] \quad (1)$$

$$\Rightarrow C = 0 \quad [\because \log 1 = 0]$$

On putting $C = 0$ in Eq. (i), we get the required particular solution as

$$\log|y| = \log|\sec x|$$

$$\therefore y = \sec x \quad (1)$$

which is the required solution.

43. Solve the following differential equation

$$\cos^2 x \frac{dy}{dx} + y = \tan x.$$

All India 2009; Delhi 2008, 2011, 2008C

Given differential equation is

$$\cos^2 x \frac{dy}{dx} + y = \tan x$$

On dividing both sides by $\cos^2 x$, we get

$$\frac{dy}{dx} + \frac{y}{\cos^2 x} = \frac{\tan x}{\cos^2 x}$$

$$\Rightarrow \frac{dy}{dx} + y \cdot \sec^2 x = \tan x \cdot \sec^2 x \quad \dots(i)$$

$$\left[\because \frac{1}{\cos^2 x} = \sec^2 x \right]$$

which is the linear differential equation of the form

$$\frac{dy}{dx} + Py = Q \quad \dots(ii)$$

On comparing Eqs. (i) and (ii), we get

$$P = \sec^2 x \text{ and } Q = \tan x \cdot \sec^2 x \quad (1)$$

$$\therefore \text{IF} = e^{\int \sec^2 x \, dx} = e^{\tan x}$$

$$[\because \int \sec^2 x \, dx = \tan x + C] \quad (1)$$

Now, solution of linear differential equation is given by

$$y \times \text{IF} = \int (Q \times \text{IF}) \, dx + C$$

$$\therefore y \times e^{\tan x} = \int \tan x \cdot \sec^2 x \cdot e^{\tan x} \, dx \quad \dots(iii)$$

On putting $\tan x = t$

$\Rightarrow \sec^2 x \, dx = dt$ in Eq. (iii), we get

$$\therefore ye^{\tan x} = \int t e^t \, dt \quad (1)$$

$$\Rightarrow ye^{\tan x} = t \int e^t \, dt - \int \left[\frac{d}{dt}(t) \int e^t \, dt \right] dt$$

[using integration by parts in $\int te^t \, dt$]

$$\Rightarrow ye^{\tan x} = te^t - \int 1 \times e^t \, dt$$

$$\Rightarrow ye^{\tan x} = te^t - e^t + C$$

$$\therefore ye^{\tan x} = \tan x \cdot e^{\tan x} - e^{\tan x} + C [\because \tan x = t]$$

On dividing both sides by $e^{\tan x}$, we get

$$y = \tan x - 1 + Ce^{-\tan x}$$

which is the required solution. (1)

44. Solve the following differential equation

$$\sec x \frac{dy}{dx} - y = \sin x.$$

All India 2009C

Given differential equation is

$$\sec x \frac{dy}{dx} - y = \sin x$$

On dividing both sides by $\sec x$, we get

$$\frac{dy}{dx} - \frac{y}{\sec x} = \frac{\sin x}{\sec x}$$

$$\Rightarrow \frac{dy}{dx} - y \cos x = \sin x \cos x \quad \dots(i)$$

which is a linear differential equation of the

$$\text{form} \quad \frac{dy}{dx} + Py = Q \quad \dots(ii)$$

On comparing Eqs. (i) and (ii), we get

$$P = -\cos x \text{ and } Q = \sin x \cos x \quad (1)$$

$$\therefore \text{IF} = e^{\int -\cos x \, dx} = e^{-\sin x}$$

$$[\because \int \cos x \, dx = \sin x + C] \quad (1)$$

Now, solution of above equation is given by

$$y \times \text{IF} = \int (Q \times \text{IF}) \, dx + C$$

$$\therefore ye^{-\sin x} = \int \sin x \cos x e^{-\sin x} \, dx$$

$$\text{On putting } \sin x = t \Rightarrow \cos x \, dx = dt$$

$$\therefore ye^{-\sin x} = \int t e^{-t} \, dt \quad (1)$$

$$\Rightarrow ye^{-\sin x} = t \int e^{-t} \, dt - \int \left[\frac{d}{dt}(t) \int e^{-t} \, dt \right] dt$$

[using integration by parts]

$$\begin{aligned} \Rightarrow ye^{-\sin x} &= -te^{-t} - \int 1 \times (-e^{-t}) \, dt \\ &= -te^{-t} + \int e^{-t} \, dt \end{aligned}$$

$$\Rightarrow ye^{-\sin x} = -te^{-t} - e^{-t} + C$$

$$\Rightarrow ye^{-\sin x} = -\sin x e^{-\sin x} - e^{-\sin x} + C$$

$$[\because \sin x = t]$$

$$y = -\sin x - 1 + C e^{\sin x} \quad (1)$$

which is the required solution.

45. Solve the following differential equation

$$(x \log x) \frac{dy}{dx} + y = 2 \log x. \quad \text{Delhi 2009, 2009C}$$

Given differential equation is

$$(x \log x) \frac{dy}{dx} + y = 2 \log x$$

On dividing both sides by $x \log x$, we get

$$\frac{dy}{dx} + \frac{y}{x \log x} = \frac{2}{x} \quad \dots(i)$$

which is a linear differential equation of the form

$$\frac{dy}{dx} + Py = Q \quad \dots(ii)$$

On comparing Eqs. (i) and (ii), we get

$$P = \frac{1}{x \log x} \text{ and } Q = \frac{2}{x} \quad (1)$$

$$\begin{aligned} \therefore \text{IF} &= e^{\int \frac{1}{x \log x} dx} = e^{\log \log x} \\ &= \log x \quad [\because e^{\log x} = x] \quad (1) \end{aligned}$$

$$\begin{aligned} \left[\because \int \frac{1}{x \log x} dx \Rightarrow \text{put } \log x = t \Rightarrow \frac{1}{x} dx = dt \right. \\ \left. \therefore \int \frac{1}{x \log x} dx = \int \frac{dt}{t} = \log |t| = \log |\log x| \right] \end{aligned}$$

Now, solution of above equation is given by

$$y \times \text{IF} = \int (Q \times \text{IF}) dx + C$$

$$\therefore y \times \log x = \int \frac{2}{x} \log x dx \quad (1)$$

$$\begin{aligned} \Rightarrow y \log x &= \log x \int \frac{2}{x} dx \\ &\quad - \int \left[\frac{d}{dx} (\log x) \int \frac{2}{x} dx \right] dx \end{aligned}$$

[using integration by parts]

$$\Rightarrow y \log x = \log x \cdot 2 \log x - \int \frac{1}{x} \cdot 2 \log x \, dx$$

$$\left[\because \int \frac{1}{x} \, dx = \log |x| + C \right]$$

$$\Rightarrow y \log x = 2 (\log x)^2 - 2 \int \frac{\log x}{x} \, dx$$

$$\Rightarrow y \log x = 2 (\log x)^2 - \frac{2(\log x)^2}{2} + C$$

$$\left[\text{in } \int \frac{\log x}{x} \, dx, \text{ put } \log x = t \Rightarrow \frac{1}{x} \, dx = dt \right]$$

$$\therefore \int t \, dt = \frac{t^2}{2} = \frac{(\log x)^2}{2} + C$$

$$y = 2 (\log x) - (\log x) + \frac{C}{\log x}$$

[dividing both sides by $\log x$] (1)

which is the required solution.

46. Solve the following differential equation

$$x \frac{dy}{dx} = y - x \tan \left(\frac{y}{x} \right) \quad \text{All India 2009}$$

Given differential equation is

$$x \frac{dy}{dx} = y - x \tan \left(\frac{y}{x} \right)$$

$$\Rightarrow \frac{dy}{dx} = \frac{y - x \tan \left(\frac{y}{x} \right)}{x} \quad \dots (i)$$

which is a homogeneous differential equation.

On putting $y = vx$

$$\Rightarrow \frac{dy}{dx} = v + x \frac{dv}{dx} \quad (1)$$

in Eq. (i), we get

$$v + x \frac{dv}{dx} = \frac{vx - x \tan v}{x} = v - \tan v$$

$$\Rightarrow x \frac{dv}{dx} = -\tan v$$

$$\Rightarrow \frac{dv}{\tan v} = -\frac{dx}{x} \quad (1)$$

$$\Rightarrow \cot v \, dv = -\frac{dx}{x} \quad \left[\because \frac{1}{\tan v} = \cot v \right] (1)$$

On integrating both sides, we get

$$\int \cot v \, dv = -\int \frac{dx}{x}$$

$$\Rightarrow \log |\sin v| = -\log |x| + C$$

$$[\because \int \cot v \, dv = \log |\sin v| + C]$$

$$\Rightarrow \log |\sin v| + \log |x| = C$$

$$\Rightarrow \log |x \sin v| = C$$

$$[\because \log m + \log n = \log mn]$$

$$\therefore \log \left| x \sin \frac{y}{x} \right| = C \quad \left[\because v = \frac{y}{x} \right] (1)$$

which is the required solution.

47. Solve the following differential equation

$$(1 + x^2) \frac{dy}{dx} + y = \tan^{-1} x.$$

Delhi 2009

The given differential equation is

$$(1+x^2) \frac{dy}{dx} + y = \tan^{-1} x$$

On dividing both sides by $(1+x^2)$, we get

$$\frac{dy}{dx} + \frac{y}{1+x^2} = \frac{\tan^{-1} x}{1+x^2} \quad \dots(i)$$

which is a linear differential equation of the form

$$\frac{dy}{dx} + Py = Q \quad \dots(ii)$$

On comparing Eqs. (i) and (ii), we get

$$P = \frac{1}{1+x^2} \text{ and } Q = \frac{\tan^{-1} x}{1+x^2} \quad (1)$$

$$\therefore \text{IF} = e^{\int \frac{1}{1+x^2} dx} = e^{\tan^{-1} x}$$

$$\left[\because \int \frac{1}{1+x^2} dx = \tan^{-1} x + C \right] \quad (1)$$

Now, solution of above equation is given by

$$y \times IF = \int (Q \times IF) dx + C$$

$$\therefore y \times e^{\tan^{-1} x} = \int \frac{\tan^{-1} x}{1+x^2} \cdot e^{\tan^{-1} x} dx \quad \dots(iii)$$

On putting $\tan^{-1} x = t$

$$\Rightarrow \frac{1}{1+x^2} dx = dt \quad (1)$$

in Eq. (iii), we get

$$ye^{\tan^{-1} x} = \int t e^t dt$$

$$\Rightarrow ye^{\tan^{-1} x} = t \int e^t dt - \int \left[\frac{d}{dt}(t) \int e^t dt \right] dt$$

[using integration by parts]

$$\Rightarrow ye^{\tan^{-1} x} = te^t - \int 1 \times e^t dt$$

$$\Rightarrow ye^{\tan^{-1} x} = te^t - e^t + C$$

$$\Rightarrow ye^{\tan^{-1} x} = \tan^{-1} x \cdot e^{\tan^{-1} x} - e^{\tan^{-1} x} + C$$

On dividing both sides by $e^{\tan^{-1} x}$, we get

$$y = \tan^{-1} x - 1 + Ce^{-\tan^{-1} x} \quad (1)$$

which is the required solution.

48. Solve the following differential equation

$$\frac{dy}{dx} + y = \cos x - \sin x.$$

Delhi 2009

Given differential equation is

$$\frac{dy}{dx} + y = \cos x - \sin x \quad \dots(i)$$

which is a linear differential equation of the form

$$\frac{dy}{dx} + Py = Q \quad \dots(ii)$$

On comparing Eqs. (i) and (ii), we get

$$P = 1 \text{ and } Q = \cos x - \sin x \quad (1)$$

$$\therefore \text{IF} = e^{\int 1 dx} = e^x \quad (1)$$

Now, solution of above equation is given by

$$y \times \text{IF} = \int (Q \times \text{IF}) dx + C$$

$$\therefore ye^x = \int e^x (\cos x - \sin x) dx$$

$$\Rightarrow ye^x = \int e^x \cos x dx - \int e^x \sin x dx$$

$$\Rightarrow ye^x = \left[\cos x \int e^x dx - \int \left\{ \frac{d}{dx} (\cos x) \int e^x dx \right\} dx - \int e^x \sin x dx \right]$$

[applying integration by parts in the first integral]

$$\Rightarrow ye^x = [e^x \cos x - \int -\sin x \cdot e^x dx] - \int e^x \sin x dx \quad (1)$$

$$\Rightarrow ye^x = e^x \cos x + \int e^x \sin x dx - \int e^x \sin x dx + C$$

$$\Rightarrow ye^x = e^x \cos x + C$$

On dividing both sides by e^x , we get

$$y = \cos x + Ce^{-x}$$

which is the required solution. (1)

49. Solve the following differential equation

$$\frac{dy}{dx} + 2y \tan x = \sin x.$$

All India 2008

Given differential equation is

$$\frac{dy}{dx} + 2y \tan x = \sin x \quad \dots(i)$$

which is a linear differential equation of the form

$$\frac{dy}{dx} + Py = Q \quad \dots(ii)$$

On comparing Eqs. (i) and (ii), we get

$$P = 2 \tan x \text{ and } Q = \sin x \quad (1)$$

$$\therefore \text{IF} = e^{\int 2 \tan x \, dx} = e^{2 \log |\sec x|}$$

$$= e^{\log \sec^2 x} = \sec^2 x \quad (1)$$

Now, solution of above equation is given by

$$y \times \text{IF} = \int (Q \times \text{IF}) \, dx + C$$

$$y \sec^2 x = \int \sin x \cdot \sec^2 x \, dx$$

$$\Rightarrow y \sec^2 x = \int \frac{\sin x}{\cos^2 x} \, dx \quad (1)$$

$$\Rightarrow y \sec^2 x = \int \sec x \tan x \, dx$$

$$\left[\because \frac{\sin x}{\cos^2 x} = \frac{\sin x}{\cos x} \cdot \frac{1}{\cos x} = \tan x \sec x \right]$$

$$\Rightarrow y \sec^2 x = \sec x + C$$

$$[\because \int \sec x \tan x \, dx = \sec x + C]$$

$$\therefore y = \frac{1}{\sec x} + \frac{C}{\sec^2 x}$$

$$\Rightarrow y = \cos x + C \cos^2 x \quad (1)$$

which is the required solution.

50. Solve the following differential equation

$$x^2 \frac{dy}{dx} = y^2 + 2xy.$$

All India 2008

Given differential equation is

$$x^2 \frac{dy}{dx} = y^2 + 2xy$$

which is a homogeneous differential equation as degree of each term is same in the equation.

Above equation can be written as

$$\frac{dy}{dx} = \frac{y^2 + 2xy}{x^2} \quad \dots(i)$$

On putting $y = vx$

$$\Rightarrow \frac{dy}{dx} = v + x \frac{dv}{dx} \quad (1)$$

in Eq. (i), we get

$$v + x \frac{dv}{dx} = \frac{v^2 x^2 + 2vx^2}{x^2} = v^2 + 2v$$

$$\Rightarrow v + x \frac{dv}{dx} = v^2 + 2v$$

$$\Rightarrow x \frac{dv}{dx} = v^2 + 2v - v \Rightarrow x \frac{dv}{dx} = v^2 + v$$

$$\Rightarrow \frac{dv}{v^2 + v} = \frac{dx}{x} \quad (1)$$

On integrating both sides, we get

$$\int \frac{dv}{v^2 + v} = \int \frac{dx}{x}$$

$$\Rightarrow \int \frac{dv}{v^2 + v + \frac{1}{4} - \frac{1}{4}} = \int \frac{dx}{x}$$

$$\Rightarrow \int \frac{dv}{\left(v + \frac{1}{2}\right)^2 - \left(\frac{1}{2}\right)^2} = \int \frac{dx}{x}$$

$$\Rightarrow \frac{1}{2 \times \frac{1}{2}} \log \left| \frac{v + \frac{1}{2} - \frac{1}{2}}{v + \frac{1}{2} + \frac{1}{2}} \right| = \log |x| + C$$

$$\left[\therefore \int \frac{dx}{x^2 - a^2} = \frac{1}{2a} \log \left| \frac{x - a}{x + a} \right| \right] \quad (1)$$

$$\begin{aligned}
 & \left[\int \frac{1}{x^2 - a^2} - \frac{2a}{x^2 - a^2} \log |x + a| \right] \\
 \Rightarrow & \log \left| \frac{v}{v+1} \right| - \log |x| = C \\
 \Rightarrow & \log \left| \frac{v}{(v+1) \cdot x} \right| = C \\
 & \left[\because \log m - \log n = \log \left(\frac{m}{n} \right) \right] \\
 \Rightarrow & \log \left| \frac{\frac{y}{x}}{\left(\frac{y}{x} + 1 \right) x} \right| = C \quad \left[\because y = vx \right] \\
 & \left[\because v = \frac{y}{x} \right] \\
 \Rightarrow & \log \left| \frac{y}{xy + x^2} \right| = C \quad (1)
 \end{aligned}$$

which is the required solution.

51. Solve the following differential equation

$$(x^2 - y^2) dx + 2xy dy = 0, \text{ given that } y = 1,$$

when $x = 1$.

Delhi 2008

Given differential equation is

$$(x^2 - y^2) dx + 2xy dy = 0$$

which is a homogeneous differential equation as degree of each term is same.

Above equation can be written as

$$(x^2 - y^2) dx = -2xy dy \Rightarrow \frac{dy}{dx} = \frac{y^2 - x^2}{2xy} \dots (i)$$

$$\text{On putting } y = vx \Rightarrow \frac{dy}{dx} = v + x \frac{dv}{dx} \quad (1)$$

in Eq. (i), we get

$$v + x \frac{dv}{dx} = \frac{v^2 x^2 - x^2}{2vx^2} = \frac{v^2 - 1}{2v}$$

$$\Rightarrow x \frac{dv}{dx} = \frac{v^2 - 1}{2v} - v$$

$$\frac{2v}{v^2 - 1} dv = \frac{1}{x} dx$$

$$= \frac{v^2 - 1 - 2v^2}{2v} = \frac{-1 - v^2}{2v} \quad (1)$$

$$\Rightarrow \frac{2v}{v^2 + 1} = -\frac{dx}{x}$$

On integrating both sides, we get

$$\int \frac{2v}{v^2 + 1} dv = -\int \frac{dx}{x}$$

On putting $v^2 + 1 = t \Rightarrow 2v dv = dt$

$$\therefore \int \frac{dt}{t} = -\log|x| + C$$

$$\Rightarrow \log|t| = -\log|x| + C$$

$$\Rightarrow \log|v^2 + 1| + \log|x| = C \quad [\because t = v^2 + 1]$$

$$\Rightarrow \log\left|\frac{y^2}{x^2} + 1\right| + \log|x| = C \quad \dots(ii)$$

$$\left[\because v = \frac{y}{x}\right] (1)$$

Also, given that $y = 1$, when $x = 1$.

On putting $x = 1$ and $y = 1$ in Eq. (ii), we get

$$\log 2 + \log 1 = C \Rightarrow C = \log 2 \quad [\because \log 1 = 0]$$

On putting $C = \log 2$ in Eq. (ii), we get

$$\log\left|\frac{y^2 + x^2}{x^2}\right| + \log x = \log 2$$

$$\Rightarrow \log\left|x\left(\frac{x^2 + y^2}{x^2}\right)\right| = \log 2$$

$$[\because \log m + \log n = \log mn]$$

$$\Rightarrow \log\left|\frac{x^2 + y^2}{x}\right| = \log 2 \Rightarrow x^2 + y^2 = 2x \quad (1)$$

which is the required solution.

52. Solve the following differential equation

$$\frac{dy}{dx} = \frac{x(2y - x)}{x(2y + x)}, \text{ if } y = 1, \text{ when } x = 1.$$

Delhi 2008

Given differential equation is

$$\frac{dy}{dx} = \frac{x(2y - x)}{x(2y + x)} \Rightarrow \frac{dy}{dx} = \frac{2xy - x^2}{2xy + x^2} \quad \dots(i)$$

which is a homogeneous differential equation because each term of numerator and denominator have same degree.

$$\text{On putting } y = vx \Rightarrow \frac{dy}{dx} = v + x \frac{dv}{dx} \quad (1)$$

in Eq. (i), we get

$$\begin{aligned} v + x \frac{dv}{dx} &= \frac{2vx^2 - x^2}{2vx^2 + x^2} = \frac{2v - 1}{2v + 1} \\ v + x \frac{dv}{dx} &= \frac{2v - 1}{2v + 1} \\ \Rightarrow x \frac{dv}{dx} &= \frac{2v - 1}{2v + 1} - v \\ \Rightarrow x \frac{dv}{dx} &= \frac{2v - 1 - 2v^2 - v}{2v + 1} \\ \frac{2v + 1}{2v^2 - v + 1} dv &= - \frac{dx}{x} \end{aligned}$$

On integrating both sides, we get

$$\begin{aligned} \int \frac{2v + 1}{2v^2 - v + 1} dv &= - \int \frac{dx}{x} \\ \Rightarrow I &= - \log |x| + C \quad \dots(ii) \end{aligned}$$

where, $I = \int \frac{2v + 1}{2v^2 - v + 1} dv$

$$\text{Let } 2v + 1 = A \cdot \frac{d}{dv} (2v^2 - v + 1) + B$$

$$\Rightarrow 2v + 1 = A(4v - 1) + B \quad \dots(iii)$$

On comparing coefficients of v and constants from both sides, we get

$$4A = 2$$

$$\Rightarrow A = \frac{1}{2} \text{ and } -A + B = 1$$

$$\Rightarrow -\frac{1}{2} + B = 1 \Rightarrow B = \frac{3}{2}$$

On putting $A = \frac{1}{2}$ and $B = \frac{3}{2}$ in Eq. (iii), we get

$$2v + 1 = \frac{1}{2}(4v - 1) + \frac{3}{2} \quad (1)$$

On integrating both sides, we get

$$I = \int \frac{2v + 1}{2v^2 - v + 1} dv$$

$$\Rightarrow I = \int \frac{\frac{1}{2}(4v - 1) + \frac{3}{2}}{2v^2 - v + 1} dv$$

$$\Rightarrow I = \frac{1}{2} \int \frac{4v - 1}{2v^2 - v + 1} dv + \frac{3}{2} \int \frac{dv}{2v^2 - v + 1}$$

$$\Rightarrow I = \frac{1}{2} \log |2v^2 - v + 1| + \frac{3}{4} \int \frac{dv}{v^2 - \frac{v}{2} + \frac{1}{2}}$$

$$\left[\begin{array}{l} \because \int \frac{4v - 1}{2v^2 - v + 1} dv \Rightarrow \text{put } 2v^2 - v + 1 = t \\ \Rightarrow (4v - 1) dv = dt \\ \text{then } \int \frac{dt}{t} = \log |t| = \log |2v^2 - v + 1| \end{array} \right]$$

$$\Rightarrow I = \frac{1}{2} \log |2v^2 - v + 1|$$

$$+ \frac{3}{4} \int \frac{dv}{v^2 - \frac{1}{2}v + \frac{1}{2} + \frac{1}{16} - \frac{1}{16}}$$

$$= \frac{1}{2} \log |2v^2 - v + 1| + \frac{3}{4} \int \frac{dv}{\left(v - \frac{1}{4}\right)^2 + \frac{7}{16}}$$

$$= \frac{1}{2} \log |2v^2 - v + 1| + \frac{3}{4} \int \frac{dv}{\left(v - \frac{1}{4}\right)^2 + \left(\frac{\sqrt{7}}{4}\right)^2} \quad (1/2)$$

$$\Rightarrow I = \frac{1}{2} \log |2v^2 - v + 1| + \frac{3}{4} \times \frac{4}{\sqrt{7}} \tan^{-1} \left(\frac{v - \frac{1}{4}}{\frac{\sqrt{7}}{4}} \right)$$

$$\left[\because \int \frac{dx}{x^2 + a^2} = \frac{1}{a} \tan^{-1} \frac{x}{a} + C \right]$$

$$\Rightarrow I = \frac{1}{2} \log |2v^2 - v + 1| + \frac{3}{\sqrt{7}} \tan^{-1} \left(\frac{4v - 1}{\sqrt{7}} \right)$$

On putting the value of I in Eq. (ii), we get

$$\frac{1}{2} \log |2v^2 - v + 1| + \frac{3\sqrt{7}}{7} \tan^{-1} \left(\frac{4v - 1}{\sqrt{7}} \right)$$

$$= -\log |x| + C \quad (1/2)$$

$$\Rightarrow \frac{1}{2} \log \left| \frac{2y^2}{x^2} - \frac{y}{x} + 1 \right| + \frac{3\sqrt{7}}{7} \tan^{-1} \left(\frac{\frac{4y}{x} - 1}{\frac{\sqrt{7}}{x}} \right)$$

$$= -\log |x| + C \quad \left[\because \text{put } v = \frac{y}{x} \right]$$

$$\Rightarrow \frac{1}{2} \log \left| \frac{2y^2}{x^2} - \frac{y}{x} + 1 \right| + \frac{3\sqrt{7}}{7} \tan^{-1} \left(\frac{4y - x}{\sqrt{7} \cdot x} \right)$$

$$= -\log |x| + C \quad \dots(iv)$$

Also, given that $y = 1$, when $x = 1$.

On putting $x = 1$ and $y = 1$ in Eq. (iv), we get

$$\frac{1}{2} \log |2| + \frac{3\sqrt{7}}{7} \tan^{-1} \left(\frac{3}{\sqrt{7}} \right) = -\log 1 + C$$

$$\Rightarrow \frac{1}{2} \log 2 + \frac{3\sqrt{7}}{7} \tan^{-1} \left(\frac{3}{\sqrt{7}} \right) = C \quad [\because \log 1 = 0]$$

$$\rightarrow \frac{1}{2} \log 2 + \frac{1}{2} \log x + \frac{3\sqrt{7}}{7} \tan^{-1} \left(\frac{3}{\sqrt{7}} \right)$$

On putting the value of C in Eq. (iv), we get

$$\frac{1}{2} \log \left(\frac{2y^2 - xy + x^2}{x^2} \right) + \frac{3\sqrt{7}}{7} \tan^{-1} \left(\frac{4y - x}{\sqrt{7}x} \right)$$

$$= -\log |x| + \frac{1}{2} \log 2 + \frac{3\sqrt{7}}{7} \tan^{-1} \left(\frac{3}{\sqrt{7}} \right)$$

$$\Rightarrow \log \left(\frac{2y^2 - xy + x^2}{x^2} \right)^{1/2} + \log x - \log (2)^{1/2}$$

$$= \frac{3\sqrt{7}}{7} \left[\tan^{-1} \left\{ \frac{\frac{3}{\sqrt{7}} - \left(\frac{4y - x}{\sqrt{7}x} \right)}{1 + \frac{3}{\sqrt{7}} \cdot \left(\frac{4y - x}{\sqrt{7}x} \right)} \right\} \right]$$

$$\left[\because \tan^{-1} A - \tan^{-1} B = \tan^{-1} \left(\frac{A - B}{1 + AB} \right) \right] \quad (1/2)$$

$$\Rightarrow \log (2y^2 - xy + x^2)^{1/2} - \log \sqrt{2}$$

$$= \frac{3\sqrt{7}}{7} \tan^{-1} \left[\frac{(4x - 4y) \cdot \sqrt{7}}{4x + 12y} \right]$$

$$\left[\because \log \left(\frac{2y^2 - xy + x^2}{x^2} \right) = \log (2y^2 - xy + x^2)^{1/2} \right. \\ \left. = \log (2y^2 - xy + x^2)^{1/2} - \log x \right]$$

$$\Rightarrow \log \sqrt{\frac{2y^2 - xy + x^2}{2}}$$

$$= \frac{3\sqrt{7}}{7} \tan^{-1} \left(\frac{(x - y) \cdot \sqrt{7}}{x + 3y} \right)$$

$$\Rightarrow \log \sqrt{\frac{2y^2 - xy + x^2}{2}}$$

$$= \frac{3\sqrt{7}}{7} \tan^{-1} \left[\frac{\sqrt{7}x - \sqrt{7}y}{x + 3y} \right] \quad (1/2)$$

which is the required solution.

6 Marks Questions

- 53.** Find the particular solution of the differential equation $(3xy + y^2)dx + (x^2 + xy)dy = 0$, for $x = 1$ and $y = 1$. Delhi 2013C

Given differential equation is

$$(3x^2 + y^2)dx + (x^2 + xy)dy = 0$$

It can be rewritten as $\frac{dy}{dx} = -\frac{3xy + y^2}{x^2 + xy}$... (i)

which is a homogeneous differential equation of degree 2.

On putting $y = vx \Rightarrow \frac{dy}{dx} = v + x \frac{dv}{dx}$

in Eq.(i), we get $v + x \frac{dv}{dx} = -\frac{3vx^2 + v^2x^2}{x^2 + vx^2}$

$$\Rightarrow x \frac{dv}{dx} = -\frac{3v + v^2}{1 + v} - v$$

$$\Rightarrow x \frac{dv}{dx} = -\left(\frac{3v + v^2 + v + v^2}{1 + v} \right) \quad (1)$$

$$\Rightarrow x \frac{dv}{dx} = -\left(\frac{2v^2 + 4v}{1 + v} \right) \Rightarrow \frac{(1 + v)dv}{2(v^2 + 2v)} = -\frac{dx}{x}$$

On integrating both sides, we get

$$\int \frac{1 + v}{2(v^2 + 2v)} dv = -\int \frac{dx}{x} \quad \dots (ii) \quad (1)$$

Again, put $v^2 + 2v = z \Rightarrow (2v + 2)dv = dz$

$$\Rightarrow (1 + v)dv = \frac{dz}{2}$$

Then, Eq. (ii) becomes,

$$\int \frac{1}{2} \times \frac{dz}{2z} = - \int \frac{dx}{x} \quad (1)$$

$$\Rightarrow \frac{1}{4} \log |z| = -\log |x| + \log |C|$$

$$\Rightarrow \frac{1}{4} [\log |z| + 4 \log |x|] = \log |C|$$

$$\Rightarrow \log |zx^4| = 4 \log |C|$$

$$\Rightarrow zx^4 = C^4 = C_1 \quad zx^4 = C_1$$

where, $C_1 = C^4$

$$\Rightarrow x^4(v^2 + 2v) = C_1 \quad [\text{put } z = v^2 + 2v]$$

$$\Rightarrow x^4 \left(\frac{y^2}{x^2} + \frac{2y}{x} \right) = C_1 \left[\text{put } v = \frac{y}{x} \right] \dots (iii) \quad (1)$$

Also, given that $y = 1$ for $x = 1$.

On putting $x = 1$ and $y = 1$ in Eq. (iii), we get

$$1 \left(\frac{1}{1} + \frac{2}{1} \right) = C_1$$

$$\Rightarrow C_1 = 3 \quad (1)$$

Also, given that $y = 1$ for $x = 1$.

So, on putting $C_1 = 3$ in Eq. (iii), we get

$$x^4 \left(\frac{y^2}{x^2} + \frac{2y}{x} \right) = 3 \Rightarrow y^2 x^2 + 2yx^3 = 3 \quad (1)$$

which is the required particular solution.

- 54.** Show that the differential equation $2ye^{x/y} dx + (y - 2xe^{x/y}) dy = 0$ is homogeneous. Find the particular solution of this differential equation, given that $x=0$, when $y=1$.

HOTS; Delhi 2013



Firstly, replace x by λx and y by λy in $f(x, y)$ of given differential equation to check that it is homogeneous. If it is homogeneous, then put $x = vy$ and $\frac{dx}{dy} = v + y \frac{dv}{dy}$ and then solve.

Given differential equation is

$2y e^{x/y} dx + (y - 2x e^{x/y}) dy = 0$. It can be written as

$$\frac{dx}{dy} = \frac{2x e^{x/y} - y}{2y e^{x/y}} \quad \dots(i)$$

$$\text{Let } F(x, y) = \frac{\left(2x e^{\frac{x}{y}} - y \right)}{\left(2y e^{\frac{x}{y}} \right)}$$

On replacing x by λx and y by λy both sides, we get

$$F(\lambda x, \lambda y) = \frac{\left(2\lambda x e^{\frac{\lambda x}{\lambda y}} - \lambda y \right)}{\left(2\lambda y e^{\frac{\lambda x}{\lambda y}} \right)}$$

$$\Rightarrow F(\lambda x, \lambda y) = \frac{\lambda(2x e^{x/y} - y)}{\lambda(2y e^{x/y})} = \lambda^0 [F(x, y)] \quad (1)$$

Thus, $F(x, y)$ is a homogeneous function of degree zero. Therefore, the given differential equation is a homogeneous differential equation. (1)

To solve it, put $x = vy$

$$\Rightarrow \frac{dx}{dy} = v + y \frac{dv}{dy} \quad (1/2)$$

$$\begin{aligned}
 &\text{in Eq.(i), we get } v + y \frac{dv}{dy} = \frac{2ve^v - 1}{2e^v} \\
 &\Rightarrow y \frac{dv}{dy} = \frac{2ve^v - 1}{2e^v} - v = \frac{2ve^v - 1 - 2ve^v}{2e^v} \\
 &\Rightarrow 2e^v dv = \frac{-dy}{y} \quad (1)
 \end{aligned}$$

On integrating both sides, we get

$$\int 2e^v dv = - \int \frac{dy}{y} \Rightarrow 2e^v = -\log|y| + C$$

Now, replace v by $\frac{x}{y}$, we get

$$2e^{x/y} + \log|y| = C \quad \dots(ii) \quad (1\frac{1}{2})$$

Also, given that $x = 0$, when $y = 1$.

On substituting $x = 0$ and $y = 1$ in Eq. (ii), we get

$$2e^0 + \log|1| = C \Rightarrow C = 2$$

On substituting the value of C in Eq. (ii), we get

$$2e^{x/y} + \log|y| = 2$$

which is the required particular solution of the given differential equation. (1)

55. Show that the differential equation

$$x \frac{dy}{dx} \sin\left(\frac{y}{x}\right) + x - y \sin\left(\frac{y}{x}\right) = 0 \text{ is homogeneous.}$$

Find the particular solution of this differential

equation, given that $x = 1$, when $y = \frac{\pi}{2}$. Delhi 2013

Given differential equation is

$$x \frac{dy}{dx} \sin\left(\frac{y}{x}\right) = y \sin\left(\frac{y}{x}\right) - x \Rightarrow \frac{dy}{dx} = \frac{y}{x} - \frac{1}{\sin \frac{y}{x}} \dots (i)$$

$$\left[\text{dividing both sides by } x \sin\left(\frac{y}{x}\right) \right]$$

Let $(x, y) = \frac{y}{x} - \frac{1}{\sin \frac{y}{x}}$

On replacing x by λx and y by λy on both sides, we get

$$F(\lambda x, \lambda y) = \frac{\lambda y}{\lambda x} - \frac{1}{\sin \frac{\lambda y}{\lambda x}} = \lambda^0 \left(\frac{y}{x} - \frac{1}{\sin \frac{y}{x}} \right) = \lambda^0 F(x, y)$$

So, given differential equation is homogeneous. (2)

On putting $y = vx$

$$\Rightarrow \frac{dy}{dx} = v + x \frac{dv}{dx} \text{ in Eq.(i), we get} \quad (1)$$

$$v + x \frac{dv}{dx} = v - \frac{1}{\sin v}$$

$$\Rightarrow x \frac{dv}{dx} = -\frac{1}{\sin v} \Rightarrow \sin v \, dv = -\frac{dx}{x}$$

On integrating both sides, we get

$$\int \sin v \, dv = -\int \frac{dx}{x}$$

$$\Rightarrow -\cos v = -\log|x| + C$$

$$\Rightarrow -\cos y/x = -\log|x| + C \left[\because v = \frac{y}{x} \right] \quad (1\frac{1}{2}) \dots (ii)$$

Also, given that $x = 1$, when $y = \frac{\pi}{2}$.

On putting $x = 1$ and $y = \frac{\pi}{2}$ in Eq. (ii), we get

$$-\cos\left(\frac{\pi}{2}\right) = -\log|1| + C$$

$$\Rightarrow -0 = -0 + C \Rightarrow C = 0$$

On putting the value of C in Eq. (ii), we get

$$\cos \frac{y}{x} = \ln|x|$$

which is the required solution. (1½)

56. Find the particular solution of the

differential equation $\frac{dx}{dy} + x \cot y = 2y + y^2 \cot y$,

($y \neq 0$), given that $x = 0$, when $y = \frac{\pi}{2}$.

All India 2013

Given differential equation is

$$\frac{dx}{dy} + x \cot y = 2y + y^2 \cot y, (y \neq 0)$$

which is a linear differential equation.

On comparing with $\frac{dx}{dy} + Px = Q$, we get

$$P = \cot y \text{ and } Q = 2y + y^2 \cot y$$

$$\therefore \text{IF} = e^{\int P dy} = e^{\int \cot y dy} = e^{\log \sin y} = \sin y \quad (1\frac{1}{2})$$

Now, the solution of above differential equation is given by

$$x \cdot (\text{IF}) = \int Q \cdot (\text{IF}) dy + C$$

$$\begin{aligned} \therefore x \sin y &= \int (2y + y^2 \cot y) \sin y dy + C \\ &= 2 \int y \sin y dy + \int \underset{\text{I}}{y^2} \underset{\text{II}}{\cos y} dy + C \\ &= 2 \int y \sin y dy + y^2 \int \cos y dy \\ &\quad - \int \left[\left(\frac{d}{dy} y^2 \right) \int \cos y dy \right] dy + C \end{aligned}$$

[using integration by parts in second integral]

$$\begin{aligned} &= 2 \int y \sin y dy + y^2 \sin y - 2 \int y \sin y dy + C \\ &= y^2 \sin y + C \end{aligned}$$

$$\Rightarrow x \sin y = y^2 \sin y + C \quad \dots(i) \quad (2)$$

Also, given that $x = 0$, when $y = \frac{\pi}{2}$.

On putting $x = 0$ and $y = \frac{\pi}{2}$ in Eq. (i), we get

$$0 = \left(\frac{\pi}{2}\right)^2 \sin \frac{\pi}{2} + C \Rightarrow C = -\frac{\pi^2}{4} \quad (1/2)$$

On putting the value of C in Eq. (i), we get

$$x \sin y = y^2 \sin y - \frac{\pi^2}{4} \Rightarrow x = y^2 - \frac{\pi^2}{4} \cdot \operatorname{cosec} y$$

which is required particular solution of given differential equation. (2)

57. Show that the differential equation

$$\left[x \sin^2 \left(\frac{y}{x} \right) - y \right] dx + x dy = 0 \text{ is homogeneous.}$$

Find the particular solution of this differential equation, given that

$$y = \frac{\pi}{4}, \text{ when } x = 1.$$

All India 2013

Given differential equation is

$$\left[x \sin^2 \left(\frac{y}{x} \right) - y \right] dx + x dy = 0$$

$$\Rightarrow \frac{dy}{dx} = \frac{y - x \sin^2 \left(\frac{y}{x} \right)}{x} \quad \dots(i)$$

Let $F(x, y) = \frac{y - x \sin^2 \left(\frac{y}{x} \right)}{x}$

On replacing x by λx and y by λy both sides, we get

$$F(\lambda x, \lambda y) = \frac{\lambda \left[y - x \sin^2 \left(\frac{y}{x} \right) \right]}{\lambda x} = \lambda^0 [F(x, y)]$$

Thus, given differential equation is a homogeneous differential equation. **(1)**

On putting $y = vx \Rightarrow \frac{dy}{dx} = v + x \frac{dv}{dx}$ in Eq. (i), we get

$$\begin{aligned} v + x \frac{dv}{dx} &= \frac{vx - x \sin^2 \left(\frac{vx}{x} \right)}{x} \\ \Rightarrow v + x \frac{dv}{dx} &= v - \sin^2 v \Rightarrow x \frac{dv}{dx} = -\sin^2 v \\ \Rightarrow \operatorname{cosec}^2 v dv &= \frac{-dx}{x} \quad (2) \end{aligned}$$

On integrating both sides, we get

$$\begin{aligned} \int \operatorname{cosec}^2 v dv + \int \frac{dx}{x} &= 0 \\ \Rightarrow -\cot v + \log |x| &= C \\ \Rightarrow -\cot \left(\frac{y}{x} \right) + \log |x| &= C \left[\because v = \frac{y}{x} \right] \dots(ii) \end{aligned}$$

Also, given that, $y = \frac{\pi}{4}$, when $x = 1$.

On putting $x = 1$ and $y = \frac{\pi}{4}$, in Eq. (ii), we get

$$-\cot\left(\frac{\pi}{4}\right) + \log|1| = C \quad (2)$$

$$\Rightarrow C = -1 \quad \left[\because \cot \frac{\pi}{4} = 1 \right]$$

On putting the value of C in Eq. (ii), we get

$$-\cot\left(\frac{y}{x}\right) + \log|x| = -1$$

$$\Rightarrow 1 + \log|x| - \cot\left(\frac{y}{x}\right) = 0$$

which is the required particular solution of given differential equation. (1)

- 58.** Find the particular solution of the differential equation $(\tan^{-1} y - x)dy = (1 + y^2)dx$, given that $x=0$, when $y=0$. All India 2013

Given differential equation is

$$(\tan^{-1} y - x)dy = (1 + y^2)dx$$

$$\Rightarrow \frac{\tan^{-1} y - x}{1 + y^2} = \frac{dx}{dy} \Rightarrow \frac{dx}{dy} = \frac{-x}{1 + y^2} + \frac{\tan^{-1} y}{1 + y^2}$$

$$\Rightarrow \frac{dx}{dy} + \frac{1}{1 + y^2} \cdot x = \frac{\tan^{-1} y}{1 + y^2}$$

which is a linear differential equation of first order. (1)

On comparing with $\frac{dx}{dy} + Px = Q$, we get

$$P = \frac{1}{1 + y^2} \quad \text{and} \quad Q = \frac{\tan^{-1} y}{1 + y^2}$$

$$\therefore \quad \text{IF} = e^{\int P dy} = e^{\int \frac{dy}{1+y^2}} = e^{\tan^{-1} y} \quad (1)$$

Now, solution of above differential equation is given by

$$\begin{aligned} x \cdot (\text{IF}) &= \int Q \cdot (\text{IF}) dy + C \\ \Rightarrow x e^{\tan^{-1} y} &= \int \frac{\tan^{-1} y}{1+y^2} \times e^{\tan^{-1} y} + C \quad (1) \end{aligned}$$

$$\text{On putting } t = \tan^{-1} y \Rightarrow dt = \frac{1}{1+y^2} dy$$

$$\begin{aligned} \therefore x \cdot e^{\tan^{-1} y} &= \int t \cdot e^t dt + C \\ \Rightarrow x \cdot e^{\tan^{-1} y} &= t \cdot e^t - \int 1 \cdot e^t dt + C \\ &\quad [\text{using integration by parts}] \\ \Rightarrow x \cdot e^{\tan^{-1} y} &= t \cdot e^t - e^t + C \\ \Rightarrow x \cdot e^{\tan^{-1} y} &= (\tan^{-1} y - 1) e^{\tan^{-1} y} + C \dots (i) \quad (1) \end{aligned}$$

Also, given that, when $x=0$, then $y=0$.

On putting $x=0, y=0$ in Eq. (i), we get

$$\begin{aligned} 0 &= (\tan^{-1} 0 - 1) e^{\tan^{-1} 0} + C \\ \Rightarrow 0 &= (0 - 1) e^0 + C \Rightarrow 0 = (0 - 1) \cdot 1 + C \\ \Rightarrow C &= 1 \quad (1) \end{aligned}$$

On putting the value of C in Eq. (i), we get

$$\begin{aligned} \therefore x \cdot e^{\tan^{-1} y} &= (\tan^{-1} y - 1) \cdot e^{\tan^{-1} y} + 1 \\ \Rightarrow x &= \tan^{-1} y - 1 + e^{-\tan^{-1} y} \end{aligned}$$

which is the required particular solution of the differential equation. (1)