

Polynomials

Question 1.

If a polynomial $p(y)$ is divided by $y + 2$, then which of the following can be the remainder:

- (a) $y + 1$
- (b) $2y + 3$
- (c) 5
- (d) $y - 1$

Answer: (c) 5

When $p(y)$ is divided by $y + 2$, then the degree of remainder $<$ deg of $(y + 2)$

Question 2.

If a polynomial $p(x)$ is divided by $b - ax$; the remainder is the value of $p(x)$ at $x =$

- (a) a
- (b) $\frac{b}{a}$
- (c) $\frac{-b}{a}$
- (d) $\frac{a}{b}$

Answer: (b) $\frac{b}{a}$

$$b - ax = 0$$

$$x = \frac{b}{a}$$

Question 3.

If the polynomials $ax^3 + 4x^2 + 3x - 4$ and $x^3 - 4x + a$, leave the same remainder when divided by $(x - 3)$, then value of a is :

- (a) $2b$
- (b) -1
- (c) 1
- (d) $-2b$

Answer: (b) -1

$$p(x) = ax^3 + 4x^2 + 3x - 4$$

$$q(x) = x^3 - 4x + a$$



$$p(3) = q(3)$$
$$a = -1$$

Question 4.

If $p(x) = 2x^4 - ax^3 + 4x^2 + 2x + 1$ is a multiple of $1 - 2x$, then find the value of a :

- (a) 25
- (b) $\frac{1}{2}$
- (c) $-\frac{1}{2}$
- (d) 8

Answer: (a) 25

$p(x)$ is a multiple of $1 - 2x$.

$1 - 2x$ is a factor of $p(x)$

Question 5.

If -2 is a zero of $p(x) = (ax^3 + bx^2 + x - 6)$ and $p(x)$ leaves a remainder 4 when divided by $(x - 2)$, then the values of a and b are (respectively):

- (a) $a = 2, b = 2$
- (b) $a = 0, b = -2$
- (c) $a = 0, b = 2$
- (d) $a = 0, b = 0$

Answer: (c) $a = 0, b = 2$

If -2 is a zero \Rightarrow

$$p(-2) = 0$$

$$\Rightarrow -2a + b = 2$$

Also, $p(2) = 4$

$$2a + b = 2 \Rightarrow a = 0 \text{ and } b = 2$$

Question 6.

If $x^{101} + 1001$ is divided by $x + 1$, then remainder is:

- (a) 0
- (b) 1
- (c) 1490
- (d) 1000

Answer: (d) 1000

$p(x)$ is divided by $x + 1$

$$p(-1) = (-1)^{101} + 1001 = 1000$$



Question 7.

If one zero of a polynomial $p(x) = ax^2 + bx + c$ ($a \neq 0$) is zero, then, which of the following is correct:

- (a) $b = 0$
- (b) $c = 0$
- (c) other zero is also zero
- (d) Nothing can be said about $p(x)$.

Answer: (b) $c = 0$

let $\alpha = 0$

Product of the roots $= \alpha s = 0$

$$= \frac{c}{a} = 0$$

Question 8.

If α, s are the zeroes of $x^2 - lx + m$, then

$$\frac{\alpha}{s} + \frac{s}{\alpha}$$

(a) $\frac{l^2 - 2m}{m}$

(b) $\frac{l^2 + 2m}{m}$

(c) $\frac{l - 2m}{m}$

(d) $l^2 - 2m$

Answer: (a) $\frac{l^2 - 2m}{m}$

$$\alpha + s = l$$

$$\alpha s = m$$

$$\Rightarrow \alpha^2 + s^2 = (\alpha + s)^2 - 2\alpha s = l^2 - 2m$$

$$\Rightarrow \frac{\alpha}{s} + \frac{s}{\alpha} = \frac{\alpha^2 + s^2}{\alpha s} = \frac{l^2 - 2m}{m}$$

Question 9.

sum of the squares of the zeroes of the polynomial $p(x) = x^2 + 7x - k$ is 25, find k .

- (a) 12
- (b) 49
- (c) -24
- (d) -12

Answer: (d) -12

$$p(x) = x^2 + 7x - k$$

let α, s be the zeroes

$$\alpha + s = -7$$

$$\alpha s = -k$$

$$\alpha^2 + s^2 = 25$$

$$(\alpha^2 + s) - 2\alpha s = 25$$

$$49 + 2k = 25$$

$$k = -12$$

Question 10.

If one zero of $3x^2 - 8x + 2k + 1$ is seven times the other, find k.

- (a) $\frac{2}{3}$
- (b) $\frac{1}{3}$
- (c) $\frac{4}{3}$
- (d) $\frac{5}{3}$

Answer: (a) $\frac{2}{3}$

$$\alpha + 7\alpha = 8\alpha = \frac{8}{3}$$

$$\alpha = \frac{1}{3}$$

$$k = \frac{2}{3}$$

Question 11.

Let, α, s, v be the zeroes of $x^3 + 4x^2 + x - 6$ such that product of two of the zeroes is 6. Find the third zero.

- (a) 6
- (b) 2
- (c) 4
- (d) 1

Answer: (a) 6

$$\alpha s v = 6,$$

$$\alpha s = 6$$

$$\Rightarrow v = 1$$

Question 12.

If α, s are the zeroes of $x^2 - 8x + \lambda$, such that $\alpha - s = 2$, then $\lambda =$

- (a) 8
- (b) 22

- (c) 60
(d) 15

Answer: (d) 15

$$\alpha + s = 8,$$

$$\alpha s = \lambda$$

$$\alpha - s = 2$$

$$\Rightarrow (\alpha - s)^2 = 4$$

$$\Rightarrow \alpha^2 + s^2 - 2\alpha s = 4$$

$$\Rightarrow (\alpha + s)^2 - 4\alpha s = 4$$

$$\Rightarrow 64 - 4\lambda = 4$$

$$\Rightarrow 4\lambda = 60$$

$$\Rightarrow X = 15$$

Question 13.

Find a and b so that the polynomial $6x^4 + 8x^3 - 5x^2 + ax + b$ is exactly divisible by $2x^2 - 5$.

- (a) $a = 20, b = -25$
(b) $a = 4, b = -5$
(c) $a = 20, b = 5$
(d) $a = -20, b = -25$

Answer: (d) $a = -20, b = -25$

Divide the given polynomial by $2x^2 - 5$ get the remainder as $(20 + a)x + (b + 25)$ which should be zero

Question 14.

If α, s are the zeroes of $p(x) = 2x^2 - 5x + 7$, write a polynomial with zeroes $2\alpha + 3s$ and $3\alpha + 2s$.

- (a) $k(x^2 + \frac{5}{2}x - 41)$
(b) $k(x^2 - \frac{5}{2}x + 41)$
(c) $k(x^2 - \frac{5}{2}x - 41)$
(d) $k(-x^2 + \frac{5}{2}x + 41)$

Answer: (b) $k(x^2 - \frac{5}{2}x + 41)$

$$\alpha + s = \frac{5}{2}$$

$$\alpha s = \frac{7}{2}$$

$$k(x^2 - \frac{5}{2}x + 41)$$

Question 15.

If sum of the two zeroes of a cubic polynomial $x^3 - ax^2 + bx - c$, is zero, then which of the following is true:

- (a) $ab = c$
- (b) $a - b = c$
- (c) $ab = \frac{c}{2}$
- (d) $a = \frac{b}{c}$

Answer: (a) $ab = c$

Let, α, s, v be the roots $\Rightarrow \alpha + s + v = a$

$v = a$

now v is a zero

$ab = c$

Question 16.

If α, s are the zeroes of $p(x) = 2x^2 + 5x + k$ such that, $\alpha^2 + s^2 + \alpha s = \frac{21}{4}$, then k equals,

- (a) 12
- (b) 4
- (c) 2
- (d) -12

Answer: (c) 2

$$\alpha + s = -\frac{5}{2}$$

$$\alpha s = \frac{k}{2}$$

$$\alpha^2 + s^2 + \alpha s = \frac{21}{4}$$

$$(\alpha + s)^2 - \alpha s = \frac{21}{4}$$

$$\frac{25}{4} - \frac{k}{2} = \frac{21}{4}$$

$$k = 2$$

Question 17.

If α, s are the zeroes of $x^2 + px + q$, then a polynomial having zeroes $\frac{1}{\alpha}$ and $\frac{1}{s}$ is,

- (a) $x^2 + px + q$
- (b) $x^2 + qx + p$
- (c) $px^2 + qx + 1$
- (d) $qx^2 + px + 1$

Answer: (d) $qx^2 + px + 1$

$$\alpha + s = -p$$

$$\alpha s = q$$


$$S = \frac{1}{\alpha} + \frac{1}{s} = \frac{\alpha + s}{\alpha s} = \frac{-p}{q}$$

$$P = \frac{1}{\alpha} \cdot \frac{1}{s} = \frac{1}{q}$$

$$k \left(x^2 + \frac{p}{q}x + \frac{1}{q} \right) = \frac{k}{q} (qx^2 + px + 1)$$

Question 18.

Find the number of zeros in the graph given:




- (a) 3
- (b) 2
- (c) 1
- (d) 0

Answer: (b) 2

Since the graph meets X-axis at two points -2 and 1, thus it has 2 zeroes.

Question 19.

Write the zero of the polynomial $p(x)$, whose graph is given :



- (a) 1
- (b) 0
- (c) -1
- (d) -2

Answer: (b) 0

Since the graph meets X-axis at $x = 0$
 \Rightarrow Zero of $p(x)$ is 'O' \Rightarrow Correct option is (b).

Question 20.

If α, s, v are the zeros of the polynomial $2x^3 - x^2 + 3x - 1$, find the value of $(\alpha sv) + (\alpha s + sv + v\alpha)$.

- (a) 2
- (b) $\frac{3}{2}$
- (c) $\frac{1}{2}$
- (d) 0

Answer: (a) 2

$$p(x) = 2x^3 - x^2 + 3x - 1$$

$$\alpha sv = -d/a = \frac{1}{2}$$

$$\alpha s + sv + v\alpha = c/a = \frac{3}{2}$$

$$\alpha s + sv + v\alpha + \alpha sv = \frac{3}{2} + \frac{1}{2} = 2$$

Question 21.

If the zeros of the polynomial $x^3 - 3x^2 + x + 1$ are $p - q, p$ and $p + q$. Find the value of q .

- (a) 1
- (b) 0
- (c) 2
- (d) $\pm\sqrt{2}$

Answer: (d) $\pm\sqrt{2}$

$$x^3 - 3x^2 + x + 1$$

zeroes are $p - q, p, p + q$

$$\text{sum of zeroes} = (p - q) + p + (p + q)$$

$$= 3p$$

$$= 3$$

$$\alpha + s + v = \frac{-b}{a}$$

$$\text{further} = \alpha s + sv + v\alpha = \frac{c}{a}$$

$$(p - q)p + p(p + q) + (p - q)(p + q) = 1$$

$$q = \pm\sqrt{2}$$

Question 22.

A quadratic polynomial has :

- (a) at least 2 zeros
- (b) exactly 2 zeros
- (c) at most 2 zeros
- (d) exactly 1 zero


Answer: (c) at most 2 zeros

A quadratic polynomial has atmost two zeroes.


Question 23.




Which of the following Linear Graphs has no zero?

(a)



(b)





Answer:
as it does not meet X axis.

Question 24.

If α, s are the roots of $cx^2 - bx + a = 0$ ($c \neq 0$), then $\alpha + s$ is:

- (a) $\frac{-b}{a}$
- (b) $\frac{b}{a}$
- (c) $\frac{c}{a}$
- (d) $\frac{b}{c}$

Answer: (d) $\frac{b}{c}$

sum of the roots $= -\frac{\text{coefficient of } x}{\text{coefficient of } x^2} = \frac{b}{c}$

Question 25.

If $P(x)$ and $D(r)$ are any two polynomials such that $D(x) \neq 0$, there exists unique polynomial $Q(x)$ and $R(x)$ such that, $P(x) = D(x) \cdot Q(x) + R(x)$ where :

- (a) $R(x) = 0$ and $\deg R(x) > \deg Q(x)$
- (b) $R(x) = 0$ or $\deg R(x) > \deg D(x)$
- (c) $\deg R(x) < \deg Q(x)$
- (d) $R(x) = 0$ or $\deg R(x) < \deg D(x)$

Answer: (b) $R(x) = 0$ or $\deg R(x) > \deg D(x)$
division algorithm

Question 26.

When we divide $x^3 + 5x + 7$ by $x^4 - 7x^2 - 6$ then quotient and remainder are (respectively):

- (a) $0, x^3 + 5x + 7$
- (b) $x, 2x + 3$
- (c) $1, x^4 - 7x^2 - 6$
- (d) $x^2, 4x - 9$

Answer: (a) $0, x^3 + 5x + 7$

Degree of the divisor is more than the degree of the dividend = quotient is zero and the remainder is $x^3 + 5x + 7$

Question 27.

The value of b , for which $2x^3 + 9x^2 - x - b$ is exactly divisible by $2x + 3$ is:

- (a) -15
- (b) 15
- (c) 9
- (d) -9

Answer: (b) 15

when $2x^3 + 9x^2 - x - b$ is divided by $2x + 3$, remainder is $-b + 15$

Question 28.

If α and s are two zeros of the polynomial $p(x)$, then which of the following is a factor of $p(x)$:

- (a) $(x - \alpha)(x - s)$
- (b) $(x + \alpha)(x + s)$
- (c) $k(x - \alpha)$
- (d) $k(x - s)$

Answer: (a) $(x - \alpha)(x - s)$

if α, s are the zeros of $p(x)$, then $(x - \alpha)(x - s)$ is a factor of $p(x)$.

Question 29.

Find a cubic polynomial with the sum, sum of the product of its zeros taken two at a time and the product of its zeros as -2 , $+5$, -3 , respectively.

- (a) $2x^3 + 5x^2 + x + 3$
- (b) $4x^3 + 5x^2 - 3x + 7$
- (c) $x^3 + 2x^2 + 5x + 3$
- (d) $2x^3 + 5x^2 + 3x + 1$

Answer: (c) $x^3 + 2x^2 + 5x + 3$

Let the polynomial be $ax^3 + bx^2 + cx + d$

$$-b/a = -2$$

$$c/a = 5$$

$$-d/a = -3$$

$$a = 1, b = 2, c = 5 \text{ and } d = 3$$

required polynomial is $x^3 + 2x^2 + 5x + 3$

Question 30.

Write a polynomial with zeros 1 , -1 and 1 .

- (a) $x^3 + x^2 + x + 1$
- (b) $x^3 - x^2 + x + 1$
- (c) $x^3 - x^2 - x - 1$
- (d) $x^3 - x^2 - x + 1$

Answer: (d) $x^3 - x^2 - x + 1$

zeros are 1 , -1 and 1 .


required polynomial is

$$k(x-1)(x+1)(x-1)$$

$$= x^3 - x^2 - x + 1$$

Question 31.

The graph of a polynomial is as shown, find the polynomial



- (a) $k(x^2 - x - 6)$

- (b) $k(x^3 + x^2 + 6x)$
 (c) $k(x^3 - x^2 - 6x)$
 (d) $k(x^3 - 6x)$

Answer: (c) $k(x^3 - x^2 - 6x)$

zeros are $-2, 0$, and 3

required polynomial $= k(x - 2)(x - 0)(x - 3)$
 $= k(x^3 - x^2 - 6x)$

Question 32.

If α , s and v are the zeroes of the polynomial $2x^3 - x^2 + 3x - 1$, find the value of $\Rightarrow (\alpha s + sv + va + asv)^2$

- (a) $\frac{3}{2}$
 (b) $\frac{5}{2}$
 (c) $\frac{1}{2}$
 (d) 4

Answer: (d) 4

$$\alpha s + sv + v\alpha + \alpha s v = \frac{3}{2} + \frac{1}{2} = 2$$

$$(\alpha s + sv + v\alpha + \alpha s v)^2 = 4$$

Question 33.

If $2 \pm \sqrt{3}$ are the two zeros of a polynomial then the following is a factor:

- (a) $x^2 - 4x + 1$
 (b) $x^2 + 4x - 1$
 (c) $4x^2 + x - 1$
 (d) $4x^2 - x + 1$

Answer: (a) $x^2 - 4x + 1$

If a, s are the zeroes $\Rightarrow (x - \alpha)(x - s)$ is a factor

$\Rightarrow (x - (2 + \sqrt{3}))(x - (2 - \sqrt{3}))$ is a factor

$\Rightarrow x^2 - 4x + 1$ is a factor.

Question 34.

If 2 is a zero of $p(x) = x^2 + 3x + k$, find k :

- (a) 10
 (b) 5
 (c) -3
 (d) -10

Answer: (d) -10

$$p(x) = x^2 + 3x + k$$

$$p(2) = 0$$

$$\Rightarrow 4 + 6 + k = 0$$

$$\Rightarrow k = -10$$

Question 35.

Given that two of the zeroes of the polynomial, $x^3 + px^2 + rx + s$ are 0, then third zero

(a) 0

(b) $\frac{p}{r}$

(c) $\frac{-p}{r}$

(d) $\frac{p}{q}$

Answer: (c) $\frac{-p}{r}$

Two zeroes are zero, let third zero = α

$$\Rightarrow \text{Sum of the roots} = \alpha + 0 + 0$$

$$\frac{\text{Coefficient of } x^2}{\text{Coefficient of } x^3}$$

$$\frac{-p}{1}$$

Question 36.

Given that one of the zeroes of the polynomial $ax^3 + bx^2 + cx + d$ is zero, then the product of the other two zeroes is:

(a) $\frac{-c}{a}$

(b) $\frac{c}{a}$

(c) 0

(d) $\frac{-b}{a}$

Answer: (b) $\frac{c}{a}$

$$\alpha\beta + \beta\gamma + \gamma\alpha = \frac{-c}{a}$$

$$\text{now } \alpha = 0$$

$$0 + \beta\gamma + 0 = \frac{-c}{a}$$

$$\beta\gamma = \frac{-c}{a}$$

Question 37.

The number of polynomials having zeroes -1 and -5 is :

(a) 2

(b) 3



- (c) 1
- (d) More than 3.

Answer: (d) More than 3.

n – number of polynomials can have zeroes -1 and -5.

Question 38.

The graph of the polynomial $f(x) = 2x - 5$ intersects the x – axis at

- (a) $(\frac{5}{2}, 0)$
- (b) $(\frac{-5}{2}, 0)$
- (c) $(\frac{-5}{2}, \frac{5}{2})$
- (d) $(\frac{5}{2}, \frac{-5}{2})$

Answer: (a) $(\frac{5}{2}, 0)$

Question 39.

If the zeroes of the quadratic polynomial $Ax^2 + Bx + C$, $C \neq 0$ are equal, then

- (a) A and B have the same sign
- (b) A and C have the same sign
- (c) B and C have the same sign
- (d) A and C have opposite signs

Answer: (b) A and C have the same sign

Question 40.

The number of polynomials having zeroes as 4 and 7 is

- (a) 2
- (b) 3
- (c) 4
- (d) more than 4

Answer: (d) more than 4

Question 41.

If one of the zeroes of the cubic polynomial $x^3 + ax^2 + bx + c$ is -1, then the product of the other two zeroes is

- (a) $b - a + 1$
- (b) $b - a - 1$

- (c) $a - b + 1$
- (d) $a - b - 1$

Answer: (a) $b - a + 1$

Question 42.

The number of zeros of a cubic polynomial is

- (a) 3
- (b) at least 3
- (c) 2
- (d) at most 3

Answer: (d) at most 3

Question 43.

Find the quadratic polynomial whose zeros are 2 and -6

- (a) $x^2 + 4x + 12$
- (b) $x^2 - 4x - 12$
- (c) $x^2 + 4x - 12$
- (d) $x^2 - 4x + 12$

Answer: (c) $x^2 + 4x - 12$

Question 44.

If 5 is a zero of the quadratic polynomial, $x^2 - kx - 15$ then the value of k is

- (a) 2
- (b) -2
- (c) 4
- (d) -4

Answer: (a) 2

Question 45.

The number of polynomials having zeroes as -2 and 5 is

- (a) 1
- (b) 2
- (c) 3
- (d) more than 3



Answer: (d) more than 3

Question 46.

The zeroes of the quadratic polynomial $x^2 + 1750x + 175000$ are

- (a) both negative
- (b) one positive and one negative
- (c) both positive
- (d) both equal

Answer: (a) both negative

Question 47.

If the zeroes of the quadratic polynomial $x^2 + (a + 1)x + b$ are 2 and -3, then

- (a) $a = -7$, $b = -1$
- (b) $a = 5$, $b = -1$
- (c) $a = 2$, $b = -6$
- (d) $a = 0$, $b = -6$

Answer: (d) $a = 0$, $b = -6$

Question 48.

Sum and the product of zeroes of the polynomial $x^2 + 7x + 10$ is

- (a) $\frac{10}{7}$ and $\frac{-10}{7}$
- (b) $\frac{7}{10}$ and $\frac{-7}{10}$
- (c) -7 and 10
- (d) 7 and -10

Answer: (c) -7 and 10

Question 49.

If $x = 2$ and $x = 3$ are zeros of the quadratic polynomial $x^2 + ax + b$, the values of a and b respectively are :

- (a) 5, 6
- (b) -5, -6
- (c) -5, 6
- (d) 5, -6

Answer: (c) -5, 6

Question 50.

The zeroes of the quadratic polynomial $3x^2 - 48$ are

- (a) both negative
- (b) one positive and one negative
- (c) both positive
- (d) both equal

Answer: (b) one positive and one negative

Question 14.

The zeroes of the quadratic polynomial $x^2 + kx + k$, $k \neq 0$,

- (a) cannot both be positive
- (b) cannot both be negative
- (c) are always unequal
- (d) are always equal

Answer: (a) cannot both be positive

Question 51.

The sum and product of the zeroes of the polynomial $x^2 - 6x + 8$ are respectively

- (a) $-\frac{3}{2}$ and -1
- (b) 6 and 8
- (c) $-\frac{3}{2}$ and 1
- (d) $\frac{3}{2}$ and 1

Answer: (b) 6 and 8

Question 52.

If the point (5,0), (0,-2) and (3,6) lie on the graph of a polynomial. Then which of the following is a zero of the polynomial?

- (a) 5
- (b) 6
- (c) not defined
- (d) -2

Answer: (a) 5

Question 53.

If α and β are the zeroes of the polynomial $5x^2 - 7x + 2$, then sum of their reciprocals is:

- (a) $\frac{14}{25}$
- (b) $\frac{7}{5}$
- (c) $\frac{2}{5}$
- (d) $\frac{7}{2}$

Answer: (d) $\frac{7}{2}$

Question 54.

If one zero of the quadratic polynomial $x^2 + 3x + k$ is 2, then the value of k is

- (a) 10
- (b) -10
- (c) 5
- (d) -5

Answer: (b) -10

Question 55.

The zeroes of the quadratic polynomial $x^2 + px + p$, $p \neq 0$ are

- (a) both equal
- (b) both cannot be positive
- (c) both unequal
- (d) both cannot be negative

Answer: (b) both cannot be positive

Question 56.

The zeroes of the quadratic polynomial $x^2 + 99x + 127$ are

- (a) both positive
- (b) both negative
- (c) one positive and one negative
- (d) both equal

Answer: (b) both negative

Fill in the blanks:

1. A quadratic equation can have _____ two roots, (exactly/atleast/atmost)

Answer: atmost

2. If α is a zero of $p(x)$, then _____ is a factor of $p(x)$.

Answer: $(x - \alpha)$

3. The sum of the zeroes of a cubic polynomial is _____

Answer: $-\frac{\text{coefficient of } x^2}{\text{coefficient of } x^3}$

4. Division Algorithm for polynomials states that, Dividend = _____ \times _____ + Remainder.


Answer: Divisor \times coefficient





5. If a polynomial $p(x)$ does not touch _____ axis, then it has no zeroes.

Answer: X – axis


Match the following:





Question 1.

<p>(a)</p> 	Linear polynomial (one zero)	Touches x axis at one point only -2
	Quadratic Polynomial (2 zeros)	intersects X-axis at $x = 1$.

<p>(b)</p> 		
<p>(c)</p> 	<p>Quadratic Polynomial (no zero)</p>	<p>Does not meet X-axis.</p>
<p>(d)</p> 	<p>Linear Polynomial (One zero)</p>	<p>Passes through origin.</p>
<p>(e)</p> 	<p>Quadratic Polynomial (One zero)</p>	<p>Meets X- axis at 2 points $x = 1$ and $x = -1$</p>


Answer:

<p>(a)</p> 	<p>Quadratic Polynomial (2 zeros)</p>	<p>Meets X- axis at 2 points $x = 1$ and $x = -1$</p>
--	---------------------------------------	---


(b) 	Linear Polynomial (One zero)	intersects X-axis at $x = 1$.
(c) 	Linear Polynomial (One zero)	Passes through origin.
(d) 	Quadratic Polynomial (one zero)	Meets X- axis at -2
(e) 	Quadratic Polynomial (no zero)	Does not meet X-axis.

Question 2.

(a) $p(x) = ax + b$	No. of Zeroes = 3	3 Zeroes	$\alpha + \beta = -\frac{d}{a}$
(b) $q(x) = ax^2 + bx + c$ ($a \neq 0$)	Cubic Polynomial	2 Zeroes	Sum of the zeroes = 0
(c) $r(x) = ax^3 + bx^2 +$	Linear	Meets X-axis at 3	$\alpha + \beta = -\frac{b}{a}$

$cx + d(a \neq 0)$	Polynomial		
(d) 	Quadratic Polynomial	One zero	$-\frac{b}{a}$

Answer:

(a) $p(x) = ax + b$	Linear Polynomial	One zero	$-\frac{b}{a}$
(b) $q(x) = ax^2 + bx + c$ ($a \neq 0$)	Quadratic Polynomial	2 Zeroes	$\alpha + s = -\frac{b}{a}$
(c) $r(x) = ax^3 + bx^2 + cx + d(a \neq 0)$	Cubic Polynomial	3 Zeroes	$\alpha s v = -\frac{d}{a}$
(d) 	No. of Zeroes = 3	Meets X-axis at 3	Sum of the zeroes = 0