

Coordinate Geometry

NCERT TEXTBOOK QUESTIONS SOLVED

EXERCISE 7.1

Q. 1. Find the distance between the following pairs of points:

- (i) $(2, 3), (4, 1)$ (ii) $(-5, 7), (-1, 3)$ (iii) $(a, b), (-a, -b)$

Sol. (i) Here $x_1 = 2, y_1 = 3, x_2 = 4$ and $y_2 = 1$

\therefore The required distance

$$\begin{aligned} &= \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \\ &= \sqrt{(4 - 2)^2 + (1 - 3)^2} \end{aligned}$$

$$\begin{aligned}
 &= \sqrt{2^2 + (-2)^2} \\
 &= \sqrt{4 + 4} = \sqrt{8} \\
 &= \sqrt{2 \times 4} = 2\sqrt{2}
 \end{aligned}$$

(ii) Here, $x_1 = -5, y_1 = 7$
 $x_2 = -1, y_2 = 3$

∴ The required distance

$$\begin{aligned}
 &= \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \\
 &= \sqrt{[-1 - (-5)]^2 + (3 - 7)^2} \\
 &= \sqrt{(-1 + 5)^2 + (-4)^2} \\
 &= \sqrt{16 + 16} \\
 &= \sqrt{32} = \sqrt{2 \times 16} = 4\sqrt{2}
 \end{aligned}$$

(iii) Here, $x_1 = a, y_1 = b$
 $x_2 = -a, y_2 = -b$

∴ The required distance


$$\begin{aligned}
 &= \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \\
 &= \sqrt{(-a - a)^2 + (-b - b)^2} \\
 &= \sqrt{(-2a)^2 + (-2b)^2} \\
 &= \sqrt{4a^2 + 4b^2} \\
 &= \sqrt{4(a^2 + b^2)} = 2\sqrt{a^2 + b^2}
 \end{aligned}$$

Q. 2. Find the distance between the points (0, 0) and (36, 15). Can you now find the distance between the two towns A and B discussed in Section 7.2 of the NCERT textbook?

Sol. Part-I

Let the points be P (0, 0) and Q (36, 15).

$$\begin{aligned}
 \therefore PQ &= \sqrt{(36 - 0)^2 + (15 - 0)^2} \\
 &= \sqrt{(36)^2 + (15)^2} \\
 &= \sqrt{1296 + 225} \\
 &= \sqrt{1521} \\
 &= \sqrt{39^2} = 39
 \end{aligned}$$



Part-II

We have P (0, 0) and Q (36, 15) as the positions of two towns.

$$\begin{aligned}
 \therefore \text{Here } x_1 &= 0, x_2 = 36 \\
 y_1 &= 0, y_2 = 15
 \end{aligned}$$

$$\begin{aligned}\therefore PQ &= \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \\ &= \sqrt{(36 - 0)^2 + (15 - 0)^2} = \mathbf{39 \text{ km.}}\end{aligned}$$

Q. 3. Determine if the points (1, 5), (2, 3) and (-2, -11) are collinear.

Sol. Let the points be A (1, 5), B (2, 3) and C (-2, -11)

A, B and C are collinear, if

$$AB + BC = AC$$

$$AC + CB = AB$$

$$BA + AC = BC$$

$$\begin{aligned}\therefore AB &= \sqrt{(2 - 1)^2 + (3 - 5)^2} \\ &= \sqrt{1^2 + (-2)^2} \\ &= \sqrt{1 + 4} = \sqrt{5} \\ BC &= \sqrt{(-2 - 2)^2 + (-11 - 3)^2} \\ &= \sqrt{(-4)^2 + (-14)^2} \\ &= \sqrt{16 + 196} = \sqrt{212} \\ AC &= \sqrt{(-2 - 1)^2 + (-11 - 5)^2} \\ &= \sqrt{(-3)^2 + (-16)^2} \\ &= \sqrt{9 + 256} = \sqrt{265}\end{aligned}$$

But $AB + BC \neq AC$

$$AC + CB \neq AB$$

$$BA + AC \neq BC$$

\therefore A, B and C are **not collinear**.

Q. 4. Check whether (5, -2), (6, 4) and (7, -2) are the vertices of an isosceles triangle.

Sol. Let the points be A (5, -2), B (6, 4) and C (7, -2).

(CBSE 2012)


$$\begin{aligned}\therefore AB &= \sqrt{(6 - 5)^2 + [4 - (-2)]^2} \\ &= \sqrt{(1)^2 + (6)^2} \\ &= \sqrt{1 + 36} = \sqrt{37} \\ BC &= \sqrt{(7 - 6)^2 + (-2 - 4)^2} \\ &= \sqrt{(1)^2 + (-6)^2} \\ &= \sqrt{1 + 36} = \sqrt{37} \\ AC &= \sqrt{(5 - 7)^2 + (-2 - (-2))^2} \\ &= \sqrt{(-2)^2 + (0)^2}\end{aligned}$$

$$= \sqrt{4+0} = 2$$

We have $AB = BC \neq AC$

$\therefore \triangle ABC$ is an **isosceles triangle**.

- Q. 5.** In a classroom, 4 friends are seated at the points A, B, C and D as shown in Fig. Champa and Chameli walk into the class and after observing for a few minutes Champa asks Chameli, "Don't you think ABCD is a square?" Chameli disagrees. Using distance single formula, find which of them is correct.



Sol. Let the number of horizontal columns represent the x -coordinates whereas the vertical rows represent the y -coordinates.

\therefore The points are:

A (3, 4), B (6, 7), C (9, 4) and D (6, 1)

$$\begin{aligned} \therefore AB &= \sqrt{(6-3)^2 + (7-4)^2} \\ &= \sqrt{(3)^2 + (3)^2} \\ &= \sqrt{9+9} = \sqrt{18} = 3\sqrt{2} \end{aligned}$$

$$\begin{aligned} BC &= \sqrt{(9-6)^2 + (4-7)^2} \\ &= \sqrt{3^2 + (-3)^2} \\ &= \sqrt{9+9} = \sqrt{18} = 3\sqrt{2} \end{aligned}$$

$$\begin{aligned} CD &= \sqrt{(6-9)^2 + (1-4)^2} \\ &= \sqrt{(-3)^2 + (-3)^2} \\ &= \sqrt{9+9} = \sqrt{18} = 3\sqrt{2} \end{aligned}$$

$$\begin{aligned} AD &= \sqrt{(6-3)^2 + (1-4)^2} \\ &= \sqrt{(3)^2 + (-3)^2} \end{aligned}$$



$$= \sqrt{9+9} = \sqrt{18} = 3\sqrt{2}$$

Since, $AB = BC = CD = AD$

i.e., All the four sides are equal.

$$\begin{aligned}\text{Also } AC &= \sqrt{(9-3)^2 + (4-4)^2} \\ &= \sqrt{(-6)^2 + (0)^2} = 6\end{aligned}$$

$$\begin{aligned}\text{and } BD &= \sqrt{(6-6)^2 + (1-7)^2} \\ &= \sqrt{(0)^2 + (-6)^2} = 6\end{aligned}$$

i.e., $BD = AC \Rightarrow$ Both the diagonals are also equal.

$\therefore ABCD$ is a square.

Thus, Champa is correct.

Q. 6. Name the type of quadrilateral formed, if any, by the following points, and give reasons for your answer:

(i) $(-1, -2), (1, 0), (-1, 2), (-3, 0)$

(ii) $(-3, 5), (3, 1), (0, 3), (-1, -4)$

(iii) $(4, 5), (7, 6), (4, 3), (1, 2)$

Sol. (i) Let the points be: $A(-1, -2), B(1, 0), C(-1, 2)$ and $D(-3, 0)$.

$$\begin{aligned}\therefore AB &= \sqrt{(1+1)^2 + (0+2)^2} \\ &= \sqrt{(2)^2 + (2)^2} = \sqrt{4+4} = \sqrt{8} \\ BC &= \sqrt{(-1-1)^2 + (2-0)^2} = \sqrt{4+4} = \sqrt{8} \\ CD &= \sqrt{(-3+1)^2 + (0-2)^2} = \sqrt{4+4} = \sqrt{8} \\ DA &= \sqrt{(-1+3)^2 + (-2-0)^2} = \sqrt{4+4} = \sqrt{8} \\ AC &= \sqrt{(-1+1)^2 + (2+2)^2} = \sqrt{0+4^2} = 4 \\ BD &= \sqrt{(-3-1)^2 + (0-0)^2} = \sqrt{(4)^2} = 4\end{aligned}$$

$$\Rightarrow AB = BC = CD = AD$$

i.e., All the sides are equal.

And $AC = BD$

Also, AC and BD (the diagonals) are equal.

$\therefore ABCD$ is a square.

(ii) Let the points be $A(-3, 5), B(3, 1), C(0, 3)$ and $D(-1, -4)$.

$$\begin{aligned}\therefore AB &= \sqrt{[3-(-3)]^2 + (1-5)^2} \\ &= \sqrt{6^2 + (-4)^2} \\ &= \sqrt{36+16} = \sqrt{52} = 2\sqrt{13} \\ BC &= \sqrt{(0-3)^2 + (3-1)^2} \\ &= \sqrt{9+4} = \sqrt{13}\end{aligned}$$

$$CD = \sqrt{(-1-0)^2 + (-4-3)^2}$$

$$= \sqrt{(-1)^2 + (-7)^2} = \sqrt{1+49} = \sqrt{50}$$

$$DA = \sqrt{[-3-(-1)]^2 + [5-(-4)]^2}$$

$$= \sqrt{(2)^2 + (9)^2}$$

$$= \sqrt{4+81} = \sqrt{85}$$

$$AC = \sqrt{[0-(-3)]^2 + (3-5)^2}$$

$$= \sqrt{(3)^2 + (-2)^2}$$

$$= \sqrt{9+4} = \sqrt{13}$$

$$BD = \sqrt{(-1-3)^2 + (-4-1)^2} = \sqrt{(-4)^2 + (-5)^2}$$

$$= \sqrt{16+25} = \sqrt{41}$$

We see that:

$$\sqrt{13} + \sqrt{13} = 2\sqrt{13}$$

$$\text{i.e., } AC + BC = AB$$

\Rightarrow A, B, C and D are collinear. Thus, ABCD is **not a quadrilateral**.

(iii) Let the points be A (4, 5), B (7, 6), C (4, 3) and D (1, 2).

$$\therefore AB = \sqrt{(7-4)^2 + (6-5)^2}$$

$$= \sqrt{3^2 + 1^2} = \sqrt{10}$$

$$BC = \sqrt{(4-7)^2 + (3-6)^2}$$

$$= \sqrt{(-3)^2 + (-3)^2} = \sqrt{18}$$

$$CD = \sqrt{(1-4)^2 + (2-3)^2}$$

$$= \sqrt{(-3)^2 + (-1)^2} = \sqrt{10}$$

$$DA = \sqrt{(4-1)^2 + (5-2)^2}$$

$$= \sqrt{9+9} = \sqrt{18}$$

$$AC = \sqrt{(4-4)^2 + (3-5)^2}$$

$$= \sqrt{0+(-2)^2} = 2$$

$$BD = \sqrt{(1-7)^2 + (2-6)^2}$$

$$= \sqrt{36+16} = \sqrt{52}$$



Since, $AB = CD$ [opposite sides of the quadrilateral are equal]
 $BC = DA$

And $AC \neq BD \Rightarrow$ Diagonals are unequal

$\therefore ABCD$ is a **parallelogram**.

Q. 7. Find the point on the x -axis which is equidistant from $(2, -5)$ and $(-2, 9)$. (CBSE 2012)

Sol. We know that any point on x -axis has its ordinate = 0.

Let the required point be $P(x, 0)$.

Let the given points be $A(2, -5)$ and $B(-2, 9)$.

$$\begin{aligned}\therefore PA &= \sqrt{(x-2)^2 + [0-(-5)]^2} \\ &= \sqrt{(x-2)^2 + 5^2} = \sqrt{x^2 - 4x + 4 + 25} = \sqrt{x^2 - 4x + 29} \\ PB &= \sqrt{[x-(-2)]^2 + (0-9)^2} \\ &= \sqrt{(x+2)^2 + (-9)^2} = \sqrt{x^2 + 4x + 4 + 81} = \sqrt{x^2 + 4x + 85}\end{aligned}$$

Since, A and B are equidistant from P ,

$$\therefore PA = PB$$

$$\Rightarrow \sqrt{x^2 - 4x + 29} = \sqrt{x^2 + 4x + 85}$$

$$\Rightarrow x^2 - 4x + 29 = x^2 + 4x + 85$$

$$\Rightarrow x^2 - 4x - x^2 - 4x = 85 - 29$$

$$\Rightarrow -8x = 56$$

$$\Rightarrow x = \frac{56}{-8} = -7$$

\therefore The required point is **$(-7, 0)$** .

Q. 8. Find the values of y for which the distance between the points $P(2, -3)$ and $Q(10, y)$ is 10 units.

Sol. The given points are $P(2, -3)$ and $Q(10, y)$.

$$\begin{aligned}\therefore PQ &= \sqrt{(10-2)^2 + [y-(-3)]^2} \\ &= \sqrt{8^2 + (y+3)^2} \\ &= \sqrt{64 + y^2 + 6y + 9} \\ &= \sqrt{y^2 + 6y + 73}\end{aligned}$$

But $PQ = 10$

$$\therefore \sqrt{y^2 + 6y + 73} = 10$$

Squaring both sides,

$$y^2 + 6y + 73 = 100$$

$$\Rightarrow y^2 + 6y - 27 = 0$$

$$\Rightarrow y^2 - 3y + 9y - 27 = 0$$

$$\Rightarrow (y-3)(y+9) = 0$$

$$\Rightarrow \text{Either } y-3 = 0 \Rightarrow y = 3$$

$$\text{or } y+9 = 0 \Rightarrow y = -9$$

\therefore The required value of y is **3 or -9**.

Q. 9. If $Q(0, 1)$ is equidistant from $P(5, -3)$ and $R(x, 6)$, find the values of x . Also find the distances QR and PR .

Sol. Here,

$$QP = \sqrt{(5-0)^2 + [(-3)-1]^2}$$

$$= \sqrt{5^2 + (-4)^2}$$

$$= \sqrt{25+16} = \sqrt{41}$$

$$QR = \sqrt{(x-0)^2 + (6-1)^2}$$

$$= \sqrt{x^2 + 5^2} = \sqrt{x^2 + 25}$$

$$\therefore QP = QR$$

$$\therefore \sqrt{41} = \sqrt{x^2 + 25}$$

Squaring both sides, we have:

$$x^2 + 25 = 41$$

$$\Rightarrow x^2 + 25 - 41 = 0$$

$$\Rightarrow x^2 - 16 = 0 \Rightarrow x = \pm 4$$

Thus, the point R is **(4, 6) or (-4, 6)**

Now,

$$QR = \sqrt{[(\pm 4) - (0)]^2 + (6-1)^2}$$

$$= \sqrt{16+25} = \sqrt{41}$$

and

$$PR = \sqrt{(\pm 4 - 5)^2 + (6+3)^2}$$

$$\Rightarrow PR = \sqrt{(4-5)^2 + (6+3)^2} \text{ or } \sqrt{(-4-5)^2 + (6+3)^2}$$

$$\Rightarrow PR = \sqrt{1+81} \text{ or } \sqrt{(-9)^2 + 9^2}$$

$$\Rightarrow PR = \sqrt{82} \text{ or } \sqrt{2 \times 9^2}$$

$$\Rightarrow PR = \sqrt{82} \text{ or } 9\sqrt{2}$$

Q. 10. Find a relation between x and y such that the point (x, y) is equidistant from the point $(3, 6)$ and $(-3, 4)$.

Sol. Let the points be $A(x, y)$, $B(3, 6)$ and $C(-3, 4)$.

$$\therefore AB = \sqrt{(3-x)^2 + (6-y)^2}$$

And

$$AC = \sqrt{[(-3)-x]^2 + (4-y)^2}$$

Since, the point (x, y) is equidistant from $(3, 6)$ and $(-3, 4)$.

$$\therefore AB = AC$$

$$\Rightarrow \sqrt{(3-x)^2 + (6-y)^2} = \sqrt{(-3-x)^2 + (4-y)^2}$$

Squaring both sides,

$$(3-x)^2 + (6-y)^2 = (-3-x)^2 + (4-y)^2$$

$$\Rightarrow (9 + x^2 - 6x) + (36 + y^2 - 12y) = (9 + x^2 + 6x) + (16 + y^2 - 8y)$$

$$\Rightarrow 9 + x^2 - 6x + 36 + y^2 - 12y - 9 - x^2 - 6x - 16 - y^2 + 8y = 0$$

$$\Rightarrow -6x - 6x + 36 - 12y - 16 + 8y = 0$$

$$\Rightarrow -12x - 4y + 20 = 0$$

$$\Rightarrow -3x - y + 5 = 0$$

$$\Rightarrow 3x + y - 5 = 0$$

[Dividing by 4]

which is the required relation between x and y .

NCERT TEXTBOOK QUESTIONS SOLVED

EXERCISE 7.2

Q. 1. Find the coordinates of the point which divides the join of $(-1, 7)$ and $(4, -3)$ in the ratio $2 : 3$.

Sol. Let the required point be $P(x, y)$.

Here, the end points are:

$(-1, 7)$ and $(4, -3)$

\therefore Ratio $= 2 : 3 = m_1 : m_2$

$$\begin{aligned} \therefore x &= \frac{m_1 x_2 + m_2 x_1}{m_1 + m_2} \\ &= \frac{(2 \times 4) + 3 \times (-1)}{2 + 3} \\ &= \frac{8 - 3}{5} = \frac{5}{5} = 1 \end{aligned}$$


And

$$\begin{aligned} y &= \frac{m_1 y_2 + m_2 y_1}{m_1 + m_2} \\ &= \frac{2 \times (-3) + (3 \times 7)}{2 + 3} \\ &= \frac{-6 + 21}{5} = \frac{15}{5} = 3 \end{aligned}$$

Thus, the required point is **$(1, 3)$** .

Q. 2. Find the coordinates of the points of trisection of the line segment joining $(4, -1)$ and $(-2, -3)$.

Sol. Let the given points be $A(4, -1)$ and $B(-2, -3)$.



Let the points P and Q trisect AB .

i.e., $AP = PQ = QB$

i.e., P divides AB in the ratio of $1 : 2$

Q divides AB in the ratio of $2 : 1$

Let the coordinates of P be (x, y) .

$$\begin{aligned} \therefore x &= \frac{m_1 x_2 + m_2 x_1}{m_1 + m_2} \\ &= \frac{1(-2) + 2(4)}{1 + 2} = \frac{-2 + 8}{3} = 2 \\ y &= \frac{m_1 y_2 + m_2 y_1}{m_1 + m_2} = 1 \end{aligned}$$



$$= \frac{1(-3) + 2 \times (-1)}{1+2} = \frac{-3-2}{3} = \frac{-5}{3}$$

∴ The required co-ordinates of P are $\left(2, \frac{-5}{3}\right)$

Let the co-ordinates of Q be (X, Y).

$$\therefore X = \frac{m_1 x_2 + m_2 x_1}{m_1 + m_2} = \frac{2(-2) + 1(4)}{2+1} = \frac{-4+4}{3} = 0$$

$$Y = \frac{m_1 y_2 + m_2 y_1}{m_1 + m_2} = \frac{2(-3) + 1(-1)}{2+1} = \frac{-6-1}{3} = \frac{-7}{3}$$

∴ The required coordinates of Q are $\left(0, \frac{-7}{3}\right)$.


Q. 3. To conduct Sports Day activities, in your rectangular shaped school ground ABCD, lines have been drawn with chalk powder at a distance of 1 m each. 100 flower pots have been placed at a distance of 1 m from each other along AD, as shown in the figure. Niharika runs $\frac{1}{4}$ th the distance

AD on the 2nd line and posts a green flag. Preet runs $\frac{1}{5}$ th the distance AD on the eighth line and posts a red flag. What is the distance between both the flags? If Rashmi has to post a blue flag exactly halfway between the line segment joining the two flags, where should she post her flag?

Sol. Let us consider 'A' as origin, then AB is the x-axis. AD is the y-axis.

Now, the position of green flag-post is $\left(2, \frac{100}{4}\right)$ or (2, 25)

And the position of red flag-post is $\left(8, \frac{100}{5}\right)$ or (8, 20)




⇒ Distance between both the flags

$$= \sqrt{(8-2)^2 + (20-25)^2}$$

$$= \sqrt{6^2 + (-5)^2} = \sqrt{36+25} = \sqrt{61}$$

Let the mid-point of the line segment joining the two flags be $M(x, y)$.



$$\therefore x = \frac{2+8}{2} \quad \text{and} \quad y = \frac{25+20}{2}$$

or $x = 5$ and $y = 22.5$.

Thus, the blue flag is on the 5th line at a distance **22.5 m** above AB.

Q. 4. Find the ratio in which the line segment joining the points $(-3, 10)$ and $(6, -8)$ is divided by $(-1, 6)$.

Sol. Let the given points are: A $(-3, 10)$ and B $(6, -8)$.

Let the point P $(-1, 6)$ divides AB in the ratio $m_1 : m_2$.

∴ Using the section formula, we have:

$$(-1, 6) = \left(\frac{x_2 m_1 + m_2 x_1}{m_1 + m_2}, \frac{m_1 y_2 + m_2 y_1}{m_1 + m_2} \right)$$

$$\Rightarrow (-1, 6) = \left(\frac{(m_1 \times 6) + [m_2 \times (-3)]}{m_1 + m_2}, \frac{[m_1 (-8)] + (m_2 \times 10)}{m_1 + m_2} \right)$$

$$\Rightarrow (-1, 6) = \frac{6m_1 + (-3m_2)}{m_1 + m_2}, \frac{-8m_1 + 10m_2}{m_1 + m_2}$$

$$\Rightarrow -1 = \frac{6m_1 - 3m_2}{m_1 + m_2} \quad \text{and} \quad 6 = \frac{-8m_1 + 10m_2}{m_1 + m_2}$$

$$\Rightarrow -1(m_1 + m_2) = 6m_1 - 3m_2 \quad \text{and} \quad 6(m_1 + m_2) = -8m_1 + 10m_2$$

$$\Rightarrow -m_1 - m_2 - 6m_1 + 3m_2 = 0 \quad \text{and} \quad 6m_1 + 6m_2 + 8m_1 - 10m_2 = 0$$

$$\Rightarrow -7m_1 + 2m_2 = 0 \quad \text{and} \quad 14m_1 - 4m_2 = 0 \quad \text{or} \quad 7m_1 - 2m_2 = 0$$

$$\Rightarrow 2m_2 = 7m_1 \quad \text{and} \quad 7m_1 = 2m_2$$

$$\Rightarrow \frac{m_1}{m_2} = \frac{2}{7} \quad \text{and} \quad \frac{m_1}{m_2} = \frac{2}{7}$$

$$\Rightarrow m_1 : m_2 = 2 : 7 \quad \text{and} \quad m_1 : m_2 = 2 : 7$$

Thus, the required ratio is **2 : 7**.

Q. 5. Find the ratio in which the line segment joining A $(1, -5)$ and B $(-4, 5)$ is divided by the x-axis. Also find the coordinates of the point of division.

Sol. The given points are: A $(1, -5)$ and B $(-4, 5)$.

Let the required ratio = $k : 1$ and the required point be P (x, y) .

Part-I: To find the ratio

Since the point P lies on x-axis,

∴ Its y-coordinate is 0.

$$x = \frac{m_1 x_2 + m_2 x_1}{m_1 + m_2} \quad \text{and} \quad 0 = \frac{m_1 y_2 + m_2 y_1}{m_1 + m_2}$$

$$\Rightarrow x = \frac{k(-4) + 1(1)}{k+1} \quad \text{and} \quad 0 = \frac{k(5) + 1(-5)}{k+1}$$

$$\Rightarrow x = \frac{-4k+1}{k+1} \quad \text{and} \quad 0 = \frac{5k-5}{k+1}$$

$$\Rightarrow x(k+1) = -4k+1 \quad \text{and} \quad 5k-5=0 \Rightarrow k=1$$

Part II : To find coordinates of P :

$$\Rightarrow x(k+1) = -4k+1 \Rightarrow x(1+1) = -4+1 \quad [\because k=1]$$

$$\Rightarrow 2x = -3$$

$$\Rightarrow x = \frac{-3}{2}$$

\therefore The required ratio $k : 1 = 1 : 1$

Coordinates of P are $(x, 0) = \left(\frac{-3}{2}, 0\right)$.

Q. 6. If $(1, 2)$, $(4, y)$, $(x, 6)$ and $(3, 5)$ are the vertices of a parallelogram taken in order, find x and y .

Sol. We have the parallelogram vertices

A $(1, 2)$, B $(4, y)$, C $(x, 6)$ and D $(3, 5)$.

Since, the diagonals of a parallelogram bisect each other.

\therefore The coordinates of P are:

$$X = \frac{x+1}{2} = \frac{3+4}{2}$$


$$\Rightarrow x+1 = 7 \Rightarrow x = 6$$

$$Y = \frac{5+y}{2} = \frac{6+2}{2}$$

$$\Rightarrow 5+y = 8 \Rightarrow y = 3$$

\therefore The required values of x and y are:

$$x = 6, \quad y = 3$$




Q. 7. Find the coordinates of a point A, where AB is the diameter of a circle whose centre is $(2, -3)$ and B is $(1, 4)$.

Sol. Here, centre of the circle is O $(2, -3)$.

Let the end points of the diameter be

A (X, Y) and B $(1, 4)$



The centre of a circle bisects the diameter.

$$\therefore 2 = \frac{x+1}{2} \Rightarrow x+1 = 4 \quad \text{or} \quad x = 3$$

$$\text{And} \quad -3 = \frac{y+4}{2} \Rightarrow y+4 = -6 \quad \text{or} \quad y = -10$$


Hence the coordinates of A are $(3, -10)$.

Q. 8. If A and B are $(-2, -2)$ and $(2, -4)$, respectively, find the coordinates of P such that

$AP = \frac{3}{7} AB$ and P lies on the line segment AB.



Sol.



Here, the given points are

$A(-2, -2)$ and $B(2, -4)$

Let the coordinates of P are (x, y) .

Since, the point P divides AB such that

$$AP = \frac{3}{7} AB \quad \text{or} \quad \frac{AP}{AB} = \frac{3}{7}$$

\Rightarrow Since $AB = AP + BP$

$$\therefore \frac{AP}{AB} = \frac{3}{7} \Rightarrow \frac{AP}{AP + AB} = \frac{3}{7}$$

$$\Rightarrow \frac{AP + BP}{AP} = \frac{7}{3}$$

$$\Rightarrow 1 + \frac{BP}{AP} = \frac{3 + 4}{3} = 1 + \frac{4}{3}$$

$$\Rightarrow \frac{BP}{AP} = \frac{4}{3}$$

$$\Rightarrow AP : PB = 3 : 4$$

i.e., $P(x, y)$ divides AB in the ratio $3 : 4$.

$$\therefore x = \frac{3 \times 2 + 4 \times (-2)}{3 + 4} = \frac{6 - 8}{7} = \frac{-2}{7}$$

$$y = \frac{3 \times (-4) + 4 \times (-2)}{3 + 4} = \frac{-12 - 8}{7} = \frac{-20}{7}$$


Thus, the coordinates of P are $\left(\frac{-2}{7}, \frac{-20}{7}\right)$.

Q. 9. Find the coordinates of the points which divide the line segment joining $A(-2, 2)$ and $B(2, 8)$ into four equal parts.

Sol. Here, the given points are:

$A(-2, 2)$ and $B(2, 8)$

Let P_1, P_2 and P_3 divide AB in four equal parts.



$$\therefore AP_1 = P_1P_2 = P_2P_3 = P_3B$$

Obviously, P_2 is the mid point of AB

\therefore Coordinates of P_2 are:

$$\left(\frac{-2+2}{2}, \frac{2+8}{2}\right) \text{ or } (0, 5)$$

Again, P_1 is the mid point of AP_2 .

\therefore Coordinates of P_1 are:

$$\left(\frac{-2+0}{2}, \frac{2+5}{2}\right) \text{ or } \left(-1, \frac{7}{2}\right)$$

Also P_3 is the mid point of $P_2 B$.

\therefore Coordinates of P_3 are:

$$\left(\frac{0+2}{2}, \frac{5+8}{2}\right) \text{ or } \left(1, \frac{13}{2}\right)$$

Thus, the coordinates of P_1 , P_2 and P_3 are:


$$(0, 5), \left(-1, \frac{7}{2}\right) \text{ and } \left(1, \frac{13}{2}\right) \text{ respectively.}$$

Q. 10. Find the area of a rhombus if its vertices are $(3, 0)$, $(4, 5)$, $(-1, 4)$ and $(-2, -1)$ taken in order.

[Hint: Area of a rhombus = $\frac{1}{2}$ (product of its diagonals)]

Sol. Let the vertices of the given rhombus are:

$A (3, 0)$, $B (4, 5)$, $C (-1, 4)$ and $D (-2, -1)$



$\therefore AC$ and BD are the diagonals of rhombus $ABCD$.

$$\begin{aligned} \therefore \text{Diagonal } AC &= \sqrt{(-1-3)^2 + (4-0)^2} = \sqrt{(-4)^2 + (4)^2} \\ &= \sqrt{16+16} = 4\sqrt{2} \end{aligned}$$

$$\begin{aligned} \text{Diagonal } BD &= \sqrt{(-2-4)^2 + (-1-5)^2} \\ &= \sqrt{(-6)^2 + (-6)^2} = \sqrt{36+36} = 2\sqrt{2} \end{aligned}$$

\therefore For a rhombus,

$$\begin{aligned} \text{Area} &= \frac{1}{2} \times (\text{Product of diagonals}) \\ &= \frac{1}{2} \times AC \times BD \\ &= \frac{1}{2} \times 4\sqrt{2} \times 2\sqrt{2} \\ &= \frac{1}{2} \times 2 \times 4 \times 2 \text{ Square units.} \\ &= 4 \times 2 = 8 \text{ Square units.} \end{aligned}$$

● **Area of Triangle**

I. If $A (x_1, y_1)$; $B (x_2, y_2)$ and $C (x_3, y_3)$ are the vertices of ΔABC , then

$$\text{the area of } \Delta ABC = \frac{1}{2} [x_1 (y_2 - y_3) + x_2 (y_3 - y_1) + x_3 (y_1 - y_2)].$$

II. The three points A , B and C are collinear if and only if area of $\Delta ABC = 0$.

NCERT TEXTBOOK QUESTIONS SOLVED

EXERCISE 7.3

Q. 1. Find the area of the triangle whose vertices are:

- (i) (2, 3), (-1, 0), (2, -4) (ii) (-5, -1), (3, -5), (5, 2)

Sol. (i) Let the vertices of the triangle be

A (2, 3), B (-1, 0) and C (2, -4)

Here, $x_1 = 2, y_1 = 3$
 $x_2 = -1, y_2 = 0$
 $x_3 = 2, y_3 = -4$

$$\therefore \text{Area of a } \Delta = \frac{1}{2} [x_1 (y_2 - y_3) + x_2 (y_3 - y_1) + x_3 (y_1 - y_2)]$$

$$\begin{aligned}\therefore \text{Area of } \Delta ABC &= \frac{1}{2} [2 \{0 - (-4)\} + (-1) \{-4 - (3)\} + 2 \{3 - 0\}] \\ &= \frac{1}{2} [2 (0 + 4) + (-1) (-4 - 3) + 2 (3)] \\ &= \frac{1}{2} [8 + 7 + 6] \\ &= \frac{1}{2} [21] = \frac{21}{2} \text{ sq. units.}\end{aligned}$$

(ii) Let the vertices of the triangle be

A (-5, -1), B (3, -5) and C (5, 2)

i.e., $x_1 = -5, y_1 = -1$
 $x_2 = 3, y_2 = -5$
 $x_3 = 5, y_3 = 2$

$$\therefore \text{Area of a } \Delta = \frac{1}{2} [x_1 (y_2 - y_3) + x_2 (y_3 - y_1) + x_3 (y_1 - y_2)]$$

$$\begin{aligned}\therefore \text{Area of } \Delta ABC &= \frac{1}{2} [-5 \{-5 - 2\} + 3 \{2 - (-1)\} + 5 \{-1 - (-5)\}] \\ &= \frac{1}{2} [-5 \{-7\} + 3 \{2 + 1\} + 5 \{-1 + 5\}] \\ &= \frac{1}{2} [(-5) (-7) + 3 (3) + 5 (4)] \\ &= \frac{1}{2} [35 + 9 + 20] \\ &= \frac{1}{2} \times 64 = 32 \text{ sq. units.}\end{aligned}$$

Q. 2. In each of the following find the value of 'k', for which the points are collinear.

- (i) (7, -2), (5, 1), (3, k) (ii) (8, 1), (k, -4), (2, -5)

Sol. The given three points will be collinear if the Δ formed by them has zero area.

(i) Let A (7, -2), B (5, 1) and C (3, k) be the vertices of a triangle.

\therefore The given points will be collinear, if ar (ΔABC) = 0

or $7 (1 - k) + 5 (k + 2) + 3 (-2 - 1) = 0$



$$\Rightarrow 7 - 7k + 5k + 10 + (-6) - 3 = 0$$

$$\Rightarrow 17 - 9 + 5k - 7k = 0$$

$$\Rightarrow 8 - 2k = 0$$

$$\Rightarrow 2k = 8$$

$$\Rightarrow k = \frac{8}{2} = 4$$

The required value of $k = 4$.

(ii) Let $(8, 1)$, $(k, -4)$ and $(2, -5)$ be the vertices of a triangle.

\therefore For the above points being collinear, $\Delta ABC = 0$

$$\text{i.e., } 8(-4 + 5) + k(-5 - 1) + 2[1 - (-4)] = 0$$

$$\Rightarrow 8(+1) + k(-6) + 2(5) = 0$$

$$\Rightarrow 8 + (-6k) + 10 = 0$$

$$\Rightarrow -6k + 18 = 0$$

$$\Rightarrow k = (-18) \div (-6) = 3$$

Thus, $k = 3$.

Q. 3. Find the area of the triangle formed by joining the mid-points of the sides of the triangle whose vertices are $(0, -1)$, $(2, 1)$ and $(0, 3)$. Find the ratio of this area to the area of the given triangle.


Sol. Let the vertices of the triangle be $A(0, -1)$, $B(2, 1)$ and $C(0, 3)$.

Let D , E and F be the mid-points of the sides BC , CA and AB respectively. Then:

$$\text{Coordinates of } D \text{ are } \left(\frac{2+0}{2}, \frac{1+3}{2}\right) \text{ i.e., } \left(\frac{2}{2}, \frac{4}{2}\right) \text{ or } (1, 2)$$

$$\text{Coordinates of } E \text{ are: } \frac{0+0}{2}, \frac{3+(-1)}{2} \text{ i.e., } (0, 1)$$

$$\text{Coordinates of } F \text{ are: } \frac{2+0}{2}, \frac{1+(-1)}{2} \text{ i.e., } (1, 0)$$



$$\text{Now, ar } (\Delta ABC) = \frac{1}{2} [0(1 - 3) + 2\{3 - (-1)\} + 0(-1 - 1)]$$

$$= \frac{1}{2} [0(-2) + 8 + 0(-2)]$$

$$= \frac{1}{2} [0 + 8 + 0] = \frac{1}{2} \times 8 = 4 \text{ sq. units}$$

$$\text{And ar } \Delta (DEF) = \frac{1}{2} [1(1 - 0) + 0(0 - 2) + 1(2 - 1)]$$

$$= \frac{1}{2} [1(1) + 0 + 1(1)]$$


$$= \frac{1}{2} [1 + 0 + 1] = \frac{1}{2} \times 2 = 1 \text{ sq. unit}$$

$$\therefore \frac{\text{ar}(\Delta DEF)}{\text{ar}(\Delta ABC)} = \frac{1}{4}$$

$$\Rightarrow \text{ar}(\Delta DEF) : \text{ar}(\Delta ABC) = 1 : 4.$$

Q. 4. Find the area of the quadrilateral whose vertices, taken in order, are $(-4, -2)$, $(-3, -5)$, $(3, -2)$ and $(2, 3)$.

Sol. Let $A(-4, -2)$, $B(-3, -5)$, $C(3, -2)$ and $D(2, 3)$ be the vertices of the quadrilateral. Let us join diagonal BD .



$$\begin{aligned} \text{Now, ar}(\Delta ABD) &= \frac{1}{2} [(-4)\{-5-3\} + (-3)\{3-(-2)\} + 2\{(-2)-(-5)\}] \\ &= \frac{1}{2} [(-4)(-8) + (-3)(5) + 2(-2+5)] \\ &= \frac{1}{2} [32 + (-15) + 6] \\ &= \frac{1}{2} [23] = \frac{23}{2} \text{ sq. units} \end{aligned}$$

$$\begin{aligned} \text{ar}(\Delta CBD) &= \frac{1}{2} [3\{-5-3\} + (-3)\{3-(-2)\} + 2\{(-2)-(-5)\}] \\ &= \frac{1}{2} [3(-8) + -3(5) + 2(3)] \\ &= \frac{1}{2} [-24 - 15 + 6] \\ &= \frac{1}{2} [-33] = \frac{33}{2} \text{ sq. units, (numerically)} \end{aligned}$$

$$\text{Since, ar (quad } ABCD) = \text{ar}(\Delta ABD) + \text{ar}(\Delta CBD)$$

$$\begin{aligned} \therefore \text{ar (quad } ABCD) &= \left(\frac{23}{2} + \frac{33}{2}\right) \text{ sq. units} \\ &= \frac{56}{2} \text{ sq. units} = 28 \text{ sq. units.} \end{aligned}$$

Q. 5. You have studied in class IX (Chapter 9, Example-3) that, a median of a triangle divides it into two triangles of equal areas. Verify this result for ΔABC whose vertices are $A(4, -6)$, $B(3, -2)$ and $C(5, 2)$.


Sol. Here, the vertices of the triangle are $A(4, -6)$, $B(3, -2)$ and $C(5, 2)$. Let D be the mid-point of BC .

∴ The coordinates of the mid point D are:

$$\left\{ \frac{3+5}{2}, \frac{-2+2}{2} \right\} \text{ or } (4, 0).$$

Since AD divides the triangle ABC into two parts i.e., ΔABD and ΔACD ,

$$\begin{aligned} \text{Now, ar } (\Delta ABD) &= \frac{1}{2} [4 \{(-2) - 0\} + 3 (0 + 6) + 4 (-6 + 2)] \\ &= \frac{1}{2} [(-8) + 18 + (-16)] \\ &= \frac{1}{2} (-6) = -3 \text{ sq. units.} \\ &= 3 \text{ sq. units (numerically)} \end{aligned} \quad \dots(1)$$



$$\begin{aligned} \text{ar } (\Delta ACD) &= \frac{1}{2} [4 (0 - 2) + 4 (2 + 6) + 5 (-6 - 0)] \\ &= \frac{1}{2} [-8 + 32 - 30] \\ &= \frac{1}{2} [-6] = -3 \text{ sq. units} \\ &= 3 \text{ sq. units (numerically)} \end{aligned} \quad \dots(2)$$

From (1) and (2)

$$\text{ar } (\Delta ABD) = \text{ar } (\Delta ACD)$$

i.e. A median divides the triangle into two triangles of equal areas.

NCERT TEXTBOOK QUESTIONS SOLVED

EXERCISE 7.4

Q. 1. Determine the ratio in which the line $2x + y - 4 = 0$ divides the line segment joining the points $A (2, -2)$ and $B (3, 7)$.

Sol. Let the required ratio be $k : 1$ and the point C divides them in the above ratio.

∴ Coordinates of C are:

$$\left(\frac{3k+2}{k+1}, \frac{7k-2}{k+1} \right)$$

Since the point C lies on the given line $2x + y - 4 = 0$,
 \therefore We have:

$$2\left(\frac{3k+2}{k+1}\right) + \left(\frac{7k-2}{k+1}\right) - 4 = 0$$

$$\Rightarrow 2(3k+2) + (7k-2) = 4 \times (k+1)$$

$$\Rightarrow 6k + 4 + 7k - 2 - 4k - 4 = 0$$

$$\Rightarrow (6 + 7 - 4)k + (4 - 2 - 4) = 0$$

$$\Rightarrow 9k + (-2) = 0$$

$$\Rightarrow 9k - 2 = 0$$

$$\Rightarrow k = \frac{2}{9}$$

\therefore The required ratio

$$= k : 1$$

$$= \frac{2}{9} : 1$$

$$= 2 : 9$$

Q. 2. Find a relation between x and y if the points (x, y) , $(1, 2)$ and $(7, 0)$ are collinear.

Sol. The given points are:

$A(x, y)$, $B(1, 2)$ and $C(7, 0)$

The points A , B and C will be collinear if

$$x(2-0) + 1(0-y) + 7(y-2) = 0$$

or if $2x - y + 7y - 14 = 0$

or if $2x + 6y - 14 = 0$

or if $x + 3y - 7 = 0$

which is the required relation between x and y .

Q. 3. Find the centre of a circle passing through the points $(6, -6)$, $(3, -7)$ and $(3, 3)$.

Sol. Let $P(x, y)$ be the centre of the circle passing through

$A(6, -6)$, $B(3, -7)$ and $C(3, 3)$

$$\therefore AP = BP = CP$$

Taking $AP = BP$, we have $AP^2 = BP^2$

$$\Rightarrow (x-6)^2 + (y+6)^2 = (x-3)^2 + (y+7)^2$$

$$\Rightarrow x^2 - 12x + 36 + y^2 + 12y + 36 = x^2 - 6x + 9 + y^2 + 14y + 49$$

$$\Rightarrow -12x + 6x + 12y - 14y + 72 - 58 = 0$$

$$\Rightarrow -6x - 2y + 14 = 0$$

$$\Rightarrow 3x + y - 7 = 0 \quad \dots(1) \text{ [Dividing by } (-2)]$$

Taking $BP = CP$, we have $BP^2 = CP^2$

$$\Rightarrow (x-3)^2 + (y+7)^2 = (x-3)^2 + (y-3)^2$$

$$\Rightarrow x^2 - 6x + 9 + y^2 + 14y + 49 = x^2 - 6x + 9 + y^2 - 6y + 9$$

$$\Rightarrow -6x + 6x + 14y + 6y + 58 - 18 = 0$$

$$\Rightarrow 20y + 40 = 0$$

$$\Rightarrow y = \frac{-40}{20} = -2 \quad \dots(2)$$

From (1) and (2),

$$3x - 2 - 7 = 0$$

$$\Rightarrow 3x = 9 \Rightarrow x = 3$$

i.e., $x = 3$ and $y = -2$
 \therefore The required centre is $(3, -2)$.

Q. 4. The two opposite vertices of a square are $(-1, 2)$ and $(3, 2)$. Find the coordinates of the other two vertices.

Sol. Let us have a square $ABCD$ such that $A(-1, 2)$ and $C(3, 2)$ are the opposite vertices.

Let $B(x, y)$ be an unknown vertex.

Since all sides of a square are equal,

$$\therefore AB = BC$$

$$\Rightarrow AB^2 = BC^2$$

$$\Rightarrow (x + 1)^2 + (y - 2)^2 = (x - 3)^2 + (y - 2)^2$$

$$\Rightarrow 2x + 1 = -6x + 9$$

$$\Rightarrow 8x = 8 \Rightarrow x = 1$$

...(1)

Since each angle of a square $= 90^\circ$,

$\therefore ABC$ is a right angled triangle.

\therefore Using Pythagoras theorem, we have:

$$AB^2 + BC^2 = AC^2$$

$$\Rightarrow [(x + 1)^2 + (y - 2)^2] + [(x - 3)^2 + (y - 2)^2] = [(3 + 1)^2 + (2 - 2)^2]$$

$$\Rightarrow 2x^2 + 2y^2 + 2x - 4y - 6x - 4y + 1 + 4 + 9 + 4 = 16$$

$$\Rightarrow 2x^2 + 2y^2 - 4x - 8y + 2 = 0$$

$$\Rightarrow x^2 + y^2 - 2x - 4y + 1 = 0$$

...(2)

Substituting the value of x from (1) into (2) we have:

$$1 + y^2 - 2 - 4y + 1 = 0$$

$$\Rightarrow y^2 - 4y + 2 - 2 = 0$$

$$\Rightarrow y^2 - 4y = 0$$

$$\Rightarrow y(y - 4) = 0$$

$$\Rightarrow y = 0 \text{ or } y = 4$$


Hence, the required other two vertices are: $(1, 0)$ and $(1, 4)$.

Q. 5. The Class X students of a secondary school in Krishinagar have been allotted a rectangular plot of land for their gardening activity. Saplings of Gulmohar are planted on the boundary at a distance of 1 m from each other. There is a triangular grassy lawn in the plot as shown in the Fig. The students are to sow seeds of flowering plants on the remaining area of the plot.

(i) Taking A as origin, find the coordinates of the vertices of the triangle.

(ii) What will be the coordinates of the vertices of ΔPQR if C is the origin?

Also calculate the areas of the triangles in these cases. What do you observe?



Sol. (i) By taking A as the origin and AD and AB as the coordinate axes, we have

$P(4, 6)$, $Q(3, 2)$ and $R(6, 5)$ as the vertices of ΔPQR .

(ii) By taking C as the origin and CB and CD as the coordinate axes, then the vertices of ΔPQR are

P (12, 2), Q (13, 6) and R (10, 3)

Now, ar (ΔPQR) [when P (4, 6), Q (3, 2) and R (6, 5) are the vertices]

$$= \frac{1}{2} [4(2-5) + 3(5-6) + 6(6-2)]$$

$$= \frac{1}{2} [-12 - 3 + 24]$$

$$= \frac{9}{2} \text{ sq. units.}$$

[taking numerical value]

ar (ΔPQR) [when P (12, 2), Q (13, 6) and R (10, 3) are the vertices.]

$$= \frac{1}{2} [12(6-3) + 13(3-2) + 10(2-6)]$$

$$= \frac{1}{2} [36 + 13 - 40]$$

$$= \frac{9}{2} \text{ sq. units.}$$

Thus, in both cases, the area of ΔPQR is the same.

Q. 6. The vertices of a ΔABC are A (4, 6), B (1, 5) and C (7, 2). A line is drawn to intersect sides AB

and AC at D and E respectively, such that $\frac{AD}{AB} = \frac{AE}{AC} = \frac{1}{4}$. Calculate the area of the

ΔADE and compare it with the area of ΔABC . (Recall Theorem 6.2 and Theorem 6.6).

Sol. We have $\frac{AD}{AB} = \frac{1}{4}$

$$\Rightarrow \frac{AB}{AD} = \frac{4}{1}$$

$$\Rightarrow \frac{AD+DE}{AD} = \frac{4}{1}$$

$$\Rightarrow \frac{AD}{AD} + \frac{DE}{AD} = \frac{4}{1} = 1 + \frac{3}{1}$$

$$\Rightarrow 1 + \frac{DE}{AD} = 1 + \frac{3}{1} \Rightarrow \frac{DE}{AD} = \frac{3}{1}$$

$$\Rightarrow AD : DE = 1 : 3$$

Thus, the point D divides AB in the ratio 1 : 3

\therefore **The coordinates of D are:**

$$\left[\frac{(1 \times 1) + (3 \times 4)}{1+3}, \frac{(1 \times 5) + (3 \times 6)}{1+3} \right]$$

$$\text{or } \left[\frac{1+12}{4}, \frac{5+18}{4} \right]$$

$$\text{or } \left(\frac{13}{4}, \frac{23}{4} \right)$$

Similarly, $AE : EC = 1 : 3$

i.e., E divides AC in the ratio 1 : 3

⇒ Coordinates of E are:

$$\left[\frac{(1 \times 7) + (3 \times 4)}{1 + 3}, \frac{1 \times 2 + 3 \times 6}{1 + 3} \right]$$

$$\text{or } \left[\frac{7 + 12}{4}, \frac{2 + 18}{4} \right]$$

$$\text{or } \left[\frac{19}{4}, 5 \right]$$

Now, ar (ΔADE)

$$= \frac{1}{2} \left[4 \left(\frac{23}{4} - 5 \right) + \frac{13}{4} (5 - 6) + \frac{19}{4} \left(6 - \frac{23}{4} \right) \right]$$

$$= \frac{1}{2} \left[(23 - 20) + \frac{13}{4} (1) + \frac{19}{4} \left(\frac{24 - 23}{4} \right) \right]$$

$$= \frac{1}{2} \left(3 - \frac{13}{4} + \frac{19}{16} \right)$$

$$= \frac{1}{2} \left[\frac{48 + 52 + 19}{16} \right] = \frac{15}{32} \text{ sq. units.}$$

Area of ΔABC

$$= \frac{1}{2} [4(5 - 2) + 1(2 - 6) + 7(6 - 5)]$$

$$= \frac{1}{2} [(4 \times 3) + 1 \times (-4) + 7 \times 1]$$

$$= \frac{1}{2} [12 + (-4) + 7]$$

$$= \frac{1}{2} (15) = \frac{15}{2} \text{ sq. units.}$$

$$\text{Now, } \frac{\text{ar}(\Delta ADE)}{\text{ar}(\Delta ABC)} = \frac{\frac{15}{32}}{\frac{15}{2}} = \frac{15}{32} \times \frac{2}{15} = \frac{1}{16}$$

$$\Rightarrow \text{ar}(\Delta ADE) : \text{ar}(\Delta ABC) = 1 : 16.$$

Q. 7. Let A (4, 2), B (6, 5) and C (1, 4) be the vertices of ΔABC .


- The median from A meets BC at D. Find the coordinates of the point D.
- Find the coordinates of the point P on AD such that AP : PD = 2 : 1.
- Find the coordinates of points Q and R on medians BE and CF respectively such that BQ : QE = 2 : 1 and CR : RF = 2 : 1.
- What do you observe?

[Note: The point which is common to all the three medians is called the centroid and this point divides each median in the ratio 2 : 1.]


- If A (x_1, y_1), B (x_2, y_2) and C (x_3, y_3) are the vertices of ΔABC , find the coordinates of the centroid of the triangle.

Sol. We have the vertices of ΔABC as A (4, 2), B (6, 5) and C (1, 4).

- Since AD is a median



\therefore Coordinates of D are: $\left(\frac{6+1}{2}, \frac{5+4}{2}\right)$ or $\left(\frac{7}{2}, \frac{9}{2}\right)$



(ii) Since $AP : PD = 2 : 1$ i.e., P divides AD in the ratio $2 : 1$.

\therefore Coordinates of P are:

$$\left[\frac{2\left(\frac{7}{2}\right) + (1 \times 4)}{2+1}, \frac{2\left(\frac{9}{2}\right) + 1 \times 2}{2+1} \right] \text{ or } \left(\frac{11}{3}, \frac{11}{3}\right)$$

(iii) $BQ : QE = 2 : 1 \Rightarrow$ [The point Q divides BE in the ratio $2 : 1$]

\therefore Coordinates of Q are:

$$\left[\frac{2\left(\frac{5}{2}\right) + 1 \times 6}{2+1}, \frac{(2 \times 3) + (1 \times 5)}{2+1} \right]$$

$$\text{or } \left[\frac{5+6}{3}, \frac{6+5}{3} \right]$$

$$\text{or } \left[\frac{11}{3}, \frac{11}{3} \right]$$

Coordinates of Q are:

$$\left(\frac{4+6}{2}, \frac{2+5}{2}\right) \text{ or } \left(5, \frac{7}{2}\right)$$

Coordinates of R are:

$$\left[\frac{2 \times 5 + 1 \times 1}{2-1}, \frac{2 \times \frac{7}{2} + 1 \times 4}{2+1} \right]$$

$$\text{or } \left[\frac{10+1}{3}, \frac{7+4}{3} \right]$$

$$\text{or } \left[\frac{11}{3}, \frac{11}{3} \right]$$

(iv) We observe that P , Q and R represent the same point.

(v) Here, we have

$A(x_1, y_1)$, $B(x_2, y_2)$, $C(x_3, y_3)$ as the vertices of ΔABC . Also AD , BE and CF are its medians.

$\therefore D$, E and F are the mid points of BC , CA and AB respectively.

We know, the centroid is a point on a median, dividing it in the ratio $2 : 1$.

Considering the median AD ,

Coordinates of AD are:

$$\left[\frac{x_2 + x_3}{2}, \frac{y_2 + y_3}{2} \right]$$

Let G be the centroid.

\therefore Coordinates of the centroid are:


$$\left[\frac{(1 \times x_1) + 2 \left(\frac{x_2 + x_3}{2} \right)}{1 + 2}, \frac{(1 \times y_1) + 2 \left(\frac{y_2 + y_3}{2} \right)}{1 + 2} \right]$$

$$= \left[\frac{x_1 + x_2 + x_3}{3}, \frac{y_1 + y_2 + y_3}{3} \right]$$

Similarly, considering the other medians we find that in each the coordinates of G


are $\frac{x_1 + x_2 + x_3}{3}, \frac{y_1 + y_2 + y_3}{3}$.

i.e., The coordinates of the centroid are $\left[\frac{x_1 + x_2 + x_3}{3}, \frac{y_1 + y_2 + y_3}{3} \right]$.



Q. 8. $ABCD$ is a rectangle formed by the points $A(-1, -1)$, $B(-1, 4)$, $C(5, 4)$ and $D(5, -1)$. P , Q , R and S are the mid points of AB , BC , CD and DA respectively. Is the quadrilateral $PQRS$ a square? a rectangle? or a rhombus? Justify your answer.

Sol. We have a rectangle whose vertices are $A(-1, -1)$, $B(-1, 4)$, $C(5, 4)$ and $D(5, -1)$.



$\therefore P$ is mid-point of AB

\therefore Coordinates of P are:

$$\left[\frac{-1 - 1}{2}, \frac{-1 + 4}{2} \right] \text{ or } \left(-1, \frac{3}{2} \right)$$

Similarly, the coordinates of Q are:



$$\left(\frac{-1+5}{2}, \frac{4+4}{2} \right) \text{ or } (2, 4)$$

Coordinates of R are:

$$\left(\frac{5+5}{2}, \frac{-1+4}{2} \right) \text{ or } \left(5, \frac{3}{2} \right)$$

Coordinates of S are:

$$\left(\frac{-1+5}{2}, \frac{-1-1}{2} \right) \text{ or } (2, -1)$$

$$\text{Now, } PQ = \sqrt{(2+1)^2 + \left(4 - \frac{3}{2}\right)^2} = \sqrt{9 + \frac{25}{4}} = \frac{\sqrt{61}}{2}$$

$$SR = \sqrt{(5-2)^2 + \left(\frac{3}{2} - 4\right)^2} = \sqrt{9 + \frac{25}{4}} = \frac{\sqrt{61}}{2}$$

$$RS = \sqrt{(2-5)^2 + \left\{-1 + \left(-\frac{3}{2}\right)\right\}^2} = \sqrt{9 + \frac{25}{4}} = \frac{\sqrt{61}}{2}$$

$$SP = \sqrt{(2+1)^2 + \left(-1 - \frac{3}{2}\right)^2} = \sqrt{9 + \frac{25}{4}} = \frac{\sqrt{61}}{2}$$

$$SR = \sqrt{(5+1)^2 + \left(\frac{3}{2} - \frac{3}{2}\right)^2} = \sqrt{6^2 + 0} = 6$$

$$QS = \sqrt{(2-2)^2 + (4+1)^2} = \sqrt{0 + 5^2} = 5$$

We see that:

$$PQ = QR = RS = SP$$

i.e., all sides of PQRS are equal.

∴ It can be a square or a rhombus.

But its diagonals are not equal.

i.e., $PR \neq QS$

∴ PQRS is a **rhombus**.

MORE QUESTIONS SOLVED

I. VERY SHORT ANSWER TYPE QUESTIONS

Q. 1. Find a point on the y-axis equidistant from $(-5, 2)$ and $(9, -2)$.

(CBSE 2012)

Sol. Let the required point on the y-axis be $P(0, y)$

$$\therefore PA = PB$$

$$\Rightarrow \sqrt{(0+5)^2 + (y-2)^2} = \sqrt{(0-9)^2 + (y+2)^2}$$

$$\Rightarrow \sqrt{5^2 + y^2 + 4 - 4y} = \sqrt{(-9)^2 + y^2 + 4 + 4y}$$

$$\Rightarrow 25 + y^2 + 4 - 4y = 81 + y^2 + 4 + 4y$$

$$\Rightarrow y^2 - y^2 - 4y - 4y = 81 + 4 - 4 - 25$$

$$\Rightarrow -8y = 85 - 29$$

$$\Rightarrow -8y = 56$$

$$\Rightarrow y = \frac{56}{-8} = -7$$

\therefore The required point is $(0, -7)$.

Q. 2. Find a point on x -axis at a distance of 4 units from the point $A(2, 1)$.

Sol. Let the required point on x -axis be $P(x, 0)$.

$$\therefore PA = 4$$

$$\Rightarrow \sqrt{(x-2)^2 + (0-1)^2} = 4$$

$$\Rightarrow x^2 - 4x + 4 + 1 = 4^2 = 16$$

$$\Rightarrow x^2 - 4x + 1 + 4 - 16 = 0$$

$$\Rightarrow x^2 - 4x - 11 = 0$$

$$\Rightarrow x = 2 \pm \sqrt{15}$$

Thus, the coordinates of P are: $(2 \pm \sqrt{15}, 0)$.

Q. 3. Find the distance of the point $(3, -4)$ from the origin.

Sol. The coordinates of origin $(0, 0)$.

\therefore Distance of $(3, -4)$ from the origin

$$= \sqrt{(3-0)^2 + (-4-0)^2}$$

$$= \sqrt{(3)^2 + (-4)^2} = \sqrt{9+16} = \sqrt{25} = 5.$$

Q. 4. For what value of x is the distance between the points $A(-3, 2)$ and $B(x, 10)$ 10 units?

Sol. The distance between $A(-3, 2)$ and $B(x, 10)$

$$= \sqrt{(x+3)^2 + (10-2)^2}$$

$$\therefore \sqrt{(x+3)^2 + (10-2)^2} = 10$$

$$\Rightarrow (x+3)^2 + (8)^2 = 10^2$$

$$\Rightarrow (x+3)^2 = 10^2 - 8^2$$

$$\Rightarrow (x+3)^2 = (10-8)(10+8) = 36$$

$$\Rightarrow x+3 = \pm \sqrt{36} = \pm 6$$

$$\text{For +ve sign, } x = 6 - 3 = 3$$

$$\text{For -ve sign, } x = -6 - 3 = -9$$

Q. 5. Find a point on the x -axis which is equidistant from the points $A(5, 2)$ and $B(1, -2)$.

Sol. The given points are:

$A(5, 2)$ and $B(1, -2)$

Let the required point on the x -axis be $C(x, 0)$.

Since, C is equidistant from A and B .

$$\therefore AC = BC$$

$$\therefore \sqrt{(x-5)^2 + (0-2)^2} = \sqrt{(x-1)^2 + (0+2)^2}$$

$$\Rightarrow (x-5)^2 + (-2)^2 = (x-1)^2 + (2)^2$$

$$\Rightarrow x^2 + 25 - 10x + 4 = x^2 + 1 - 2x + 4$$

$$\Rightarrow -10x + 2x = 5 - 29$$

$$\Rightarrow -8x = -24$$

$$\Rightarrow x = \frac{-24}{-8} = 3$$

\therefore The required point is $(0, 3)$.

Q. 6. Establish the relation between x and y when $P(x, y)$ is equidistant from the points $A(-1, 2)$ and $B(2, -1)$.

Sol. $\because P$ is equidistant from A and B

$$\therefore PA = PB$$

$$\sqrt{(x+1)^2 + (y-2)^2} = \sqrt{(x-2)^2 + (y+1)^2}$$

$$\Rightarrow (x+1)^2 + (y-2)^2 = (x-2)^2 + (y+1)^2$$

$$\Rightarrow x^2 + 1 + 2x + y^2 - 4y + 4 = x^2 + 4 - 4x + y^2 + 1 + 2y$$

$$\Rightarrow 2x - 4x + 5 = -4x + 2y + 5$$

$$\Rightarrow 2x + 4x + 5 = 2y + 4y + 5$$

$$\Rightarrow 6x = 6y$$

$$\Rightarrow x = y$$

which is the required relation.

Q. 7. Show that the points $(7, -2)$, $(2, 3)$ and $(-1, 6)$ are collinear.

Sol. Here, the vertices of a triangle are $(7, -2)$, $(2, 3)$ and $(-1, 6)$

\therefore Area of the triangle

$$= \frac{1}{2} [x_1(y_2 - y_3) + x_2(y_3 - y_1) + x_3(y_1 - y_2)]$$

$$= \frac{1}{2} [7(3 - 6) + 2(6 + 2) + (-1)(-2 - 3)]$$

$$= \frac{1}{2} [7 \times (-3) + 2 \times 8 + (-1)(-5)]$$

$$= \frac{1}{2} [-21 + 16 + 5]$$

$$= \frac{1}{2} [0] = 0$$

Since area of triangle = 0

\therefore The vertices of the triangle are collinear.

Thus, the given points **are collinear**.

Q. 8. Find the distance between the points

$$\left(-\frac{8}{5}, 2\right) \text{ and } \left(\frac{2}{5}, 2\right)$$

(CBSE 2009)

Sol. Distance between $\left(-\frac{8}{5}, 2\right)$ and $\left(\frac{2}{5}, 2\right)$ is given by

$$\sqrt{\left(\frac{2}{5} + \frac{8}{5}\right)^2 + (2-2)^2} = \sqrt{2^2 - 0^2} = 2 \text{ units.}$$

Q. 9. If the mid point of the line joining the points $P(6, b-2)$ and $Q(-2, 4)$ is $(2, -3)$, find the value of b .
(CBSE 2009 F)

Sol. Here, $P(6, b-2)$ and $Q(-2, 4)$ are the given points.

\therefore Mid point of PQ is given by:

$$\left[\frac{6+(-2)}{2}, \frac{4+b-2}{2} \right]$$

$$\text{or } \left[\frac{6-2}{2}, \frac{4-2+b}{2} \right]$$

$$\text{or } \left[2, \frac{2+b}{2} \right]$$

$$\therefore \frac{2+b}{2} = -3 \Rightarrow 2+b = -6$$

$$\Rightarrow b = -6 - 2$$

$$\Rightarrow b = -8$$

Q. 10. In the given figure, ABC is a triangle. D and E are the mid points of the sides BC and AC respectively. Find the length of DE. Prove that $DE = \frac{1}{2} AB$ (CBSE 2011)

Sol. Co-ordinates of the mid point of BC are:


$$= \left(\frac{-6+2}{2}, \frac{-1+(-2)}{2} \right)$$

$$= \left(-2, \frac{-3}{2} \right) \Rightarrow E \left(-2, \frac{-3}{2} \right)$$

Co-ordinates of the mid point of AC are:

$$= \left(\frac{-6+4}{2}, \frac{-1+(-2)}{2} \right)$$

$$= \left(-1, \frac{-3}{2} \right) \Rightarrow D \left(-1, \frac{-3}{2} \right)$$



$$\begin{aligned} \text{Now, } DE &= \sqrt{(-2+1)^2 + \left(-\frac{3}{2} + \frac{3}{2}\right)^2} \\ &= \sqrt{(-1)^2 + 0} = 1 \\ AB &= \sqrt{(4-2)^2 + (-2+2)^2} \\ &= \sqrt{2^2 + 0} = 2 \end{aligned} \quad \left. \vphantom{\begin{aligned} DE &= \sqrt{(-2+1)^2 + \left(-\frac{3}{2} + \frac{3}{2}\right)^2} \\ &= \sqrt{(-1)^2 + 0} = 1 \\ AB &= \sqrt{(4-2)^2 + (-2+2)^2} \\ &= \sqrt{2^2 + 0} = 2 \end{aligned}} \right\} \Rightarrow DE = \frac{1}{2} AB \text{ Hence proved.}$$

II. SHORT ANSWER TYPE QUESTIONS

Q. 1. Points P (5, -3) is one of the two points of trisection of the line segment joining the points A (7, -2) and B (1, -5) near to A. Find the coordinates of the other point of trisection.

(AI CBSE 2010)

Sol.



Since P is near to A

∴ other point Q is the mid point of PB

$$\Rightarrow x = \frac{5+1}{2} = 3$$


$$\Rightarrow y = \frac{-3-5}{2} = -\frac{8}{2} = -4$$

Thus, the point Q is (3, -4)

Q. 2. Find the area of the quadrilateral ABCD whose vertices are A (1, 0), B (5, 3), C (2, 7) and D (-2, 4). [AI CBSE 2009]

$$\begin{aligned} \text{Sol. Area of } \triangle ABC &= \frac{1}{2} [1(3-7) + 5(7-0) + 2(0-3)] \\ &= \frac{1}{2} [-4 + 35 - 6] = \frac{1}{2} \times 25 \\ &= \frac{25}{2} \text{ sq. units.} \end{aligned}$$

$$\begin{aligned} \text{Area of } \triangle ACD &= \frac{1}{2} [1(7-4) + 2(4-0) + (-2)(0-7)] \\ &= \frac{1}{2} [3 + 8 + 14] = \frac{1}{2} \times 25 \\ &= \frac{25}{2} \text{ sq. units.} \end{aligned}$$




∴ Area of the quad. ABCD

$$\begin{aligned} &= \frac{25}{2} \text{ sq. units} + \frac{25}{2} \text{ sq. units.} \\ &= \mathbf{25 \text{ sq. units.}} \end{aligned}$$

Q. 3. Points P, Q, R and S, in this order, divide a line segment joining A (2, 6), B (7, -4) in five equal parts. Find the coordinates of P and R. [AI CBSE 2009 Comptt.]

Sol.



∴ P, Q, R and S divide AB in five equal parts.

$$\therefore AP = PQ = QR = RS = SB$$

Now, P divides AB in the ratio 1 : 4

∴ Coordinates of P are:

$$\left[\frac{1 \times 7 + 4 \times 2}{1 + 4}, \frac{1 \times (-4) + 4 \times 6}{1 + 4} \right]$$

$$\text{or } \left[\frac{7+8}{5}, \frac{-4+24}{5} \right] \text{ or } (3, 4)$$

Again, R divides AB in the ratio 3 : 2

∴ Coordinates of R are:

$$\left[\frac{2 \times 2 + 3 \times 7}{2 + 3}, \frac{2 \times 6 + 3 \times (-4)}{2 + 3} \right] \text{ or } \left[\frac{4+21}{5}, \frac{0}{5} \right] \text{ or } (5, 0)$$



Q. 4. A $(-4, -2)$, B $(-3, -5)$, C $(3, -2)$ and D $(2, k)$ are the vertices of a quad. ABCD. Find the value of k , if the area of the quad is 28 sq. units.

Sol. Area of quad ABCD = 28 sq. units

$$\therefore [\text{ar}(\Delta ABD)] + [\text{ar}(\Delta BCD)] = 28 \text{ sq. units}$$

$$\Rightarrow \frac{1}{2} [-4(-5-k) - 3(k+2) + 2(-2+5)]$$

$$+ \frac{1}{2} [-3(-2-k) + 3(k+5) + 2(-5+2)] = 28$$

$$\Rightarrow \frac{1}{2} [20 + 4k - 3k - 6 + 6] + \frac{1}{2} [6 + 3k + 3k + 15 - 6] = 28$$


$$\Rightarrow \frac{1}{2} [k + 20] + \frac{1}{2} [6k + 15] = 28$$

$$\Rightarrow k + 20 + 6k + 15 = 56$$

$$\Rightarrow 7k + 35 = 56$$

$$\Rightarrow 7k = 56 - 35 = 21$$

$$\Rightarrow k = \frac{21}{7} = 3$$



Q. 5. Find the point on y -axis which is equidistant from the points $(5, -2)$ and $(-3, 2)$.

Sol. Let the required point be $P(0, y)$

\therefore The given points are A $(5, -2)$ and B $(-3, 2)$

$$\therefore PA = PB$$

$$\Rightarrow PA^2 = PB^2$$

$$\therefore (5-0)^2 + (-2-y)^2 = (-3-0)^2 + (2-y)^2$$

$$\Rightarrow 5^2 + (-2-y)^2 = (-3)^2 + (2-y)^2$$

$$\Rightarrow 25 + 4 + y^2 + 4y = 9 + 4 + y^2 - 4y$$

$$\Rightarrow 25 + 4y = 9 - 4y$$

$$\Rightarrow 8y = -16 \Rightarrow y = -2$$

Thus, the required point is $(0, -2)$

Q. 6. Find the point on y -axis which is equidistant from $(-5, 2)$ and $(9, -2)$. (CBSE 2009 C)

Sol. Let the required point on Y -axis be $P(0, y)$.

The given points are A $(-5, 2)$ and B $(9, -2)$

$$\therefore AP = BP$$

$$\therefore \sqrt{(0+5)^2 + (y-2)^2} = \sqrt{(0-9)^2 + (y+2)^2}$$

$$\Rightarrow 5^2 + (y-2)^2 = 9^2 + (y+2)^2$$

$$\Rightarrow 25 + y^2 - 4y + 4 = 81 + y^2 + 4 + 4y$$

$$\Rightarrow -4y - 4y = 81 + 4 - 4 - 25$$

$$\Rightarrow -8y = 56$$

$$\Rightarrow y = \frac{56}{-8} = -7$$

\therefore The required point = $(0, -7)$

Q. 7. Find the value of x for which the distance between the points $P(4, -5)$ and $Q(12, x)$ is 10 units.

(CBSE 2009 C)

Sol. The given points are $P(4, -5)$ and $Q(12, x)$ such that $PQ = 10$



$$\begin{aligned}
 \therefore \sqrt{(12-4)^2 + (x+5)^2} &= 10 \\
 \Rightarrow (12-4)^2 + (x+5)^2 &= 10^2 \\
 \Rightarrow 8^2 + (x+5)^2 &= 100 \\
 \Rightarrow 64 + x^2 + 25 + 10x &= 100 \\
 \Rightarrow x^2 + 10x - 11 &= 0 \\
 \Rightarrow (x-1)(x+11) &= 0 \\
 \Rightarrow x = 1 \text{ or } x = -11
 \end{aligned}$$

Q. 8. Find the relation between x and y if the points $(2, 1)$, (x, y) and $(7, 5)$ are collinear.

(AI CBSE 2009)

Sol. Here,

$$\begin{aligned}
 x_1 &= 2, & y_1 &= 1 \\
 x_2 &= x, & y_2 &= y \\
 x_3 &= 7, & y_3 &= 5
 \end{aligned}$$

$$\begin{aligned}
 \therefore \text{Area of triangle} &= \frac{1}{2} [x_1 (y_2 - y_3) + x_2 (y_3 - y_1) + x_3 (y_1 - y_2)] \\
 &= \frac{1}{2} [2 (y - 5) + x (5 - 1) + 7 (1 - y)] \\
 &= \frac{1}{2} [2y - 10 + 5x - x + 7 - 7y] \\
 &= \frac{1}{2} [-5y + 4x - 3]
 \end{aligned}$$

$$\therefore \text{ar}(\Delta) = 0$$

$$\therefore \frac{1}{2} [-5y + 4x - 3] = 0$$

$$\Rightarrow 4x - 5y - 3 = 0$$

which is the required relation.

Q. 9. If $A(-2, 4)$, $B(0, 0)$ and $C(4, 2)$ are the vertices of ΔABC , then find the length of the median through the vertex A .

(CBSE 2009 C)

Sol. $\because AD$ is the median on BC


$\therefore D$ is the mid-point of BC .

\Rightarrow Coordinates of D are:

$$\left(\frac{0+4}{2}, \frac{0+2}{2} \right) \text{ i.e., } (2, 1)$$

Now, the length of the median

$$\begin{aligned}
 AD &= \sqrt{(2+2)^2 + (1-4)^2} \\
 &= \sqrt{(4)^2 + (-3)^2} \\
 &= \sqrt{16+9} = \sqrt{25} = 5 \text{ units.}
 \end{aligned}$$



Q. 10. If the points $A(4, 3)$ and $B(x, 5)$ are on the circle with the centre $O(2, 3)$, find the value of x .

(AI CBSE 2009)

Sol. Let $O(2, 3)$ be the centre of the circle.

$$\begin{aligned}
 \therefore OA &= OB \Rightarrow OA^2 = OB^2 \\
 \Rightarrow (4-2)^2 + (3-3)^2 &= (x-2)^2 + (5-3)^2
 \end{aligned}$$

$$\begin{aligned}
 \Rightarrow 2^2 &= (x-2)^2 + 2^2 \\
 \Rightarrow (x-2)^2 &= 0 \\
 \Rightarrow x-2 &= 0 \\
 \Rightarrow x &= 2
 \end{aligned}$$

Thus, the required value of x is **2**.

Q. 11. If the vertices of a Δ are $(2, 4)$, $(5, k)$ and $(3, 10)$ and its area is 15 sq. units, then find the value of ' k '. (AI CBSE 2008)

Sol. The area of the given Δ

$$\begin{aligned}
 &= \frac{1}{2} [2(k-10) + 5(10-4) + 3(4-k)] \\
 &= \frac{1}{2} [2k - 20 + 50 - 20 + 12 - 3k] \\
 &= \frac{1}{2} [-k + 22]
 \end{aligned}$$

But $\text{ar}(\Delta) = 15$ [given]

$$\therefore \frac{1}{2} [-k + 22] = 15$$

$$\Rightarrow -k + 22 = 30$$

$$\Rightarrow -k = 30 - 22 = 8$$

$$\Rightarrow k = -8$$

Q. 12. The vertices of a triangle are: $(1, k)$, $(4, -3)$, $(-9, 7)$ and its area is 15 sq. units. Find the value of k . (AI CBSE 2008)

Sol. Area of the given triangle

$$= \frac{1}{2} [1(-3-7) + 4(7-k) - 9(k+3)] = 15$$

$$\Rightarrow \frac{1}{2} [-10 + 28 - 4k - 9k - 27] = 15$$

$$\Rightarrow -13k - 9 = 30$$

$$\Rightarrow -13k = 39$$

$$\Rightarrow k = \frac{39}{-13} \Rightarrow k = -3$$

Q. 13. Find the area of a ΔABC whose vertices are $A(-5, 7)$, $B(-4, -5)$ and $C(4, 5)$. (AI CBSE 2008)

Sol. Here,

$$\begin{aligned}
 x_1 &= -5, & y_1 &= 7 \\
 x_2 &= -4, & y_2 &= -5 \\
 x_3 &= 4, & y_3 &= 5
 \end{aligned}$$

$$\begin{aligned}
 \text{Now, } \text{ar}(\Delta) &= \frac{1}{2} [x_1(y_2 - y_3) + x_2(y_3 - y_1) + x_3(y_1 - y_2)] \\
 &= \frac{1}{2} [(-5)(-5-5) + (-4)(5-7) + 4(7+5)] \\
 &= \frac{1}{2} [50 + 8 + 48] \\
 &= \frac{1}{2} \times 106 = 53
 \end{aligned}$$

\therefore The required $\text{ar}(\Delta ABC) = \mathbf{53 \text{ sq. units.}}$

Q. 14. Find the value of k such that the points $(1, 1)$, $(3, k)$ and $(-1, 4)$ are collinear.

(AI CBSE 2008)

Sol. For the three points, to be collinear, the area of triangle formed by them must be zero.

$$\therefore \text{Area of triangle} = 0$$

$$\Rightarrow \frac{1}{2} [1(k - 4) + (3)(4 - 1) + (-1)(1 - k)] = 0$$

$$\Rightarrow \frac{1}{2} [k - 4 + 9 - 1 + k] = 0$$

$$\Rightarrow \frac{1}{2} [2k + 4] = 0$$

$$\Rightarrow k + 2 = 0$$

$$\Rightarrow k = -2$$

Q. 15. For what value of p , the points $(-5, 1)$, $(1, p)$ and $(4, -2)$ are collinear?

(CBSE 2008)

Sol. Since the points are collinear,

\therefore The area of the Δ formed by these points must be zero.

$$\text{i.e., } \frac{1}{2} [-5(p + 2) + 1(-2 - 1) + 4(1 - p)] = 0$$

$$\Rightarrow -5p - 10 - 3 + 4 - 4p = 0$$

$$\Rightarrow -9p - 9 = 0$$

$$\Rightarrow -9p = 9$$

$$\Rightarrow p = \frac{9}{-9} = -1$$

Q. 16. For what value of p , are the points $(2, 1)$, $(p, -1)$ and $(-1, 3)$ collinear?

(CBSE 2008)

Sol. \therefore The given points are collinear.

\therefore The area of a triangle formed by these points must be zero.

$$\text{i.e., } \text{Area of triangle} = 0$$

$$\Rightarrow \frac{1}{2} [2(-1 - 3) + p(3 - 1) + (-1)(1 + 1)] = 0$$

$$\Rightarrow \frac{1}{2} [-8 + 2p - 2] = 0$$

$$\Rightarrow \frac{1}{2} [-10 + 2p] = 0$$


$$\Rightarrow -5 + p = 0$$

$$\Rightarrow p = 5$$

Q. 17. Find the ratio in which the line $3x + 4y - 9 = 0$ divides the line segment joining the points $(1, 3)$ and $(2, 7)$.

(CBSE 2008)

Sol. Let the ratio be $k : 1$.



\therefore Coordinates of R are:

$$\left(\frac{2k + 1}{k + 1}, \frac{7k + 3}{k + 1} \right)$$

Since, R lies on the line $3x + 4y - 9 = 0$

$$\therefore 3\left(\frac{2k+1}{k+1}\right) + 4\left(\frac{7k+3}{k+1}\right) - 9 = 0$$

$$\Rightarrow 6k + 3 + 28k + 12 - 9k + 9 = 0$$

$$\Rightarrow (6k + 28k - 9k) + (3 + 12 - 9) = 0$$

$$\Rightarrow 25k + 6 = 0$$


$$\Rightarrow k = \frac{-6}{25}$$

\therefore The required ratio is

$$-6 : 25 \text{ or } 6 : 25$$

Q. 18. If the point P (x, y) is equidistant from the points A (3, 6) and B (-3, 4), prove that $3x + y - 5 = 0$. (AI CBSE 2008)

Sol.



\therefore P is equidistant from A and B.

$$\therefore AP = BP$$

$$\Rightarrow AP^2 = BP^2$$

$$\Rightarrow (x - 3)^2 + (y - 6)^2 = (x + 3)^2 + (y - 4)^2$$

$$\Rightarrow x^2 - 6x + 9 + y^2 - 12y + 36 = x^2 + 6x + 9 + y^2 - 8y + 16$$

$$\Rightarrow x^2 + y^2 - 6x + 45 = x^2 + y^2 + 6x - 8y + 25$$

$$\Rightarrow (-6x - 6x) + (-12y + 8y) + 45 - 25 = 0$$

$$\Rightarrow -12x + 20 - 4y = 0$$

$$\Rightarrow -3x - y + 5 = 0$$


$$\text{or } 3x + y - 5 = 0$$

Q. 19. The coordinates of A and B are (1, 2) and (2, 3). If P lies on AB, then find the coordinates of P such that:

$$\frac{AP}{PB} = \frac{4}{3}$$

(AI CBSE 2008)

Sol. $\therefore \frac{AP}{PB} = \frac{4}{3}$
 $\therefore AP : PB = 4 : 3$



Here, P divides AB internally in the ratio 4 : 3.

\therefore P has coordinates as:

$$\left[\frac{4 \times 2 + 3 \times 1}{4 + 3}, \frac{4 \times 3 + 3 \times 2}{4 + 3} \right]$$

$$\text{or } \left[\frac{8 + 3}{7}, \frac{12 + 6}{7} \right]$$

$$\text{or } \left[\frac{11}{7}, \frac{18}{7} \right]$$

- Q. 20.** If $A(4, -8)$, $B(3, 6)$ and $C(5, -4)$ are the vertices of a ΔABC , $D(4, 1)$ is the mid-point of BC and P is a point on AD joined such that $\frac{AP}{PD} = 2$, find the coordinates of P .

(AI CBSE 2008)

Sol. $\because D$ is the mid-point of BC

$$\therefore \text{We have } D\left[\frac{3+5}{2}, \frac{6-4}{2}\right] \text{ or } D[4, 1]$$

$$\text{Since, } \frac{AP}{PD} = \frac{2}{1}$$


$$\Rightarrow AP : PD = 2 : 1$$

\therefore Coordinates of P are:

$$\left[\frac{2 \times 4 + 1 \times 4}{2+1}, \frac{2 \times 1 + 1 \times (-8)}{2+1}\right]$$

$$\text{or } \left[\frac{8+4}{3}, \frac{2-8}{3}\right]$$

$$\text{or } \left[\frac{12}{3}, \frac{-6}{3}\right] \text{ or } [4, -2]$$



- Q. 21.** Show that the triangle PQR formed by the points $P(\sqrt{2}, \sqrt{2})$, $Q(-\sqrt{2}, -\sqrt{2})$ and $R(-\sqrt{6}, -\sqrt{6})$ is an equilateral triangle.

OR

Name the type of triangle PQR formed by the points $P(\sqrt{2}, \sqrt{2})$, $Q(-\sqrt{2}, -\sqrt{2})$ and $R(-\sqrt{6}, -\sqrt{6})$.

[NCERT Exemplar]

Sol. We have, $P(\sqrt{2}, \sqrt{2})$

$$Q(-\sqrt{2}, -\sqrt{2})$$

$$\text{and } R(-\sqrt{6}, -\sqrt{6})$$

$$\therefore PQ = \sqrt{(\sqrt{2} + \sqrt{2})^2 + (\sqrt{2} + \sqrt{2})^2} = \sqrt{(2\sqrt{2})^2 + (2\sqrt{2})^2}$$

$$= \sqrt{4 \times 2 + 4 \times 2} = \sqrt{8+8} = \sqrt{16} = 4$$

$$PR = \sqrt{(\sqrt{2} + \sqrt{6})^2 + (\sqrt{2} - \sqrt{6})^2}$$

$$= \sqrt{2+6+2\sqrt{12}+2+6-2\sqrt{12}} = \sqrt{2+6+2+6} = \sqrt{16} = 4$$

$$RQ = \sqrt{[(-\sqrt{2}) + \sqrt{6}]^2 + (-\sqrt{2} - \sqrt{6})^2}$$

$$= \sqrt{2+6-2\sqrt{12}+2+6+2\sqrt{12}} = \sqrt{2+6+2+6} = \sqrt{16} = 4$$

$$\text{Since, } PQ = PR = RQ = \text{each } (= 4)$$

$\therefore PQR$ is an equilateral triangle.

- Q. 22.** The line joining the points $(2, -1)$ and $(5, -6)$ is bisected at P . If P lies on the line $2x + 4y + k = 0$, find the value of k .

(AI CBSE 2008)

Sol. We have $A(2, -1)$ and $B(5, -6)$.

$\therefore P$ is the mid point of AB ,

\therefore Coordinates of P are: $\left[\frac{2+5}{2}, \frac{-1-6}{2} \right]$ or $\frac{7}{2}, \frac{-7}{2}$

Since P lies on the line $2x + 4y + k = 0$

\therefore We have:

$$2x + 4y + k = 0 \Rightarrow 2\left(\frac{7}{2}\right) + 4\left(\frac{-7}{2}\right) + k = 0$$

$$\Rightarrow 7 - 14 + k = 0$$

$$\Rightarrow -7 + k = 0 \Rightarrow k = 7$$

Q. 23. Find the point on y -axis which is equidistant from the points $(5, -2)$ and $(-3, 2)$.

(CBSE 2009)

Sol. \therefore Let P is on the y -axis

\therefore Coordinates of P are: $(0, y)$

Since, $PA = PB$

$$\therefore PA^2 = PB^2$$

$$\Rightarrow (5 - 0)^2 + (-2 - y)^2 = (-3 - 0)^2 + (2 - y)^2$$

$$\Rightarrow 25 + 4 + 4y + y^2 = 9 + 4 - 4y + y^2$$

$$\Rightarrow 25 + 4y = 9 - 4y$$


$$\Rightarrow 8y = -16$$

$$\Rightarrow y = \frac{-16}{8} = -2$$

\therefore The required point is $(0, -2)$.

Q. 24. The line joining the points $(2, 1)$ and $(5, -8)$ is trisected at the points P and Q . If point P lies on the line $2x - y + k = 0$, find the value of k . (CBSE 2009)

Sol.



$\therefore AB$ is trisected at P and Q

\therefore Coordinates of P are:

$$\left[\frac{1 \times 5 + 2 \times 2}{1 + 2}, \frac{1 \times (-8) + 2 \times 1}{1 + 2} \right]$$

$$\text{or } \left(\frac{9}{3}, \frac{-6}{3} \right) \text{ or } (3, -2)$$

Since, $P(3, -2)$ lies on $2x - y + k = 0$

\therefore We have:

$$2(3) - (-2) + k = 0$$

$$\Rightarrow 6 + 2 + k = 0$$

$$\Rightarrow 8 + k = 0 \Rightarrow k = -8$$



Q. 25. If $P(x, y)$ is any point on the line joining the points $A(a, 0)$ and $B(0, b)$, then show that:

$$\frac{x}{a} + \frac{y}{b} = 1 \quad (\text{CBSE 2009})$$

Sol. $\because P$ lies on the line joining A and B .

$\therefore A, B$ and P are collinear.

\Rightarrow The area of a Δ formed by $A(a, 0)$, $B(0, b)$ and $P(x, y)$ is zero.

\therefore We have:

$$x_1[y_2 - y_3] + x_2[y_3 - y_1] + x_3[y_1 - y_2] = 0$$

$$\Rightarrow x[0 - b] + a[b - y] + 0[y - 0] = 0$$

$$\Rightarrow -bx + ab - ay = 0$$

$$\Rightarrow -(bx + ay) = -ab$$

$$\Rightarrow bx + ay = ab$$

$$\Rightarrow \frac{bx}{ab} + \frac{ay}{ab} = \frac{ab}{ab} \quad [\text{Dividing by } ab]$$

$$\Rightarrow \frac{x}{a} + \frac{y}{b} = 1$$

Q. 26. Find the point on x -axis which is equidistant from the points $(2, -5)$ and $(-2, 9)$.

(CBSE 2009)

Sol. \because The required point ' P ' is on x -axis.

\therefore Coordinates of P are $(x, 0)$.

\therefore We have

$$AP = PB$$

$$\Rightarrow AP^2 = PB^2$$

$$\Rightarrow (2 - x)^2 + (-5 - 0)^2 = (-2 - x)^2 + (9 - 0)^2$$

$$\Rightarrow 4 - 4x + x^2 + 25 = 4 + 4x + x^2 + 81$$

$$\Rightarrow 4x + 25 = 4x + 81$$


$$\Rightarrow -8x = 56$$

$$\Rightarrow x = \frac{56}{-8} = -7$$

\therefore The required point is $(-7, 0)$.

Q. 27. The line segment joining the points $P(3, 3)$ and $Q(6, -6)$ is trisected at the points A and B such that A is nearer to P . It also lies on the line given by $2x + y + k = 0$. Find the value of k . (CBSE 2009)

Sol. $\because PQ$ is trisected by A such that



$$PA : AQ = 1 : 2$$

\therefore The coordinates of A are:

$$\left[\frac{1 \times 6 + 2 \times 3}{1 + 2}, \frac{1 \times (-6) + 2 \times 3}{1 + 2} \right]$$

$$\text{or } \left[\frac{6 + 6}{3}, \frac{-6 + 6}{3} \right]$$

$$\text{or } \left[\frac{12}{3}, \frac{0}{3} \right] \text{ or } (4, 0).$$

Since, A (4, 0) lies on the line $2x + y + k = 0$

$$\therefore 2(4) + (0) + k = 0$$

$$\Rightarrow 8 + k = 0 \Rightarrow k = -8$$

Q. 28. Find the ratio in which the points (2, 4) divides the line segment joining the points A (-2, 2) and B (3, 7). Also find the value of y. (AI CBSE 2009)

Sol. Let P (2, y) divides the join of A (-2, 2) and B (3, 7) in the ratio k:1

\therefore Coordinates of P are:

$$\frac{3k-2}{k+1}, \frac{7k+2}{k+1}$$

$$\Rightarrow \frac{3k-2}{k+1} = 2 \text{ and } \frac{7k+2}{k+1} = y$$

$$\text{Now, } \frac{3k-2}{k+1} = 2 \Rightarrow 3k-2 = 2k+2 \Rightarrow k = 4$$

$$\text{And } \frac{7k+2}{k+1} = 7 \Rightarrow \frac{7(4)+2}{4+1} = y$$

$$\Rightarrow \frac{30}{5} = y \Rightarrow 6 = y$$


Thus, $y = 6$ and $k = 4$

Q. 29. Find the area of the quadrilateral ABCD whose vertices are:

A (-4, -2), B (-3, -5), C (3, -2) and D (2, 3)

(AI CBSE 2009)

$$\begin{aligned} \text{Sol. Area of } (\Delta ABC) &= \frac{1}{2} [(-4)(-5+2) + (-3)(-2+2) + 3(-2+5)] \\ &= \frac{1}{2} [-4(-3) + (-3)(0) + 3(3)] \\ &= \frac{1}{2} [-12 + 0 + 9] \\ &= \frac{1}{2} [21] = \frac{21}{2} \text{ sq. units.} \end{aligned}$$



$$\begin{aligned} \text{Also, ar } (\Delta ACD) &= \frac{1}{2} [(-4)(-2-3) + 3(3+2) + 2(-2+2)] \\ &= \frac{1}{2} [20 + 15 + 0] \\ &= \frac{35}{2} \text{ sq. units.} \end{aligned}$$


$$\therefore \text{ar (quad. ABCD)} = \text{ar } (\Delta ABC) + \text{ar } (\Delta ACD)$$

$$= \frac{21}{2} + \frac{35}{2} \text{ sq. units.}$$

$$= \frac{56}{2} = 28 \text{ sq. units.}$$

Q. 30. Find the ratio in which the point $(x, 2)$ divides the line segment joining the points $(-3, -4)$ and $(3, 5)$. Also find the value of x . (AI CBSE 2009)

Sol. Let the required ratio = $k : 1$



\therefore Coordinates of the point P are:

$$\left(\frac{3k - 3}{k + 1}, \frac{5k - 4}{k + 1} \right)$$

But the coordinates of P are $(x, 2)$

$$\therefore \frac{5k - 4}{k + 1} = 2 \Rightarrow 5k - 4 = 2k + 2$$

$$\Rightarrow 3k = 6$$

$$\Rightarrow k = \frac{6}{3} = 2$$

\therefore The required ratio is $2 : 1$

$$\begin{aligned} \text{Now, } x &= \frac{3k - 3}{k + 1} \\ &= \frac{3(2) - 3}{2 + 1} \\ &= \frac{6 - 3}{3} = \frac{3}{3} = 1 \end{aligned}$$

Q. 31. Find the area of the triangle formed by joining the mid-points of the sides of triangle whose vertices are $(0, -1)$, $(2, 1)$, and $(0, 3)$. (AI CBSE 2009)

Sol. We have the vertices of the given triangle as $A(0, -1)$, $B(2, 1)$ and $C(0, 3)$. Let D , E and F be the mid-points of AB , BC and AC .

$$\therefore \text{Coordinates of } D \text{ are } \left[\frac{0+2}{2}, \frac{-1+1}{2} \right] \text{ or } (1, 0)$$

$$E \text{ are } \left[\frac{2+0}{2}, \frac{1+3}{2} \right] \text{ or } (1, 2)$$

$$F \text{ are } \left[\frac{0+0}{2}, \frac{3+(-1)}{2} \right] \text{ or } (0, 1)$$

\therefore Coordinates of the vertices of $\triangle DEF$ are $(1, 0)$, $(1, 2)$ and $(0, 1)$.

$$\begin{aligned} \text{Now, area of } \triangle DEF &= \frac{1}{2} [1(2 - 1) + 1(1 - 0) + 0(0 - 2)] \\ &= \frac{1}{2} \times 2 = 1 \text{ sq. units.} \end{aligned}$$


Q. 32. Find the area of the $\triangle ABC$ with $A(1, -4)$, and the mid-point of sides through A being $(2, -1)$ and $(0, -1)$. [NCERT Exemplar]

Sol. Let the co-ordinates of B and C are (a, b) and (x, y) respectively.

Sides through A are AB and AC

$$\begin{aligned}\therefore (2, -1) &= \left(\frac{1+a}{2}, \frac{-4+b}{2} \right) \\ &= \frac{1+a}{2} = 2 \text{ and } \frac{-4+b}{2} = -1 \\ &= 1+a = 4 \text{ and } -4+b = -2 \\ &= a = 3 \text{ and } b = 2\end{aligned}$$

$$\begin{aligned}\text{Also, } (0, -1) &= \left(\frac{1+x}{2}, \frac{-4+y}{2} \right) \\ &= \frac{1+x}{2} = 0 \text{ and } \frac{-4+y}{2} = -1 \\ &= 1+x = 0 \text{ and } -4+y = -2 \\ &= x = -1 \text{ and } y = 2\end{aligned}$$




Thus, the co-ordinates of the vertices of $\triangle ABC$ are: A(1, -4), B(3, 2) and C(-1, 2)

\therefore Area of $\triangle ABC$

$$\begin{aligned}&= \frac{1}{2} [1(2-2) + 3(2+4) - 1(-4-2)] \\ &= \frac{1}{2} [0 + 18 + 6] \\ &= \frac{1}{2} [24] \\ &= 12 \text{ sq. units}\end{aligned}$$

Q. 33. Find the ratio in which the point (x, -1) divides the line segment joining the points (-3, 5) and (2, -5). Also find the value of x. (AI CBSE 2009)

Sol. Let the required ratio is k : 1



\therefore The coordinates of P are:

$$\left[\frac{2k-3}{k+1}, \frac{-5k+5}{k+1} \right]$$

But the coordinates of P are (x, -1)

$$\therefore \frac{-5k+5}{k+1} = -1 \Rightarrow -5k+5 = -k-1$$

$$\Rightarrow 2k = 3 \text{ or } k = \frac{3}{2}$$

$$\text{Also, } x = \frac{2k-3}{k+1} = \frac{2\left(\frac{3}{2}\right)-3}{\frac{3}{2}+1} = \frac{3-3}{\frac{5}{2}} = 0$$

$$\therefore x = 0$$

$$\text{And } k = \frac{3}{2}$$

Q. 34. If the mid-point of the line segment joining the point A(3, 4) and B(k, 6) is P(x, y) and $x + y - 10 = 0$, then find the value of k. [NCERT Exemplar]

Sol. \therefore Mid point of the line segment joining A(3, 4) and B(k, 6)

$$= \left(\frac{3+k}{2}, \frac{4+6}{2} \right) = \left(\frac{3+k}{2}, 5 \right)$$

$$\therefore \left(\frac{3+k}{2}, 5 \right) = (x, y) \Rightarrow \frac{3+k}{2} = x \text{ and } 5 = y$$

Since, $x + y - 10 = 0$

$$\Rightarrow \frac{3+k}{2} + 5 - 10 = 0 \quad \left[\because x = \frac{3+k}{2} \text{ and } y = 5 \right]$$

$$\Rightarrow 3 + k + 10 - 20 = 0$$

$$\Rightarrow 3 + k = 10$$

$$\Rightarrow k = 10 - 3 = 7$$

Thus, the required value of k = 7

Q. 35. Point P, Q, R and S divide the line segment joining the points A (1, 2) and B (6, 7) in 5 equal parts. Find the co-ordinates of the points P, Q and R. [AI. CBSE (Foreign) 2014]

Sol. $A(1, 2) \bullet \xrightarrow{\quad} P \bullet \xrightarrow{\quad} Q \bullet \xrightarrow{\quad} R \bullet \xrightarrow{\quad} S \bullet \xrightarrow{\quad} B(6, 7)$

\therefore P, Q, R and S, divide AB into five equal parts.

$$\therefore AP = PQ = QR = RS = SB$$

Now, P divides AB in the ratio 1 : 4

Let, the co-ordinates of P be x and y.

\therefore Using the section formula i.e.,

$$x = \frac{mx_2 + nx_1}{m+n}, y = \frac{my_2 + ny_1}{m+n}, \text{ we have}$$

$$\therefore x = \frac{1(6) + 4(1)}{1+4} = \frac{6+4}{5} = 2$$

$$y = \frac{1(7) + 4(2)}{1+4} = \frac{7+8}{5} = 3$$

$$(x, y) = (2, 3)$$

Next, Q divides AB in the ratio 2 : 3

\therefore Co-ordinates of Q are :

$$\left[\frac{2(6) + 3(1)}{2+3}, \frac{2(7) + 3(2)}{5} \right] \text{ or } \left[\frac{15}{5}, \frac{20}{5} \right] \text{ or } (3, 4)$$

Now, R divides AB in the ratio 3 : 2

\Rightarrow Co-ordinates of R are :


$$\left[\frac{3(6) + 2(1)}{3+2}, \frac{3(7) + 2(2)}{5} \right] \text{ or } \left(\frac{20}{5}, \frac{25}{5} \right) \text{ or } (4, 5)$$

The co-ordinates of P, Q and R are respectively :

(2, 3), (3, 4) and (4, 5).


TEST YOUR SKILLS

- The line-segment joining the points $(3, -4)$ and $(1, 2)$ is trisected at the points P and Q .
If the coordinates of P and Q are $(p, -2)$ and $\left(\frac{5}{3}, q\right)$ respectively, find the values of p and q .
[CBSE 2005]
- In the figure, find the coordinates of A .
[AI CBSE 2005]




- The line joining the points $(2, 1)$ and $(5, -8)$ is trisected at the points P and Q . If P lies on the line $2x - y + k = 0$, find the value of k .
[AI CBSE 2006]
- If the coordinates of the mid-points of the sides of a Δ are $(10, 5)$, $(8, 4)$ and $(6, 6)$, then find the coordinates of its vertices.
[AI CBSE 2006]
- Find the coordinates of the points which divide the line segment joining the points $(-4, 0)$, and $(0, 6)$ in three equal parts.
[CBSE 2005C]
- Find the coordinates of the point equidistant from the points $A(1, 2)$, $B(3, -4)$ and $C(5, -6)$.
[CBSE 2005C]
- Prove that the points $A(-4, -1)$, $B(-2, -4)$, $C(4, 0)$ and $D(2, 3)$ are the vertices of a rectangle.
[CBSE 2005C]
- Find the coordinates of the points which divide the line-segment joining the points $(-4, 0)$ and $(0, 6)$ in four equal parts.
[CBSE 2005C]
- Find the coordinates of the points which divide the line-segment joining the points $(2, -2)$ and $(-7, 4)$ in three equal parts.
[CBSE 2011]
- Find the coordinates of the point equidistant from the points $A(5, 1)$, $B(-3, -7)$ and $C(7, -1)$.
[AI CBSE 2005 Comptt]
- The vertices of a ΔABC are given by $A(2, 3)$ and $B(-2, 1)$ and its centroid is $G\left(1, \frac{2}{3}\right)$.
Find the coordinates of the third vertex C of the ΔABC .
[AI CBSE 2005C]
- If the points (x, y) is equidistant from the points $(a + b, b - a)$ and $(a - b, a + b)$. Prove that $bx = ay$.
[AI CBSE 2005 Comptt]
- Two vertices of a ΔABC are given by $A(6, 3)$ and $B(-1, 7)$ and its centroid is $G(1, 5)$. Find the coordinates of the third vertex of the ΔABC .
[AI CBSE 2005C]
- Two of the vertices of a ΔABC are given by $A(6, 4)$ and $B(-2, 2)$ and its centroid is $G(3, 4)$. Find the coordinates of the third vertex C of the ΔABC .
[AI CBSE 2005C]
- The coordinates one end point of a diameter of a circle are $(4, -1)$. If the coordinates of the centre be $(1, -3)$ find the coordinates of the other end of the diameter. [CBSE 2006]
- Show that the points $A(1, 2)$, $B(5, 4)$, $C(3, 8)$ and $D(-1, 6)$ are the vertices of a square.
[CBSE 2006]

17. Find the value of P for which the points $(-1, 3)$, $(2, p)$ and $(5, -1)$ are collinear.
[CBSE 2006]
18. Find the distance of the point $(-6, 8)$ from the origin.
[AI CBSE 2006]
19. Find the coordinates of the point equidistant from three given points $A(5, 3)$, $B(5, -5)$ and $C(1, -5)$.
[AI CBSE 2006]
20. Find the value of p for which the points $(-5, 1)$, $(1, p)$ and $(4, -2)$ are collinear.
[AI CBSE 2006]
21. Find the coordinates of the point on the line joining $P(1, -2)$ and $Q(4, 7)$ that is twice as far from P as from Q .
[AI CBSE 2006]
22. Find the perimeter of a triangle with vertices $(0, 4)$, $(0, 0)$ and $(3, 0)$.
[CBSE 2011, NCERT Exemplar Problem]
23. In the following figure, find the length of the median AD .
[CBSE 2006C]



24. Prove that the points $(3, 0)$, $(6, 4)$ and $(-1, 3)$ are vertices of right angled triangle. Also prove that these are the vertices of an isosceles triangle.
[CBSE 2006C]
25. In what ratio is the line segment joining the points $(-2, -3)$ and $(3, 7)$ divided by the y -axis? Also, find the coordinates of the point of division.
[CBSE 2006C]
26. If $A(5, -1)$, $B(-3, -2)$ and $C(-1, 8)$ are the vertices of $\triangle ABC$, find the length of median through A and the coordinates of the centroid.
[CBSE 2006C]
27. Find the value of k if the points $A(2, 3)$, $B(4, k)$ and $C(6, -3)$ are collinear.
[CBSE 2006C]
28. In the figure, A and B are the end points of a diameter of a circle having its centre at $(1, 2)$. If the coordinates of A are $(-3, 5)$ find the coordinates of point B .
[AI CBSE 2006C]
29. If $(-2, -1)$, $(a, 0)$, $(4, b)$ and $(1, 2)$ are the vertices of a parallelogram, find the value of ' a ' and ' b '.
[AI CBSE 2006C]
30. The vertices of a triangle are $(-1, 3)$, $(1, -1)$ and $(5, 1)$. Find the lengths of medians through vertices $(-1, 3)$ and $(5, 1)$.
[AI CBSE 2006C]
31. By distance formula, show that the points $(1, -1)$, $(5, 2)$ and $(9, 5)$ are collinear.
[AI CBSE 2006C]
32. Show that the points $(7, 10)$, $(-2, 5)$ and $(3, -4)$ are the vertices of an isosceles right triangle.
[CBSE 2007]
33. In what ratio the line $x - y - 2 = 0$ divides the line segment joining $(3, -1)$ and $(8, 9)$?
[CBSE 2007]
34. Find the ratio in which the line joining the points $(6, 4)$ and $(1, -7)$ is divided by x -axis.
[CBSE 2007]



35. Find the ratio in which the point $(-3, k)$ divides the line segment joining the points $(-5, -4)$ and $(-2, 3)$. Hence find the value of k . [AI CBSE 2007]
36. Three consecutive vertices of a parallelogram are $(-2, -1)$, $(1, 0)$ and $(4, 3)$. Find the coordinates of the fourth vertex. [AI CBSE 2007]
37. For what value P , are the points $(2, 1)$, $(p, -1)$ and $(-1, 3)$ collinear?
38. Show that the point $P(-4, 2)$ lies on the line segment joining the points $A(-4, 6)$ and $B(-4, -6)$.
39. Find the value (s) of k for which the points $[(3k - 1), (k - 2)]$, $[k, (k - 7)]$ and $[(k - 1), (-k - 2)]$ are collinear. [AI CBSE (Foreign) 2014]

Hint: Here, $x_1 = (3k - 1)$ $x_2 = k$ $x_3 = (k - 1)$
 $y_1 = (k - 2)$ $y_2 = (k - 7)$ $y_3 = (-k - 2)$

For the given points to be collinear, we have
 $(3k - 1)[(k - 7) - (-k - 2)] + k[(-k - 2) - (k - 2)] + (k - 1)[(k - 2) - (k - 7)] = 0$
 $\Rightarrow 6k^2 - 17k + 5 - 2k^2 + 5k - 5 = 0$ or $4k^2 - 12 = 0 \Rightarrow k = 0, 3$

40. If the point $A(0, 2)$ is equidistant from the point $B(3, p)$ and $C(p, 5)$, find p . [CBSE (Delhi) 2014]

Hint: Here, $AB = AC$

$\therefore \sqrt{3^2 + (p - 2)^2} = \sqrt{p^2 + 3^2} \Rightarrow 9 + (p - 2)^2 = p^2 + 9$
 $\Rightarrow (p - 2)^2 = p^2 \Rightarrow p^2 + 4 - 4p = p^2 \Rightarrow p = 1$


41. Find the ratio in which the point $P(x, 2)$ divides the line segment joining the points $A(12, 5)$ and $B(4, -3)$. Also find the value of x . [CBSE (Delhi) 2014]

Hint: Here, $x_1 = 12, x_2 = 4, y_1 = 5$ and $y_2 = -3$
 Let $P(x, 2)$ divide the given line segment in the ratio $k : 1$
 \therefore co-ordinates of P are :

$x = \frac{k(4) + 1(12)}{k + 1}, \quad 2 = \frac{k(-3) + 1(5)}{k + 1}$
 $\Rightarrow [x(k + 1) = 4k + 12], \quad [2(k + 1) = -3k + 5]$
 $2(k + 1) = -3k + 5 \Rightarrow 2k + 2 = -3k + 5 \Rightarrow 5k = 3$ or $k = \frac{3}{5}$

$x(k + 1) = k(4) + 12 \Rightarrow x\left[\left(\frac{3}{5}\right) + 1\right] = \frac{3}{5}(4) + 12$
 $\Rightarrow \frac{8}{5}x = \frac{72}{5} \quad \text{or} \quad x = \frac{72}{5} \times \frac{5}{8} = 9$
 i.e., $x = 9$ and ratio $= 3 : 5$

42. If the point $P(k - 1, 2)$ is equidistant from the point $A(3, k)$ and $B(k, 5)$ then find the values of k . [AI. CBSE 2014]

Hint: 

$$AP = BP \Rightarrow \sqrt{(k-1)-3]^2 + (2-k)^2} = \sqrt{(k-1-k)^2 + (2-5)^2}$$

$$\Rightarrow \sqrt{(k-4)^2 + (2-k)^2} = \sqrt{(-1)^2 + (-3)^2}$$

$$\Rightarrow \text{Solving it we get } k = 5$$

43. Find the ratio in which the line segment joining the points A(3, -3) and B(-2, 7) is divided by x-axis. Also find the co-ordinates of the point of division. [AI CBSE 2014]

Hint: Let the x-axis meets AB at P(x, 0)

Let (k : 1) is the required ratio.

∴ Co-ordinates of P are given as

$$x = \frac{3k-2}{k+1}$$


$$y = \frac{-3k+7}{k+1} = 0 \Rightarrow k = \frac{7}{3}$$

Now,

$$x = \frac{3\left(\frac{7}{3}\right)-2}{\frac{7}{3}+1} = \frac{3}{2}$$

We get $x = \frac{3}{2}$ and Required ratio as 7 : 3

⇒ co-ordinates of the point of division $\left(\frac{3}{2}, 0\right)$



44. The mid-point P of the line segment joining the points A(-10, 4) and B(-2, 0) lies on the line segment joining the points C(-9, -4) and D(-4, y). Find the ratio in which P divides CD. Also find the value of y. [AI. CBSE (Foreign) 2014]

Hint: Mid point of AB [where A (-10, 4) and B (-2, 0)] is P(-6, 2)

Let P(-6, 2) divide CD, [where C(-9, -4) and D(-4, y)] in k : 1

$$\therefore \frac{k(-4)+1(-9)}{k+1} = -6 \Rightarrow k = \frac{3}{2} \text{ and } 2 = \frac{k(y)+1(-4)}{k+1} \Rightarrow y = 6$$

∴ Required ratio = 3 : 2 and y = 6.

ANSWERS

Test Your Skills

1. $p = \frac{7}{3}; q = 0$

2. A(10, -2)

3. $k = -8$ or $k = -13$

4. (4, 5), (8, 3) and (12, 3)

5. $\left(\frac{-8}{3}, 2\right), \left(\frac{-4}{3}, 3\right)$

6. (-1, -2)

7. ____

8. $\left(-3, \frac{3}{2}\right), (2, 3)$ and $\left(-1, \frac{9}{2}\right)$

9. (-1, 0) and (-4, 2)



10. $(2, -4)$

13. $(-2, 5)$

16. ____

19. $(3, -1)$

22. 12 units

25. $2 : 3, (0, 1)$

28. $(5, -1)$

31. ____

34. $4 : 7$

37. $p = 5$

11. $(3, -2)$

14. $(5, 6)$

17. $p = 1$

20. $p = -1$

23. 5 cm

26. $\sqrt{65}; \left(\frac{1}{3}, \frac{5}{3}\right)$

29. $a = 1, b = 3$

32. ____

35. $2 : 1, k = \frac{2}{3}$

38. ____

12. ____

15. $(-2, -5)$

18. 10

21. $(2, 1)$

24. ____

27. $k = 0$

30. $\left(\frac{5}{3}, 1\right)$

33. $2 : 3$

36. $(1, 2), (1, 4)$

