
# 12. Basic principles and techniques in organic chemistry



- Acyclic or open chain compounds consist of straight or branched chain compounds.
- Alicyclic or closed chain or ring compounds contain carbon atoms joined in the form of ring (homocyclic). In some rings (heterocyclic), atoms other than carbon are present.
- Benzenoid aromatic compounds (include benzene and other related compounds)
- Non-benzenoid compounds (do not contain benzene ring)
- Functional group: An atom or group of atoms joined in a specific manner which is responsible for the characteristic chemical properties of the organic compound. For example, alcohol, aldehyde etc.
- Homologous series: A group or a series of organic compounds each containing a characteristic functional group. Successive members differ from each other in molecular formula by a -CH<sub>2</sub> unit.

## **Melting Point**


- The normal melting point of a solid is defined as the temperature at which the solid state and liquid state of that solid are in equilibrium at an external pressure of one atmosphere.
- Generally, the capillary tube method is used for the determination of the melting point of a compound but this method is less useful for oil and fat.
- Pure compounds have sharp and well-defined melting point, while impure compounds do not have sharp melting point.
- The concept of melting point is used for the identification and characterisation of organic compounds.

# **Boiling Point**

- Boiling point of a liquid is defined as the temperature at which the vapour pressure of the liquid becomes equal to the atmospheric pressure.
- For the determination of boiling point of a liquid, a micro boiling point apparatus based on Thiele's tube is used.
- Boiling point of a liquid decreases as the atmospheric pressure or applied pressure decreases.
- It is the physical property of the any compound by which we can identify its purity.
- Isomerism:







#### Structural isomerism

1. Chain isomerism: Two or more compounds having the same molecular formula, but different carbon skeletons

2. Position isomerism: Two or more compounds differing in the position of functional group on the carbon skeleton

$$\begin{array}{ccc} & & \text{OH} \\ & | \\ \text{CH}_3 - \text{CH}_2 - \text{CH}_2 - \text{CH}_2 - \text{OH} & \text{CH}_3 - \text{CH}_2 - \text{CH} - \text{CH}_3 \\ \\ & \text{Butan-1-ol} & \text{Butan-2-ol} \end{array}$$

3. Functional group isomerism: Two or more compounds having the same molecular formula, but different functional groups

$$CH_3 - CH_2 - C - CH_3$$
  $CH_3 - CH_2 - CH_2 - C - H_3$ 

Butanone Butanal

4. Metamerism: Two or more compounds arising due to different alkyl chains on either side of the functional group in a molecule

#### Stereoisomerism:

Compounds having the same constitution and sequence of covalent bonds, but different relative positions of their atoms or groups in space





# Fundamental concepts in organic reaction mechanism:

#### Fission of a covalent bond

Heterolytic cleavage: Formation of cations and anions takes place.

The increasing order of stability of carbocations is

Homolytic cleavage: Formation of free radicals takes place.

The increasing order of stability of alkyl radicals is

$${\overset{\circ}{C}}{H}_3 < {\overset{\circ}{C}}{H}_2{CH}_3 < {\overset{\circ}{C}}{H}({CH}_3)_2 < {\overset{\circ}{C}}({CH}_3)_3$$

### **Nucleophiles and electrophiles**

Nucleophile (Nu:): Nucleus seeking. For example: hydroxide (HO<sup>-</sup>), cyanide (CN<sup>-</sup>),

Electrophile (E<sup>+</sup>): Electron seeking. For example: carbonyl group (>C=O) or alkyl halides

