## 14. Alkenes

## Alkenes:

- General formula is  $C_nH_{2n}$ .
- Isomerism
- 1.Position isomer

$$\begin{array}{c} ^{1}{\rm CH_{2} = \overset{2}{\rm CH} - \overset{3}{\rm CH_{2}} - \overset{4}{\rm CH_{2}} - \overset{5}{\rm CH_{3}} } \\ & {\rm Pent} - 1 - {\rm ene} \\ ^{1}{\rm CH_{3} - CH} = \overset{3}{\rm CH} - \overset{4}{\rm CH_{2}} - \overset{5}{\rm CH_{3}} \\ & {\rm Pent} - 2 - {\rm ene} \end{array}$$

## 2. Geometrical isomerism

• Cis-isomer: Two identical atoms or groups are on the same side of the double bond

$$C = C$$
H

 $C = C$ 
H

 $CH_3$ 
 $CH_3$ 

• Trans-isomer: Two identical atoms or groups lie on the opposite side of the double bond

$$C = C$$
 $CH_3$ 
 $trans$ -But-2-ene

## Alkenes:

- General formula is  $C_nH_{2n}$ .
- Alkenes are unsaturated hydrocarbons containing at least one double bond.
- Carbon-carbon double bond in alkenes consist of one sigma bond and one pi bond.
- Prepared by:
  - Partial reduction of alkynes
  - Dehydrohalogenation by heating alkyl halides with acloholic KOH
  - Dehalogenation by reacting vicinal dihalides with Zn metal
  - Dehydration of alcohols using
    - Concentrated H<sub>2</sub>SO<sub>4</sub>
    - Heated Al<sub>2</sub>O<sub>3</sub>
- Properties:
  - Colourless, odourless, insoluble in water and fairly soluble in non-polar solvents
  - Undergo addition reactions:







- Addition of hydrogen to form alkanes
- Addition of halogen to form dihalides
- Addition of hydrogen halide to form alkyl halides
- Addition of H<sub>2</sub>SO<sub>4</sub>
- Addition of water
- Oxidation of alkenes with
  - Baeyer's reagent converts them to vicinal glycols
  - Acidic KMnO<sub>4</sub> or acidic K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub> oxidises them to give ketones or acids (depending upon the nature of alkenes)
- Undergoes ozonolysis to form aldehydes
- Polymerises at high temperature in presence of suitable catalyst
- Undergoes combustion to form a large amount of heat
- Uses:
  - Manufacture of synthetic chemicals, polythene, raw materials for detergents
  - For ripeninng of fruits
  - producing oxy-ethylene





