
4. Graphs

• Finding coordinates of any points P and Q, in the given graph.

Consider the given graph.



Here, the coordinates of the points P and Q are (0, 5) and (4, 9) respectively

• Cartesian plane and the terms associated with it

To identify the position of an object or a point in a plane, we require two perpendicular lines: one of them is horizontal and the other is vertical.


Example:

Put an eraser on a book and then describe the position of the eraser.

Solution:

In order to identify the position of the eraser on the book, we take the adjacent edges as perpendicular lines. Take 1 unit = 1 cm along the vertical and horizontal lines.


Now, it is seen that the eraser is at a distance of 11 cm from the vertical line and 10 cm from the horizontal line.



Thus, conventionally, the position of the eraser can be written as (11, 10).

Cartesian system

A Cartesian system consists of two perpendicular lines: one of them is horizontal and the other is vertical. The horizontal line is called the x- axis and the vertical line is called the y-axis. The point of intersection of the two lines is called origin, and is denoted by O.



- XOX' is called the x-axis; YOY' is called the y-axis; the point O is called the origin.
- Positive numbers lie on the directions of OX and OY.
- Negative numbers lie on the directions of OX' and OY'.
- OX and OY are respectively called positive x-axis and positive y-axis.
- OX' and OY' are respectively called negative x-axis and negative y-axis. The axes divide the plane into four equal parts. The four parts are called quadrants, numbered I, II, III and IV, in anticlockwise from positive x-axis, OX.
- The plane is also called co-ordinate plane or Cartesian plane or xy -plane.

Coordinate Geometry

Example:

Name the quadrant or the axis in which the points (5, -4), (2, 7) and (0, -9) lie?

Solution

The coordinates of the point (5, -4) are of the form (+, -).

(5, -4) lie in quadrant IV

The coordinates of the point (2, 7) are of the form (+, +).

(2, 7) lie in quadrant I.

The coordinates of the point (0, -9) are of the form (0, b).

(0, -9) lie on the y-axis

The coordinates of a point on the coordinate plane can be determined by the following conventions.

The x-coordinate of a point is its perpendicular distance from the y-axis, measured along the x-axis (positive along the positive x-axis and negative along the negative x-axis).

The x-coordinate is also called the abscissa.








The y-coordinate of a point is its perpendicular distance from the x-axis, measured along the y-axis (positive along the positive y-axis and negative along the negative y-axis)

The y-coordinate is also called the ordinate.

In stating the coordinates of a point in the coordinate plane, the *x*-coordinate comes first and then the *y*-coordinate. The coordinates are placed in brackets.





What are the coordinates of points A, B and C in the given figure?

Solution:

It is observed that x-coordinate of point A is 5 y-coordinate of point A is 2 Coordinates of point A are (5, 2). x-coordinate of point C is -5 y-coordinate of point C is 2 Coordinates of point C are (-5, 2).

Note: The coordinates of the origin are (0, 0). Since the origin has zero distance from both the axes, its abscissa and ordinate are both zero.

• Relationship between the signs of the coordinates of a point and the quadrant of the point in which it lies:

The 1st quadrant is enclosed by the positive *x*-axis and positive *y*-axis. So, a point in the 1st quadrant is in the form (+, +). The 2nd quadrant is enclosed by the negative *x*-axis and positive *y*-axis. So, a point in the 2nd quadrant is in the form (-, +). The 3rd quadrant is enclosed by the negative *x*-axis and the negative *y*-axis. So, the point in the 3rd quadrant is in the form (-, -).





The 4^{th} quadrant is enclosed by the positive *x*-axis and the negative *y*-axis. So, the point in the 4^{th} quadrant is in the form (+, -).

• Location of a point in the plane when its coordinates are given

Example: Plot the following ordered pairs of numbers (x, y) as points in the coordinate plane. [Use the scale 1 cm = 1 unit]

x	-3	4	-3	0
y	4	-3	-3	2


Solution:

x	-3	4	-3	0
v	4	-3	-3	2

Taking 1 cm = 1 unit, we draw the x-axis and y-axis.

The pairs of numbers in the given table can be represented as (-3, 4), (4, -3) and (-3, -3), (0, 2).

These points can be located in the coordinate plane as:



NB: The coordinates of the point on the x-axis are of the form (a, 0) and the coordinates of the point on the y-axis are of the form (0, b), where a, b are real numbers.

• We can plot a point in the Cartesian plane, if the coordinates of the points are given.

Example:

Plot the points A (5, -3) and B (-2, 5) on the Cartesian plane.

Solution:

To plot A (5, -3):

(1) Move 5 units along OX and mark the endpoint as M.






(2) From M and perpendicular to the x-axis, move 3 units along OY'. Mark the endpoint as A. This is the location of the point (5, -3) on the Cartesian plane.

To plot B (-2, 5):

- (1) Move 2 units along OX' and mark the endpoint as N.
- (2) From N and perpendicular to the x-axis, move 5 units along OY. Mark the endpoint as B. This is the location of the point (-2, 5) on the Cartesian plane.

Points A and B are plotted in the following graph.



- The graph of x = a is a straight line parallel to the y-axis, situated at a distance of a units from y-axis.
- The graph of y = b is a straight line parallel to the x-axis, situated at a distance of b units from x-axis.

Example:

Represent the equation 2y + 5 = 0, on Cartesian plane.

Solution:


$$2y + 5 = 0$$

$$\Rightarrow 2y = -5$$

$$\Rightarrow$$
 $y = \frac{-5}{2} = -2.5$, which is of the form $y = b$.

The graph of this equation can be drawn as follows:





• Graphical solution of linear equation in two variables:

Every point on the graph of a linear equation in two variables is a solution of the linear equation and moreover, every solution of the linear equation is a point on the graph of the linear equation.

Example:

A bag contains some Re 1 coins and some Rs 2 coins. The total worth of coins is Rs 45. Find the number of Re 1 coins, if there are 10 coins of Rs 2.

Solution:

Let there be *x* coins of Re 1 and *y* coins of Rs 2.

Thus,
$$1x + 2y = 45$$

$$\Rightarrow x + 2y = 45$$

This is the required linear equation of the given information. The three solutions of this equation have been given in the tabular form as follows:


X	5	15	25
y	20	15	10

By plotting the points (5, 20), (15, 15) and (25, 10), we obtain the following graph.









From the above graph, it can be seen that the value of x corresponding to y = 10 is 25.

Therefore, there are 25 coins of Re 1, if there are 10 coins of Rs 2.

