

## System Of Particles & Rotational Motion

### Very Short Answer Type Questions

1. The centre of gravity of a body on the earth coincides with its centre of mass for a 'small' object whereas for an 'extended' object it may not. What is the qualitative meaning of 'small' and 'extended' in this regard?

For which of the following the two coincides? A building, a pond, a lake, a mountain?

2. Why does a solid sphere have smaller moment of inertia than a hollow cylinder of same mass and radius, about an axis passing through their axes of symmetry?
3. The variation of angular position  $\theta$ , of a point on a rotating rigid body, with time  $t$  is shown in Fig. 7.7. Is the body rotating clock-wise or anti-clockwise?




Fig. 7.7

4. A uniform cube of mass  $m$  and side  $a$  is placed on a frictionless horizontal surface. A vertical force  $F$  is applied to the edge as shown in Fig. 7.8. Match the following (most appropriate choice)




Fig. 7.8

- |                       |  |
|-----------------------|--|
| (a) $mg/4 < F < mg/2$ | (i) Cube will move up.                                       |
| (b) $F > mg/2$        | (ii) Cube will not exhibit motion.                           |
| (c) $F > mg$          | (iii) Cube will begin to rotate and slip at A.               |
| (d) $F = mg/4$        | (iv) Normal reaction effectively at $a/3$ from A, no motion. |

5. A uniform sphere of mass  $m$  and radius  $R$  is placed on a rough horizontal surface (Fig. 7.9). The sphere is struck horizontally at a height  $h$  from the floor. Match the following:




Fig. 7.9

- |                |   |
|----------------|---|
| (a) $h = R/2$  | (i) Sphere rolls without slipping with a constant velocity and no loss of energy. |
| (b) $h = R$    | (ii) Sphere spins clockwise, loses energy by friction.                            |
| (c) $h = 3R/2$ | (iii) Sphere spins anti-clockwise, loses energy by friction.                      |
| (d) $h = 7R/5$ | (iv) Sphere has only a translational motion, loses energy by friction.            |

## Short Answer Type Questions

- The vector sum of a system of non-collinear forces acting on a rigid body is given to be non-zero. If the vector sum of all the torques due to the system of forces about a certain point is found to be zero, does this mean that it is necessarily zero about any arbitrary point?
- A wheel in uniform motion about an axis passing through its centre and perpendicular to its plane is considered to be in mechanical (translational plus rotational) equilibrium because

no net external force or torque is required to sustain its motion. However, the particles that constitute the wheel do experience a centripetal acceleration directed towards the centre. How do you reconcile this fact with the wheel being in equilibrium?

3. How would you set a half-wheel into uniform motion about an axis passing through the centre of mass of the wheel and perpendicular to its plane? Will you require external forces to sustain the motion?
4. A door is hinged at one end and is free to rotate about a vertical axis (Fig. 7.10). Does its weight cause any torque about this axis? Give reason for your answer




Fig. 7.10

5.  $(n-1)$  equal point masses each of mass  $m$  are placed at the vertices of a regular  $n$ -polygon. The vacant vertex has a position vector  $a$  with respect to the centre of the polygon. Find the position vector of centre of mass

## Long Answer Type Questions

1. Find the centre of mass of a uniform (a) half-disc, (b) quarter-disc.
2. Two discs of moments of inertia  $I_1$  and  $I_2$  about their respective axes (normal to the disc and passing through the centre), and rotating with angular speed  $\omega_1$  and  $\omega_2$  are brought into contact face to face with their axes of rotation coincident.
  - (a) Does the law of conservation of angular momentum apply to the situation? why?
  - (b) Find the angular speed of the two-disc system.
  - (c) Calculate the loss in kinetic energy of the system in the process.
  - (d) Account for this loss.

3. A disc of radius  $R$  is rotating with an angular speed  $\omega_0$  about a horizontal axis. It is placed on a horizontal table. The coefficient of kinetic friction is  $\mu_k$ .
- (a) What was the velocity of its centre of mass before being brought in contact with the table?
  - (b) What happens to the linear velocity of a point on its rim when placed in contact with the table?
  - (c) What happens to the linear speed of the centre of mass when disc is placed in contact with the table?
  - (d) Which force is responsible for the effects in (b) and (c).
  - (e) What condition should be satisfied for rolling to begin?
  - (f) Calculate the time taken for the rolling to begin.
4. Two cylindrical hollow drums of radii  $R$  and  $2R$ , and of a common height  $h$ , are rotating with angular velocities  $\omega$  (anti-clockwise) and  $\omega$  (clockwise), respectively. Their axes, fixed are parallel and in a horizontal plane separated by  $(3R + \delta)$ . They are now brought in contact ( $\delta \rightarrow 0$ ).
- (a) Show the frictional forces just after contact.
  - (b) Identify forces and torques external to the system just after contact.
  - (c) What would be the ratio of final angular velocities when friction ceases?
5. A uniform square plate  $S$  (side  $c$ ) and a uniform rectangular plate  $R$  (sides  $b, a$ ) have identical areas and masses (Fig. 7.11).




Fig. 7.11

Show that

- (i)  $I_{xR} / I_{xS} < 1$ ;
- (ii)  $I_{yR} / I_{yS} > 1$ ;
- (iii)  $I_{zR} / I_{zS} > 1$ .

6. A uniform disc of radius  $R$ , is resting on a table on its rim. The coefficient of friction between disc and table is  $\mu$  (Fig 7.12). Now the disc is pulled with a force  $F$  as shown in the figure. What is the maximum value of  $F$  for which the disc rolls without slipping ?




Fig. 7.12