



# Mathematics

for

## GUJCET-ME - 2017

### Important Instructions :

1. The Mathematics test consists of 40 questions. Each question carries 1 mark. For each correct response, the candidate will get 1 mark. For each incorrect response,  $\frac{1}{4}$  mark will be deducted. The maximum marks are 40.
2. This Test is of **1 hour** duration.
3. Use **Black Ball Point Pen only** for writing particulars on OMR Answer Sheet and marking answers by darkening the circle '●'.
4. Rough work is to be done on the space provided for this purpose in the Test Booklet only.
5. **On completion of the test, the candidate must handover the Answer Sheet to the Invigilator in the Room / Hall. The candidates are allowed to take away this Test Booklet with them.**
6. Use of White fluid for correction is not permissible on the Answer Sheet.
7. Each candidate must show on demand his / her Admission Card to the Invigilator.
8. No candidate, without special permission of the Superintendent or Invigilator, should leave his / her seat.
9. Use of Manual Calculator is permissible.
10. The candidates are governed by all Rules and Regulations of the Board with regard to their conduct in the Examination Hall. All cases of unfair means will be dealt with as per Rules and Regulations of the Board.
11. No part of the Test Booklet and Answer Sheet shall be detached under any circumstances.
12. The candidates will write the Correct Test Booklet Set No. as given in the Test Booklet / Answer Sheet in the Attendance Sheet. (Patrak - 01)

Student Bro



## MATHEMATICS

1. If the length of the subnormal of a curve is constant and if it passes through the origin, then the equation of curve is \_\_\_\_\_.
- (A)  $x^2 + y^2 = k^2; k \in R$  (B)  $y^2 = kx; k \in R$   
 (C)  $x^2 = ky^2; k \in R$  (D)  $x^2 - y^2 = k^2; k \in R$

**Answer (B)**

**Sol.**  $y \frac{dy}{dx} = k_1$

$$\Rightarrow \int y dy = \int k_1 dx$$

$$\Rightarrow \frac{y^2}{2} = k_1 x + c$$

at  $x = 0, y = 0$

$$\therefore y^2 = 2k_1 x$$

$$y^2 = kx, k \in R$$

2. The integrating factor of the differential equation

$$\frac{dy}{dx} = \frac{1}{x+y+2} \text{ is}$$

- (A)  $e^{x+y+2}$  (B)  $e^y$   
 (C)  $e^{-y}$  (D)  $\log|x+y+2|$

**Answer (C)**

**Sol.**  $\frac{dx}{dy} = x+y+2$

$$\frac{dx}{dy} - x = y+2$$

$$\text{I.F.} = e^{\int -dy} = e^{-y}$$

3. If  $\vec{a} + \vec{b} + \vec{c} = \vec{0}$  and  $|\vec{a}| = 3, |\vec{b}| = 5, |\vec{c}| = 7$  and  $(\vec{a}, \vec{b}) = \alpha$ , then  $\alpha =$  \_\_\_\_\_.

- (A)  $\frac{2\pi}{3}$  (B)  $\frac{\pi}{6}$   
 (C)  $\frac{\pi}{3}$  (D)  $\frac{5\pi}{6}$

**Answer (C)**

**Sol.**  $|\vec{a} + \vec{b}|^2 = |\vec{c}|^2$

$$|\vec{a}|^2 + |\vec{b}|^2 + 2\vec{a} \cdot \vec{b} = |\vec{c}|^2$$

$$\cos \alpha = \frac{1}{2}$$

$$\alpha = \frac{\pi}{3}$$

4. For  $A(1, -2, 4), B(5, -1, 7), C(3, 6, -2), D(4, 5, -1)$ , the projection of  $\vec{AB}$  on  $\vec{CD}$  is \_\_\_\_\_.

- (A)  $(2\sqrt{3}, -2\sqrt{3}, 2\sqrt{3})$  (B)  $\frac{3}{13}(4, 1, 3)$   
 (C)  $(1, -1, 1)$  (D)  $(2, -2, 2)$

**Answer (D)**

**Sol.**  $\vec{AB} = (4, 1, 3)$

$$\vec{CD} = (1, -1, 1)$$

Now projection of  $\vec{AB}$  on  $\vec{CD} =$

$$\left( \frac{\vec{AB} \cdot \vec{CD}}{|\vec{CD}|} \right) \cdot \widehat{CD}$$

$$= \left( \frac{4-1+3}{\sqrt{3}} \right) \cdot \frac{(1, -1, 1)}{\sqrt{3}}$$

$$= (2, -2, 2)$$

5. The position vector of point A is  $(4, 2, -3)$ . If  $p_1$  is perpendicular distance of A from XY-plane and  $p_2$  is perpendicular distance from Y-axis, then  $p_1 + p_2 =$  \_\_\_\_\_.

- (A) 8  
 (B) 3  
 (C) 2  
 (D) 7

**Answer (A)**

**Sol.**  $p_1 = |z|$

$$p_2 = \sqrt{x_1^2 + y_1^2}$$

$$p_1 + p_2 = 5 + 3 = 8$$

6. Plane  $ax + by + cz = 1$  intersect axes in A, B, C respectively. If  $G\left(\frac{1}{6}, -\frac{1}{3}, 1\right)$  is a centroid of  $\Delta ABC$ , then  $a + b + 3c =$  \_\_\_\_\_.

- (A)  $\frac{4}{3}$  (B) 4  
 (C) 2 (D)  $\frac{5}{6}$

**Answer (C)**

**Sol.**  $A\left(\frac{1}{a}, 0, 0\right) B\left(0, \frac{1}{b}, 0\right) C\left(0, 0, \frac{1}{c}\right)$

centroid  $\Rightarrow \left(\frac{1}{3a}, \frac{1}{3b}, \frac{1}{3c}\right) = \left(\frac{1}{6}, \frac{-1}{3}, 1\right)$

$\therefore a = 2, b = -1, c = 1/3$

$\therefore a + b + 3c = 2$

Only (C) is correct

7. The direction angles of the line  $x = 4z + 3, y = 2 - 3z$  are  $\alpha, \beta$  and  $\gamma$ , then  $\cos\alpha + \cos\beta + \cos\gamma =$  \_\_\_\_\_.

(A)  $\frac{2}{\sqrt{26}}$

(B)  $\frac{8}{\sqrt{26}}$

(C) 1

(D) 2

**Answer (A)**

**Sol.**  $\frac{x-3}{4} = \frac{y-2}{-3} = \frac{z}{1}$

$\cos\alpha = \frac{4}{\sqrt{26}} \cos\beta = \frac{-3}{\sqrt{26}} \cos\gamma = \frac{1}{\sqrt{26}}$

$\therefore \cos\alpha + \cos\beta + \cos\gamma = \frac{2}{\sqrt{26}}$

8. If the normal of the plane makes an angles  $\frac{\pi}{4}, \frac{\pi}{4}$

and  $\frac{\pi}{2}$  with positive X-axis, Y-axis and Z-axis respectively and the length of the perpendicular line segment from origin to the plane is  $\sqrt{2}$ , then the equation of the plane is \_\_\_\_\_.

(A)  $x + y + z = \sqrt{2}$

(B)  $x + y + z = 1$

(C)  $x + y = 2$

(D)  $x = \sqrt{2}$

**Answer (C)**

**Sol.** Direction ratio of normal to the plane is

$\left(\cos\frac{\pi}{4}, \cos\frac{\pi}{4}, \cos\frac{\pi}{2}\right)$

$\left(\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, 0\right)$

$\therefore \frac{x}{\sqrt{2}} + \frac{y}{\sqrt{2}} = \sqrt{2}$

$x + y = 2$

9. If  $f : R - \left\{\frac{3}{5}\right\} \rightarrow R - \left\{\frac{3}{5}\right\}; f(x) = \frac{3x+1}{5x-3}$ , then \_\_\_\_\_.

(A)  $f^{-1}(x) = 2f(x)$

(B)  $f^{-1}(x) = f(x)$

(C)  $f^{-1}(x) = -f(x)$

(D)  $f^{-1}(x)$  doesnot exists

**Answer (B)**

**Sol.**  $f(x) = \frac{3x+1}{5x-3}$

we know that if

$f(x) = \frac{ax+b}{cx+d}$

if  $a + d = 0$

$f(x) = f^{-1}(x)$

10.  $f : R \rightarrow R, f(x) = 3x + 2$

$g : R \rightarrow R, g(x) = 6x + 5$

for the given functions  $(g \circ f^{-1})(10) =$  \_\_\_\_\_.

(A) 21

(B) 29

(C) 7

(D)  $\frac{8}{3}$

**Answer (A)**

**Sol.**  $g(f^{-1}(x))$

$= g\left(\frac{x-2}{3}\right)$

$= 6\left(\frac{x-2}{3}\right) + 5$

$= 2x + 1, \text{ at } x = 10$

$= 21$

11. Relation  $S = \{(1, 2), (2, 1), (2, 3)\}$  is defined on the set  $\{1, 2, 3\}$  is \_\_\_\_\_.

(A) not transitive

(B) symmetric

(C) reflexive

(D) equivalence

**Answer (A)**

**Sol.** As (1,1) is not there,

$\therefore$  it is not transitive

12.  $\tan^{-1}(\cot x) + \cot^{-1}(\tan x) =$  \_\_\_\_ (where,  $0 < x < \frac{\pi}{2}$ )

(A)  $\frac{\pi}{2}$

(B)  $2x$

(C)  $\pi - 2x$

(D)  $\pi - x$

**Answer (C)**

**Sol.**  $\frac{\pi}{2} - \cot^{-1}(\cot x) + \frac{\pi}{2} - \tan^{-1}(\tan x)$

$= \pi - 2x$

13.  $\cos\left(2\left(\tan^{-1}\frac{1}{5} + \tan^{-1}5\right)\right) = \underline{\hspace{2cm}}$ .

(A)  $\frac{1}{\sqrt{2}}$

(B) 0

(C) 1

(D) -1

**Answer (D)**

**Sol.**  $\tan^{-1}\frac{1}{5} + \tan^{-1}5 = \frac{\pi}{2}$

$\therefore \cos 2 \times \frac{\pi}{2} = \cos \pi = -1$

14. For  $\triangle ABC$  if  $A = \tan^{-1} 2$ ,  $B = \tan^{-1} 3$ , then  $C = \underline{\hspace{2cm}}$ .

(A)  $\frac{\pi}{6}$

(B)  $\frac{\pi}{4}$

(C)  $\frac{\pi}{3}$

(D)  $\frac{5\pi}{6}$

**Answer (B)**

**Sol.**  $A + B + C = \pi$

$A + B = \frac{3\pi}{4}$

$\therefore C = \frac{\pi}{4}$

15. If the area of the triangle with vertices  $(2, 5)$ ,  $(7, k)$  and  $(3, 1)$  is 10, then find the value of  $k$ .

(A) -5 or 35

(B) 5 or -35

(C) 15 or -5

(D) -5 or -25

**Answer (B)**

**Sol.**  $10 = \frac{1}{2} \begin{vmatrix} 2 & 5 \\ 7 & k \\ 3 & 1 \\ 2 & 5 \end{vmatrix}$

$k = 5$  or  $-35$

16. If  $k = p + q + r$ , then the value of

$\begin{vmatrix} k+r & p & q \\ r & k+p & q \\ r & p & k+q \end{vmatrix}$  is \_\_\_\_\_.

(A)  $2k^2$

(B)  $2k^3$

(C)  $k^3$

(D)  $3k^2$

**Answer (B)**

**Sol.**  $C_1 \rightarrow C_1 + C_2 + C_3$

and taking common

$k + p + q + r$

$= 2k \begin{vmatrix} 1 & p & q \\ 1 & k+p & q \\ 1 & p & k+q \end{vmatrix}$

$= R_3 \rightarrow R_3 - R_1$

$= R_2 \rightarrow R_2 - R_1$

$= 2k \begin{vmatrix} 1 & p & q \\ 0 & k & 0 \\ 0 & 0 & k \end{vmatrix}$

$= 2k^3$

17. If maximum and minimum values of

$D = \begin{vmatrix} 1 & -\cos\theta & -1 \\ \cos\theta & 1 & -\cos\theta \\ 1 & \cos\theta & 1 \end{vmatrix}$  are  $p$  and  $q$

respectively, then the value of  $2p + 3q$  is \_\_\_\_\_.

(A) 16

(B) 6

(C) 14

(D) 8

**Answer (C)**

**Sol.**  $D = 1(1 + \cos^2\theta) + \cos\theta(\cos\theta + \cos\theta) - 1(\cos^2\theta - 1)$

$D = 1 + \cos^2\theta + 2\cos^2\theta - \cos^2\theta + 1$

$= 2(1 + \cos^2\theta)$

$\therefore p = 4, q = 2$

$\therefore 2p + 3q = 14$

18. If  $A = \begin{bmatrix} 1 & 4 & 4 \\ 4 & 1 & 4 \\ 4 & 4 & 1 \end{bmatrix}$ , then  $A^2 - 6A = \underline{\hspace{2cm}}$ .

(A)  $27 I_3$

(B)  $5 I_3$

(C)  $20 I_3$

(D)  $30 I_3$

**Answer (A)**

**Sol.**  $\begin{bmatrix} 1 & 4 & 4 \\ 4 & 1 & 4 \\ 4 & 4 & 1 \end{bmatrix} \Rightarrow A^2 - 6A = 27 I_3$



19. If  $\begin{bmatrix} 1 & x & 3 \\ 2 & 4 & 5 \\ 3 & 2 & x \end{bmatrix} \begin{bmatrix} x \\ 2 \\ 0 \end{bmatrix} = 0$ , then  $x$  \_\_\_\_\_.

- (A)  $\frac{7}{3}$  (B)  $\frac{5}{3}$   
 (C)  $-\frac{5}{3}$  (D)  $-\frac{7}{3}$

**Answer (C)**

**Sol.**  $\begin{bmatrix} 20 & 20 + 2x & 21 + 4x \end{bmatrix} \begin{bmatrix} x \\ 2 \\ 0 \end{bmatrix} = 0$

$\Rightarrow 24x + 40 = 0$

$\Rightarrow x = -\frac{5}{3}$

20.  $\frac{d}{dx} \left( \sqrt{3} \sin \left( 2x + \frac{\pi}{3} \right) + \cos \left( 2x + \frac{\pi}{3} \right) \right) =$  \_\_\_\_\_.

- (A)  $4 \cos 2x$  (B)  $-4 \sin 2x$   
 (C)  $4 \sin 2x$  (D)  $-4 \cos 2x$

**Answer (B)**

**Sol.**  $\frac{d}{dx} \left( 2 \sin \left( 2x + \frac{\pi}{3} + \frac{\pi}{6} \right) \right)$

$\frac{d}{dx} \left( 2 \sin \left( 2x + \frac{\pi}{2} \right) \right)$

$\frac{d}{dx} (2 \cos 2x)$

$\Rightarrow -4 \sin 2x$

21. For the curve  $f(x) = (x - 5)^2$ , applying mean value theorem on  $[4, 6]$  the tangent at \_\_\_\_\_ is parallel to the chord joining  $A(4, 1)$ ,  $B(6, 1)$ .

- (A)  $(4, 6)$  (B)  $\left( \frac{9}{2}, \frac{1}{4} \right)$   
 (C)  $(0, 5)$  (D)  $(5, 0)$

**Answer (D)**

**Sol.**  $f'(c) = 2(c - 5) = \frac{f(6) - f(4)}{6 - 4} = 0$

(using LMVT)

$\therefore c = 5$

$\therefore (5, 0)$

22. Function  $f(x) = \begin{cases} (\log_2 2x)^{\log_x 8}; & x \neq 1 \\ (k-1)^3; & x = 1 \end{cases}$  is continuous at  $x = 1$ , then  $k =$  \_\_\_\_\_

- (A)  $e + 1$  (B)  $e^{1/3}$   
 (C)  $e^3$  (D)  $e - 1$

**Answer (A)**

**Sol.**  $f(1^-) = f(1^+) = f(1)$

$\lim_{x \rightarrow 1} e^{(\log_2 2x - 1) \log_x 8}$

$\lim_{x \rightarrow 1} e^{(\log_2 2x - 1) \frac{\log_2 8}{\log_2 x}}$

$e^3 = (k - 1)^3$

$k = e + 1$

23.  $\int \frac{dx}{\cos x \sqrt{1 + \cos 2x + \sin 2x}} =$  \_\_\_\_\_ + C;  $\left( 0 < x < \frac{\pi}{4} \right)$

- (A)  $2 + \sqrt{\cot x}$  (B)  $\sqrt{\tan x + 1}$   
 (C)  $\sqrt{2 + 2 \tan x}$  (D)  $\sqrt{2 + 2 \cot x}$

**Answer (C)**

**Sol.**  $\int \frac{\sec^2 x dx}{\sqrt{2 + 2 \tan x}}$

$\tan x = t$

$\sec^2 x dx = dt$

$\therefore \sqrt{2 + 2 \tan x} + C$

24. If  $\int \frac{\sin 2x}{\sin 5x \sin 3x} dx = \frac{1}{3} \log |\sin 3x| - \frac{1}{5} \log |f(x)| + C$ ,

then  $f(x) =$  \_\_\_\_\_.

- (A)  $\sin 5x$  (B)  $\sin 4x$   
 (C)  $\sin 2x$  (D)  $\sin 6x$

**Answer (A)**

**Sol.**  $\int \frac{\sin(5x - 3x)}{\sin 5x \sin 3x} dx$

$\int \frac{\sin 5x \cos 3x - \cos 5x \sin 3x}{\sin 5x \sin 3x} dx$

$\int (\cot 3x - \cot 5x) dx$

$\therefore \frac{1}{3} \log |\sin 3x| - \frac{1}{5} \log |\sin 5x| + C$

$\Sigma P(x_i) = 1$

**FREE!**




**JEE & NEET**

**GUJCET**

ગુજરાતી -- English -- हिन्दी


- PYQ Mock Test
- Chapter & Topic-Wise PYQs (Updated 2026)
- NCERT Line by Line MCQs.
- AR, Statement, Diagram Based Questions.
- Detailed Step-by-Step Solutions



# Exam Bro

CLICK HERE!

Download Now



25. The probability distribution of a discrete random variable  $X$  is given in the following table :

|         |        |              |          |
|---------|--------|--------------|----------|
| $X = x$ | 0      | 1            | 2        |
| $P(x)$  | $4C^3$ | $4C - 13C^2$ | $7C - 1$ |

;  $C > 0$  then  $C =$  \_\_\_\_.

- (A) 2  
 (B) 1  
 (C)  $\frac{1}{4}$   
 (D) 1 and  $-\frac{1}{4}$

**Answer (C)**

**Sol.**  $4C^3 + 4C - 13C^2 + 7C - 1 = 1$

$\therefore C = \frac{1}{4}$

26. The probability function of a binomial distribution is

$P(x) = \binom{6}{x} p^x q^{6-x}$ ,  $x = 0, 1, 2, \dots, 6$ . If  $2P(2) = 3P(3)$ , then  $p =$  \_\_\_\_\_.

- (A)  $\frac{1}{3}$  (B)  $\frac{1}{4}$   
 (C)  $\frac{1}{2}$  (D)  $\frac{1}{5}$

**Answer (A)**

**Sol.**  $2P(2) = 3P(3)$

$2 \times {}^6C_2 p^2 q^4 = 3 \times {}^6C_3 p^3 q^3$

On solving  $p = \frac{1}{3}$

(as  $p + q = 1$ )

27. The mean and standard deviation of random variable

$X$  are 10 and 5 respectively, then  $E\left(\frac{X-15}{5}\right)^2 =$

- (A) 4 (B) 3  
 (C) 2 (D) 5

**Answer (C)**

**Sol.**  $v = 5^2 = 25$

$\therefore E(x)^2 = 25 + 10^2 = 125$

$\therefore E\left(\frac{x^2 + 225 - 30x}{25}\right) = \frac{1}{25}(E(x)^2 + 225 - 30E(x))$

$= 2$

28. The corner points of the feasible region determined by the system of linear constraints are  $(0, 10)$ ,  $(5, 5)$ ,  $(25, 20)$  and  $(0, 30)$ . Let  $Z = px + qy$ , where  $p, q > 0$ . Condition on  $p$  and  $q$  so that the maximum of  $Z$  occurs at both the points  $(25, 20)$  and  $(0, 30)$  is \_\_\_\_\_.

- (A)  $5p = 2q$  (B)  $2p = 5q$   
 (C)  $p = 2q$  (D)  $q = 3p$

**Answer (A)**

**Sol.**  $25p + 20q = 30q$

$5p = 2q$

29. Two curves  $a_i x^2 + b_i y^2 = 1$ ;  $i = 1, 2$  where,  $a_1 \neq a_2$ ,  $b_1 \neq b_2$ ,  $a_1, a_2, b_1, b_2 \neq 0$  may intersect orthogonally if \_\_\_\_\_.

- (A)  $a_1 a_2 = b_1 b_2$   
 (B)  $a_1^{-1} - a_2^{-1} = b_1^{-1} - b_2^{-1}$   
 (C)  $a_1^{-1} + a_2^{-1} = b_1^{-1} + b_2^{-1}$   
 (D)  $a_1 b_2 = a_2 b_1$

**Answer (B)**

**Sol.** If  $a_1 x^2 + b_1 y^2 = 1$  and  $a_2 x^2 + b_2 y^2 = 1$  are orthogonal to each other

$\frac{1}{a_1} - \frac{1}{a_2} = \frac{1}{b_1} - \frac{1}{b_2}$

30. The sides of two squares are  $x$  and  $y$  respectively, such that  $y = x + x^2$ . The rate of change of area of second square with respect to area of first square is \_\_\_\_\_.

- (A)  $x^2 + 3x - 1$  (B)  $2x^2 - 3x + 1$   
 (C)  $2x^2 + 3x + 1$  (D)  $1 + 2x$

**Answer (C)**

**Sol.**  $\frac{dA_2}{dA_1} = \frac{2y \frac{dy}{dt}}{2x \frac{dx}{dt}}$

$\frac{y}{x} \frac{dy}{dx}$

$\therefore \frac{y}{x} \times (1 + 2x)$

$2x^2 + 3x + 1$

31. The approximate value of  $\sqrt[3]{-0.99}$  is \_\_\_\_\_.

- (A)  $-0.9967$  (B)  $-0.9976$   
 (C)  $-1.0033$  (D)  $-1$

**Answer (A)**

**Sol.** - .9967

32.  $\int \frac{dx}{\sqrt{x^{10} - x^2}}$ ;  $x > 1 = \underline{\hspace{2cm}} + C.$

- (A)  $\frac{1}{4} \log|\sqrt{x^{10} - x^2} + x^2|$  (B)  $\frac{1}{2} \log|x^{10} - x^2|$   
(C)  $-\frac{1}{4} \sec^{-1}(x^4)$  (D)  $\frac{1}{4} \sec^{-1}(x^4)$

**Answer (D)**

**Sol.**  $\int \frac{4x^3 dx}{4x^3 \times x \sqrt{x^8 - 1}}$

$\int \frac{4x^3 dx}{4x^4 \sqrt{x^8 - 1}}$

$\left( \begin{array}{l} x^4 = t \\ 4x^3 dx = dt \end{array} \right)$

$\int \frac{dt}{4t\sqrt{t^2 - 1}} = \frac{1}{4} \sec^{-1}(x^4) + C$

33.  $\int e^{\sin x} (x \cos x - \sec x \tan x) dx = \underline{\hspace{2cm}} + C;$

$0 < x < \frac{\pi}{2}.$

- (A)  $e^{\sin x}(x - \sec x)$  (B)  $e^{\sin x}(\sec x - x)$   
(C)  $e^{\sin x}x \cos x$  (D)  $e^{\sin x}(x + \sec x)$

**Answer (A)**

**Sol.**

$\int x e^{\sin x} \cos x - \int e^{\sin x} dx - \left[ (\sec x \cdot e^{\sin x}) - \int e^{\sin x} dx \right]$   
 $e^{\sin x} (x - \sec x) + C$

34.  $\int \sin(11x) \cdot \sin^9 x dx = \underline{\hspace{2cm}} + C.$

- (A)  $\frac{\sin(10x) \cdot \sin^{10} x}{10}$  (B)  $\frac{\sin^{11} x}{11}$   
(C)  $\frac{\sin(9x) \cdot \sin^9 x}{9}$  (D)  $\frac{\cos(10x) \cdot \cos^{10} x}{10}$

**Answer (A)**

**Sol.**  $\int \sin(10x + x) \sin^9 x dx$

$\int \sin 10x \cos x \sin^9 x dx + \int \cos 10x \sin^{10} x dx$

$\frac{\sin^{10} x}{10} \sin 10x - \frac{10}{10} \int \sin^{10} x \cos 10x dx + \int \cos(10x) \sin^{10} x dx$

$\frac{\sin^{10} x}{10} \sin(10x) + c$

35.  $\int_{-\log 3}^{\log 3} \cot^{-1} \left( \frac{e^x - 1}{e^x + 1} \right) dx = \underline{\hspace{2cm}}$

- (A)  $\frac{\pi}{2} \log 3$  (B)  $\pi \log 3$   
(C) 0 (D)  $\pi \log 9$

**Answer (B)**

**Sol.**  $I = \int_{-\log 3}^{\log 3} \cot^{-1} \left( \frac{e^x - 1}{e^x + 1} \right) dx = \int_{-\log 3}^{\log 3} \cot^{-1} \left( \frac{1 - e^x}{1 + e^x} \right) dx$

adding

$2I = \pi \int_{-\log 3}^{\log 3} dx$

$I = \pi \log 3$

36.  $\int_0^{100\pi} |\cos x| dx = \underline{\hspace{2cm}}.$

- (A) 200 (B) 100  
(C) 50 (D) 0

**Answer (A)**

**Sol.**  $200 \int_0^{\pi/2} \cos x dx = 200$

37.  $\int_0^{\pi/2} (x - [\sin x]) dx = \underline{\hspace{2cm}}.$

(where  $[x]$  = greatest integer not greater than  $x$ ).

- (A)  $\frac{\pi^2}{8} - 2$  (B)  $\frac{\pi^2}{4} - 1$   
(C)  $\frac{\pi^2}{8} - 1$  (D)  $\frac{\pi^2}{8}$

**Answer (D)**

**Sol.**  $\int_0^{\pi/2} x dx$

$\frac{\pi^2}{8}$

38. Area bounded between two latus-rectum of the

ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ ;  $a > b$  is  $\underline{\hspace{2cm}}.$

(where,  $e$  is eccentricity of the ellipse)

- (A)  $2b(be + a \sin^{-1} e)$  (B)  $8b(be + a \sin^{-1} e)$   
(C)  $b(be + a \sin^{-1} e)$  (D)  $4b(be + a \sin^{-1} e)$

**Answer (A)**

**Sol.**  $2b(be + a \sin^{-1} e)$

39. The area of the region bounded by the curves  $f(x) = \sin \pi x$  and X-axis is \_\_\_\_\_ where  $x \in [-1, 2]$ .

(A)  $8 \pi$

(B)  $\frac{8}{\pi}$

(C)  $\frac{6}{\pi}$

(D)  $6 \pi$

**Answer (C)**

**Sol.**  $3 \int_0^1 \sin(\pi x) dx$   
 $\frac{6}{\pi}$

40. The order and degree of the differential equation

$$\left(\frac{d^2y}{dx^2}\right)^3 + 3\frac{dy}{dx} = \sqrt{x}; x > 0 \text{ are _____ respectively.}$$

(A) 2 and 6

(B) 3 and 2

(C) 2 and 3

(D) 2 and degree is undefined

**Answer (C)**

**Sol.** order  $\Rightarrow$  2

degree  $\Rightarrow$  3



**Sb**  
**Student Bro**



**FREE!**




**JEE & NEET**

**GUJCET**

ગુજરાતી -- English -- हिन्दी

- PYQ Mock Test
- Chapter & Topic-Wise PYQs (Updated 2026)
- NCERT Line by Line MCQs.
- AR, Statement, Diagram Based Questions.
- Detailed Step-by-Step Solutions



# Exam Bro

CLICK HERE!

Download Now

